

AQUECA Report

Module-A. Disinfection efficacy of AQUAECA DESY.

Objective-1. Disinfection efficacy at 13A, pH-6.7-6.9

Material and Methods

A. Microorganisms used in the study

- Bacteria
 - *Staphylococcus aureus* (MTCC 9886)
 - *Enterococcus faecalis* (MTCC 439)
 - *Escherichia coli* (MTCC K12)
 - *Pseudomonas aeruginosa* (MTCC 424)
- Spores
 - *Bacillus subtilis* (MTCC 441)
- Fungi
 - *Candida albicans* (SC5314)
 - *Candida glabrata* (CBS138)

B. Sample preparation

1. Bacterial culture were grown in LB broth.
2. Fungal culture were grown in YPD (Yeast Extract-1%, Dextrose-2% and Peptone- 2%).
3. *Bacillus subtilis* sporulation was performed in sporulation media (SM). Composition of SM used was- Nutrient broth- 0.8%, KCl- 0.1%, MgSO₄·7H₂O- 0.012%, pH adjusted to 7.6 with 1 M NaOH, volume adjusted to 1 litre with ddH₂O. Autoclaved and allowed to cool to 50°C. Just prior to use, following sterile solutions were added: Ca(NO₃)₂- 1mM, MnCl₂- 0.01mM, FeSO₄-0.001mM. 0.5 ml of an overnight grown *B subtilis* culture was inoculated in liquid DSM media in Erlenmeyer flasks. Flasks were kept at 37 °C in an orbital shaker at 200 rpm for 7 days. Generation of spores was monitored microscopically. After 7 days of incubation > 90% of spore population was observed.
4. All cultures were maintained at 37°C.
5. Viable cell density of at least 10¹⁰ cells was maintained for bacteria and fungus.

C. Methodology

1. 1ml aliquot of each culture harbouring 10^{10} cells (as determined by total aerobic plate count or spectrophotometrically at 600 nm wavelength) was transferred into microfuge tube and centrifuged at 10,000 rpm for 5 min. Supernatant was discarded and pellet was stored for further use.
2. For calculating the disinfection efficacy (Log reduction) total aerobic plate count procedure was followed.
3. For determining the disinfection efficacy of DESY, pellet obtained from step-1 was resuspended in 1ml of Desy obtained from AQUAECA at 13A. 100 μ l of the resuspended pellet was removed after 1min, 2min, and 5 min exposure times and spread plated on relevant plates to calculate the log reduction (**Fig.1**).
4. As control the pellet was resuspende in sterile deionised water and 100 μ l of the resuspended pellet was removed after 5 min exposure times and spread plated on relevant plates. (**Fig.1**).
5. For statistical significance disinfection efficacy of **DESY** for each time point was determined using three biological replicates for each organism.

Results

Disinfection efficacy of Bacteria:

1. The bacterial disinfection efficacy of DESY obtained from AQUAECAat 13 A was tested using representative gram negative (*Escherichia coli* K12 and *Pseudomonas aeruginosa*) and gram positive bacteria (*Staphylococcus aureus* and *Enterococcus feacalis*).
2. Bacteria was exposed to DESY at room temperature. It was observed that both gram negative and gram positive bacteria could be efficiently destroyed by DESY even at 1 min exposure time (**Fig. 1**). No bacterial growth was observed at 2min and 5min time points as well (**Fig.1**).
3. Total aerobic plate count showed that 10 log reduction in bacterial counts was achieved via DESY treatment at 1min exposure time.

Disinfection efficacy of Fungi:

1. Fungal disinfection efficacy of DESY was tested using representative pathogenic fungi *Candida albicans* and *Candida glabrata*.

2. Fungi was exposed to DESY at room temperature. It was observed that both *Candida* species could be efficiently destroyed by DESY even at 1min exposure times (**Fig. 1**). No fungal growth was observed at 2min and 5min exposure time points as well (**Fig.1**)
3. Total aerobic plate count showed that 10 log reduction in fungal counts was achieved via DESY treatment at a exposure time of 1 min).

Disinfection efficacy of Spores:

1. Highly resistant bacteria spore disinfection efficacy of DESY was also analysed using representative *Bacillus subtilis* spores.
2. 10^6 spores were exposed to DESY at room temperature. It was observed that bacterial spores could be efficiently disinfected by DESY even at 1min exposure time (6 log reduction) (**Fig. 2**).
3. Total aerobic plate count method was used for determine disinfection efficacy.

Conclusions

A 10 log disinfection efficacy of bacterial culture was achieved via DESY treatment at room temperature with an exposure time of 1 min. DESY treatment was also effective on fungal cells, where a 10 log disinfection efficacy was observed at room temperature with an exposure time of 1 min. A 6 log disinfection efficacy of representative highly resistant *Bacillus subtilis* spores was also achieved with DESY at room temperature with an exposure time of 1 min.

**Dr Mohan Kamthan
Assistant Professor
Department of Biochemistry
School of Chemical and Lifesciences
Jamia Hamdard**




Figure 1. Disinfection efficacy of DESY at 13A. (A) Representative plates for total aerobic plate count. **(B)** Log reduction in bacteria and Fungi upon DESY treatment. **(C)** Relation between log reduction and % reduction.




Figure 2. Disinfection efficacy of DESY at 13A against *Bacillus subtilis* spores. (A) Representative plates for total aerobic plate count. **(B)** Relation between log reduction and % reduction.