

ASSIGNMENT 4

MANIKANTA VALLEPU - AI20BTECH11014

Download all python codes from

https://github.com/manik2255/AI1103-PROBABILITY-AND-RANDOM-VARIABLES/blob/main/ASSIGNMENT_4/assignment_4.py

and latex-tikz codes from

https://github.com/manik2255/AI1103-PROBABILITY-AND-RANDOM-VARIABLES/blob/main/ASSIGNMENT_4/ASSIGNMENT_4.tex

1 GATE 2017 MA PROBLEM.49

Let X and Y be independent and identically distributed random variables with probability density function $f(x) = \begin{cases} e^{-x} & x > 0 \\ 0 & \text{otherwise} \end{cases}$. Then $\Pr(\max(X, Y) < 2)$ equals

2 SOLUTION

Given,

$$f(x) = \begin{cases} e^{-x} & x > 0 \\ 0 & \text{otherwise} \end{cases} \quad (2.0.1)$$

We know,

$$\Pr(X < x) = \int_0^x f(x) dx \quad (2.0.2)$$

Using (2.0.1) in (2.0.2),

$$\Pr(X < x) = \int_0^x e^{-x} dx \quad (2.0.3)$$

Let $F_X(x)$ be the cumulative distribution function of random variable X .

$$F_X(x) = \int_0^x f(x) dx \quad (2.0.4)$$

$F_X(x)$ can be obtained from the uniform distribution of a random variable U on $(0,1)$ and let $U = e^{-x}$.

$$0 < U < 1 \quad (2.0.5)$$

As for random variable X also,

$$0 < F_X(x) < 1 \quad (2.0.6)$$

This similarity between U and $F_X(x)$ is used to generate the random variable X from U .

$$F_X(x) = \Pr(X < x) \quad (2.0.7)$$

$$= \Pr(-\log_e U < x) \quad (2.0.8)$$

$$= \Pr(U > e^{-x}) \quad (2.0.9)$$

$$= 1 - \Pr(U < e^{-x}) \quad (2.0.10)$$

$$= 1 - F_U(e^{-x}) \quad (2.0.11)$$

From uniform distribution,

$$F_U(x) = x, 0 < x < 1 \quad (2.0.12)$$

Using (2.0.11) in (2.0.12),




Fig. 1: CDF of random variable X

$$F_X(x) = \Pr(X < x) \quad (2.0.13)$$

$$= 1 - e^{-x} \quad (2.0.14)$$

$$\Pr(\max(X, Y) < 2) = \Pr(X < 2, Y < 2) \quad (2.0.15)$$

As X and Y are independent random variables,

$$\Pr(\max(X, Y) < 2) = \Pr(X < 2) \Pr(Y < 2) \quad (2.0.16)$$

using (2.0.14),

$$\Pr(\max(X, Y) < 2) = (1 - e^{-2})(1 - e^{-2}) \quad (2.0.17)$$

$$= 0.748. \quad (2.0.18)$$