Glucose Monitoring Report

Basic Information

Name: krishnan R

Type of Diabetes: Other

Device SN Code: 7D9043CCB543A

Sex: --

Age: --

Device Model: i3

Glucose Overview

7Days:03/23/2026 ~ 03/30/2026

Percentage of time used: 90.8%

Glucose Metrics

Mean Glucose (MG)[1] Goal: <154mg/dL	191 mg/dL
Glucose Management Indicator (GMI)[2] Goal: <7%	7.9 %
Glucose Variability (CV)[3] Goal: ≤36%	31.5 %

Time in Ranges[4]



Ambulatory Glucose Profile (AGP)



Low Glucose Condition

Hypoglycemia event(<70mg/dL)[5] **1** times Average Duration **33** minute(s)

Serious hypoglycaemia event(<54mg/dL)[6] **0** times



High Glucose Condition

Hyperglycaemia Event (>180mg/dL)[7] **20** time(s) Average Duration **275** minute(s)

Serious Hyperglycaemia Event (>250mg/dL)[8] **10** times Average Duration **166** minute(s)



Multi-day Glucose Details



Date	03/23	03/24	03/25	03/26	03/27	03/28	03/29	03/30								
Number of glucose value	69	480	480	480	480	480	480	165								
Highest mg/dL	257	294	278	353	352	336	234	303								
Lowest mg/dL	109	42	68	120	154	88	85	187								
MG mg/dL	191	179	168	199	237	186	158	236								
TIR	44.9%	43.4%	57.6%	44.1%	12.5%	58.2%	66.5%	0%								
TAR	55.1%	55%	41.6%	55.9%	87.5%	41.8%	33.5%	100%								
TBR	0%	1.6%	0.8%	0%	0%	0%	0%	0%								
LAGE[9] mg/dL	148	252	210	233	198	248	149	116								
MAGE[10] mg/dL	--*	82	81	78	81	0	62	--*								
MODD[11] mg/dL	--*	--*	64	55	56	82	58	--*								
SDBG[12] mg/dL	38	59	47	57	47	74	40	30								
CV	19.8%	33.2%	28.2%	28.6%	20%	39.4%	25.1%	12.8%								

* The data is less than 24 h and cannot be calculated.

Daily Glucose Profiles

Fingerstick blood · Exercise · Diet · Medication · Insulin · Multi Log · Other

TAR · TIR · TBR



MG	Highest	Lowest
191mg/dL	257mg/dL	109mg/dL

TAR	TIR	TBR
55.1%	44.9%	--%
1h54min	1h33min	0min

Time	Health Log	Details	Glucose (mg/dL)
22:40	Diet	chappati 3 Pre-meal glucose: 244 mg/dL 2h-PG: 132 mg/dL	237



MG	Highest	Lowest
179mg/dL	294mg/dL	42mg/dL

TAR	TIR	TBR
55%	43.4%	1.6%
13h12min	10h24min	24min

Time	Health Log	Details	Glucose (mg/dL)
10:46	Diet	thinai arisi pongal Pre-meal glucose: 236 mg/dL 2h-PG: 246 mg/dL	233



MG	Highest	Lowest
168mg/dL	278mg/dL	68mg/dL

TAR	TIR	TBR
41.6%	57.6%	0.8%
10h	13h48min	12min



MG	Highest	Lowest
199mg/dL	353mg/dL	120mg/dL

TAR	TIR	TBR
55.9%	44.1%	--%
13h24min	10h36min	0min



MG	Highest	Lowest
237mg/dL	352mg/dL	154mg/dL

TAR	TIR	TBR
87.5%	12.5%	--%
21h	3h	0min



MG	Highest	Lowest
186mg/dL	**336**mg/dL	**88**mg/dL
TAR	TIR	TBR
41.8%	**58.2**%	**--**%
10h3min	13h57min	0min



MG	Highest	Lowest
158mg/dL	**234**mg/dL	**85**mg/dL
TAR	TIR	TBR
33.5%	**66.5**%	**--**%
8h3min	15h57min	0min



MG	Highest	Lowest
236mg/dL	**303**mg/dL	**187**mg/dL
TAR	TIR	TBR
100%	**--**%	**--**%
8h15min	0min	0min

Explanation of metrics

1. Mean Glucose (MG): Refers to the average of all glucose measurements during the monitoring period to evaluate the overall glucose level.
2. Glucose Management Indicator (GMI): GMI is derived from at least 15 days of CGM data.The GMI may be similar to, higher than, or lower than the laboratory HbA1c.
3. Glucose Variability (CV): The degree of dispersion of all glucose measurements during the monitoring period is more suitable for comparing glucose variability in different populations and for evaluating glucose fluctuations.
4. Time in Ranges (TIR,TAR,TBR): Refers to the percentage of time during the monitoring period when glucose was within the target range, above the target range (distinguishing between high glucose and very high glucose), and below the target range (distinguishing between low glucose and very low glucose), respectively, and is used to evaluate the level of glycaemic control.
5. Hypoglycemia Event: Event starts when the glucose value is lower than the low limit of the target range (e.g., 3.9 mmol/L or 70 mg/dL for type 1 diabetes) for at least 15 minutes and ends when the glucose value is greater than or equal to the low limit of the target range for at least 15 minutes.
6. Serious Hypoglycaemia Event: An event starts when the glucose value remains below the hypoglycemia threshold (or TBR2 boundary value, e.g., 3.0 mmol/L or 54 mg/dL for type 1 diabetes) for at least 15 minutes, and ends when the glucose value stays at or above the hypoglycemia threshold (or TBR2 boundary value) for at least 15 minutes.
7. Hyperglycaemia Event: Event starts when the glucose value remains above the high limit of the target range (e.g., 10.0 mmol/L or 180 mg/dL for type 1 diabetes) for at least 15 minutes, and ends when the glucose value stays at or below the high limit of the target range for at least 15 minutes.
8. Serious Hyperglycaemia Event: Event starts when the glucose value remains above the high glucose threshold (or TAR2 boundary value, e.g., 13.9 mmol/L or 250 mg/dL for type 1 diabetes) for at least 15 minutes, and ends when the glucose value stays at or below the high glucose threshold (or TAR2 boundary value) for at least 15 minutes.
9. Largest Amplitude of Glycemic Excursion(LAGE): Refers to the difference between the maximum and minimum glucose values during the monitoring period and evaluates the maximum glucose fluctuation.
10. Mean Amplitude of Glycemic Excursion(MAGE): After removing all glucose fluctuations whose amplitude did not exceed a certain threshold (SDBG), the average value obtained by calculating the fluctuation amplitude based on the direction of the first valid fluctuation was used to evaluate the degree of intraday glucose fluctuation.
11. Mean of Daily Differences(MODD): It refers to the average of the absolute value of the difference between the relative measurements subtracted from each other over 2 consecutive days (i.e., a complete 48-hour period), reflecting the reproducibility of glucose from one day to another, and is used to evaluate the degree of fluctuation of glucose from day to day; MODD on the corresponding date reflects the degree of fluctuation of glucose values of the same day and the previous day.
12. Standard Deviation of Blood Glucose(SDBG): Refers to the standard deviation of measured glucose values during the monitoring date and evaluates the degree of overall deviation from the mean glucose value.
13. Ambulatory Glucose Profile (AGP): The AGP graph reflects your glucose fluctuations. The 50% median line shows the median of all glucose values at each time point — the smoother the curve, the more stable the glucose level. The 25%–75% shaded area covers 50% of glucose readings, and the 5%–95% shaded area covers 90% of glucose readings, indicating daily glucose fluctuations. A wider band represents greater fluctuations, while a narrower band indicates more stable glucose levels.

References

[1] Clinical Targets for Continuous Glucose Monitoring Data Interpretation: Recommendations From the International Consensus on Time in Rang [J]. Diabetes Care 2019;42(8):1593–1603. https://doi.org/10.2337/dci19-0028

[2] Glycemic Targets: Standards of Care in Diabetes—2023 [J].Diabetes Care 2023;46(Supplement_1):S97–S110.https://doi.org/10.2337/dc23-S006

[3] International Diabetes Center captūrAGP® Version 5.0 [J] Final 3.11.2023

[4] Continuous glucose monitoring and metrics for clinical trials: an international consensus statement [J].Lancet Diabetes Endocrinol 2023; 11: 42–57. https://doi.org/10.1016/S2213-8587(22)00319-9