

Gate Questions

EE24BTECH11013-DASARI MANIKANTA

- 1) The energy levels of a particle of mass m in a potential of the form

$$V(x) = \begin{cases} \infty, & x \leq 0 \\ \frac{1}{2}m\omega^2 x^2, & x > 0 \end{cases}$$

are given, in terms of quantum number $n = 0, 1, 2, 3, \dots$, by

- a) $\left(n + \frac{1}{2}\right)\hbar\omega$
 - b) $\left(2n + \frac{1}{2}\right)\hbar\omega$
 - c) $\left(2n + \frac{3}{2}\right)\hbar\omega$
 - d) $\left(n + \frac{3}{2}\right)\hbar\omega$
- 2) The electromagnetic field due to a point charge must be described by Lienard-Weichert potentials when
- a) the point charge is highly accelerated,
 - b) the electric and magnetic fields are not perpendicular.
 - c) the point charge is moving with velocity close to that of light.
 - d) the calculation is done for the radiation zone, i.e far away from the charge.
- 3) The strangeness quantum number is conserved in
- a) strong, weak and electromagnetic interactions .
 - b) weak and electromagnetic interactions only.
 - c) strong and weak interactions only.
 - d) strong and electromagnetic interactions only.
- 4) The eigenvalues and eigenvectors of the matrix

$$\begin{bmatrix} 5 & 4 \\ 1 & 2 \end{bmatrix}$$

are

- a) 6, 1 and $\begin{bmatrix} 4 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \end{bmatrix}$
 - b) 2, 5 and $\begin{bmatrix} 4 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \end{bmatrix}$
 - c) 6, 1 and $\begin{bmatrix} 1 \\ 4 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \end{bmatrix}$
 - d) 2, 5 and $\begin{bmatrix} 1 \\ 4 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \end{bmatrix}$
- 5) A vector field is defined everywhere as $\mathbf{F} = -\frac{y^2}{L}\hat{i} + z\hat{k}$. The net flux of \mathbf{F} associated with a cube of side L , with one vertex at the origin and sides along the positive X , Y , and Z axes, is
- a) $2L^3$
 - b) $4L^3$

- c) $8L^3$
 d) $10L^3$
- 6) If $\mathbf{r} = x\hat{i} + y\hat{j}$, then
 a) $\nabla \cdot \mathbf{r} = 0$ and $\nabla|\mathbf{r}| = \frac{\mathbf{r}}{r}$
 b) $\nabla \cdot \mathbf{r} = 2$ and $\nabla|\mathbf{r}| = \hat{r}$
 c) $\nabla \cdot \mathbf{r} = 2$ and $\nabla|\mathbf{r}| = \frac{\mathbf{r}}{r}$
 d) $\nabla \cdot \mathbf{r} = 3$ and $\nabla|\mathbf{r}| = \frac{\mathbf{r}}{r}$
- 7) Consider a vector $\mathbf{p} = 2\hat{i} + 3\hat{j} + 2\hat{k}$ in the coordinate system $(\hat{i}, \hat{j}, \hat{k})$. The axes are rotated anti-clockwise about the Y axis by an angle of 60° . The vector \mathbf{p} in the rotated coordinate system $(\hat{i}', \hat{j}', \hat{k}')$ is
 a) $(1 - \sqrt{3})\hat{i}' + 3\hat{j}' + (1 + \sqrt{3})\hat{k}'$
 b) $(1 + \sqrt{3})\hat{i}' + 3\hat{j}' + (1 - \sqrt{3})\hat{k}'$
 c) $(1 - \sqrt{3})\hat{i}' + (3 + \sqrt{3})\hat{j}' + 2\hat{k}'$
 d) $(1 - \sqrt{3})\hat{i}' + (3 - \sqrt{3})\hat{j}' + 2\hat{k}'$
- 8) The contour integral $\oint \frac{dz}{z^4 + a^4}$ is to be evaluated on a circle of radius $2a$ centered at the origin. It will have contributions only from the points
 a) $\frac{1+i}{\sqrt{2}}a$ and $\frac{1-i}{\sqrt{2}}a$
 b) ia and $-ia$
 c) $ia, -ia, \frac{-i}{\sqrt{2}}a$ and $\frac{1-i}{\sqrt{2}}a$
 d) $\frac{1+i}{\sqrt{2}}a, \frac{1-i}{\sqrt{2}}a, \frac{-i}{\sqrt{2}}a$ and $\frac{1-i}{\sqrt{2}}a$
- 9) Inverse Laplace transform of $\frac{s+1}{s^2-4}$ is
 a) $\cos 2x + \frac{1}{2} \sin 2x$
 b) $\cos x + \frac{1}{2} \sin x$
 c) $\cosh x + \frac{1}{2} \sinh x$
 d) $\cosh 2x + \frac{1}{2} \sinh 2x$
- 10) The points, where the series solution of the Legendre differential equation

$$(1-x^2)\frac{d^2y}{dx^2} - 2x\frac{dy}{dx} + \frac{3}{2}\left(\frac{3}{2} + 1\right)y = 0$$
 will diverge, are located at
 a) 0 and 1
 b) 0 and -1
 c) -1 and 1
 d) $\frac{3}{2}$ and $\frac{5}{2}$
- 11) Solution of the differential equation $x\frac{dy}{dx} + y = x^4$, with the boundary condition that $y = 1$ at $x = 1$, is
 a) $y = 5x^4 - 4$
 b) $y = \frac{x^4}{5} + \frac{4x}{5}$
 c) $y = \frac{4x^4}{5} + \frac{1}{5x}$
 d) $y = \frac{x^4}{5} + \frac{4}{5x}$

12) Match the following


P. rest mass	1. timelike vector
Q. charge	2. Lorentz invariant
R. four-momentum	3. tensor of rank 2
S. electromagnetic field	4. conserved and Lorentz invariant

- a) $P - 2, Q - 4, R - 3, S - 1$
 b) $P - 4, Q - 2, R - 1, S - 3$
 c) $P - 2, Q - 4, R - 1, S - 3$
 d) $P - 4, Q - 2, R - 3, S - 1$

13) The moment of inertia of a uniform sphere of radius r about an axis passing through its center is given by

$$\frac{2}{5} \left(\frac{4\pi}{3} r^3 \rho \right)$$

A rigid sphere of uniform mass density ρ and radius R has two smaller spheres of radius $R/2$ hollowed out of it, as shown in the figure. The moment of inertia of the resulting body about the Y axis is:



- a) $\frac{\pi \rho R^5}{4}$
 b) $\frac{5\pi \rho R^5}{12}$
 c) $\frac{7\pi \rho R^5}{12}$
 d) $\frac{3\pi \rho R^5}{4}$

14) The Lagrangian of a particle of mass m is

$$L = \frac{m}{2} \left[\left(\frac{dx}{dt} \right)^2 + \left(\frac{dy}{dt} \right)^2 + \left(\frac{dz}{dt} \right)^2 \right] - \frac{V}{2} (x^2 + y^2) + W \sin \omega t$$

where V , W , and ω are constants. The conserved quantities are:

- a) energy and z-component of linear momentum only.
 b) energy and z-component of angular momentum only.
 c) z-components of both linear and angular momenta only.
 d) energy and z-components of both linear and angular momenta.

- 15) Three particles of mass m , each situated at $x_1(t)$, $x_2(t)$, and $x_3(t)$ respectively, are connected by two springs of spring constant k and un-stretched length ℓ . The system is free to oscillate only in one dimension along the straight line joining all three particles. The Lagrangian of the system is:

$$\begin{aligned} \text{a) } L &= \frac{m}{2} \left[\left(\frac{dx_1}{dt} \right)^2 + \left(\frac{dx_2}{dt} \right)^2 + \left(\frac{dx_3}{dt} \right)^2 \right] - \frac{k}{2} (x_1 - x_2 - \ell)^2 - \frac{k}{2} (x_3 - x_2 - \ell)^2 \\ \text{b) } L &= \frac{m}{2} \left[\left(\frac{dx_1}{dt} \right)^2 + \left(\frac{dx_2}{dt} \right)^2 + \left(\frac{dx_3}{dt} \right)^2 \right] - \frac{k}{2} (x_1 - x_3 - \ell)^2 - \frac{k}{2} (x_3 - x_2 - \ell)^2 \\ \text{c) } L &= \frac{m}{2} \left[\left(\frac{dx_1}{dt} \right)^2 + \left(\frac{dx_2}{dt} \right)^2 + \left(\frac{dx_3}{dt} \right)^2 \right] - \frac{k}{2} (x_1 - x_2 + \ell)^2 - \frac{k}{2} (x_3 - x_2 + \ell)^2 \\ \text{d) } L &= \frac{m}{2} \left[\left(\frac{dx_1}{dt} \right)^2 + \left(\frac{dx_2}{dt} \right)^2 + \left(\frac{dx_3}{dt} \right)^2 \right] - \frac{k}{2} (x_1 - x_2 - \ell)^2 - \frac{k}{2} (x_3 - x_2 + \ell)^2 \end{aligned}$$

16) The Hamiltonian of a particle is $H = \frac{p^2}{2m} + pq$, where q is the generalized coordinate and p is the corresponding canonical momentum. The Lagrangian is

- a) $\frac{m}{2} \left(\frac{dq}{dt} + q \right)^2$
- b) $\frac{m}{2} \left(\frac{dq}{dt} - q \right)^2$
- c) $\frac{m}{2} \left(\frac{dq}{dt} \right)^2 + q \frac{dq}{dt} - q^2$
- d) $\frac{m}{2} \left(\frac{dq}{dt} \right)^2 - q \frac{dq}{dt} + q^2$

17) A toroidal coil has N closely-wound turns. Assume the current through the coil to be I and the toroid is filled with a magnetic material of relative permittivity μ_r . The magnitude of magnetic induction \mathbf{B} inside the toroid, at a radial distance r from the axis, is given by

- a) $\mu_r \mu_0 N I r$
- b) $\frac{\mu_r \mu_0 N I}{r}$
- c) $\frac{\mu_r \mu_0 N I}{2\pi r}$
- d) $\frac{2\pi \mu_r \mu_0 N I r}{r}$