

Early Detection of Turner Syndrome through Machine Learning Algorithms

Manish M, Sanjay. M, Manoj Kumar. V Department of Computer Science and Application, Jeppiaar College of Arts and Science, Padur, Chennai.

Abstract

Turner Syndrome (TS) is a chromosomal disorder affecting females, characterized by the absence or partial alteration of one X chromosome. This non-inherited condition arises randomly during gamete formation and can lead to a range of physical and developmental features. This research investigates the application of a Random Forest machine learning algorithm for the detection of TS. By leveraging the power of machine learning, we aim to develop a more efficient and potentially earlier diagnostic tool for this complex condition, ultimately contributing to improved management and care for individuals with Turner Syndrome.

Keywords: Turner Syndrome, X chromosomes, Random Forest Algorithm, Machine Learning.

1. Introduction

Turner Syndrome (TS) is a complex chromosomal disorder exclusively affecting females, characterized by the absence or partial alteration of one of the two X chromosomes. Typically, females possess two X chromosomes (XX); however, in TS, one X chromosome is

either completely missing (monosomy X), partially deleted, or structurally altered.




Fig. 1.1

This cytogenetic abnormality occurs randomly, usually during the formation of reproductive cells (gametes) – either the egg or the sperm – and is not typically inherited from the parents. The absence or abnormality of this crucial genetic material disrupts normal development, leading to a wide spectrum of physical characteristics, including short stature, ovarian insufficiency, and characteristic facial features, as well as potential developmental challenges. Early and accurate diagnosis of TS is crucial for timely intervention and management, which can significantly

improve the quality of life for affected individuals. Traditional diagnostic methods, such as karyotyping, can be time-consuming and may not always detect subtle chromosomal variations. Therefore, there is a need for more efficient and potentially earlier diagnostic tools. This research explores the application of a Random Forest machine learning algorithm for the detection of TS. Random Forest, an ensemble learning method, has demonstrated considerable success in various classification tasks due to its ability to handle high-dimensional data and complex relationships. By training the Random Forest model on relevant clinical and genetic data, we aim to develop a robust and accurate diagnostic tool for TS. This approach has the potential to expedite the diagnostic process, enabling earlier interventions and ultimately contributing to improved management and care for individuals with Turner Syndrome.

2. Literature review

Turner syndrome: narrative review of genetics and clinical aspects of management Elizabeth B. Fudge Turner syndrome (TS), affecting approximately 1 in 2,500 females, arises from the loss or alteration of an X chromosome, leading to diverse health issues like short stature, ovarian failure, cardiovascular problems, and neurocognitive deficits. Diagnosis can

occur at birth for pronounced cases, but milder presentations often lead to later detection, contributing to increased morbidity and mortality. This review examines TS's genetic mechanisms and clinical management, focusing on short stature, gonadal failure, cardiovascular disease, and neurocognitive challenges, highlighting advancements and areas needing further research.

An Automated Approach to Diagnose Turner Syndrome Using Ensemble Learning Methods by Qing Zhao; Guohong Yao; Faheem Akhtar; Jianqiang Li; Yan Pei

All Authors This research explores using ensemble learning and facial recognition for Turner syndrome (TS) diagnosis, a common chromosomal disorder often difficult to diagnose early without cytogenetic testing. Facial features are extracted using PCA and KPCA, and ensemble models (majority voting and stacking) combine base learners (SVM) for classification. Achieving 88.1% accuracy, the proposed method demonstrates potential for automated TS diagnosis, aiding clinicians. While stacking proved most effective, further research will focus on identifying more discriminative TS facial characteristics to improve diagnostic accuracy and potentially apply this approach to other genetic disorders.

The Changing Face of Turner Syndrome by Claus H Gravholt, Mette Viuff, Jesper Just, Kristian Sandahl, Sara Brun, Janielle van der Velden, Niels H Andersen, Anne Skakkebaek Turner syndrome (TS), characterized by missing or altered X chromosomes in females, presents a spectrum of health issues including short stature, ovarian dysfunction, congenital malformations, endocrine disorders, and neurocognitive deficits, leading to increased morbidity and mortality. Recent advances in understanding TS's genomic architecture, including candidate genes and altered pathways, alongside improved clinical management approaches, are being reviewed, though epidemiological studies face challenges like Berkson's bias; international registries may offer solutions for future research into areas like HRT effects and quality of life.

Turner Syndrome by Andreas Schüring, Frank Nawroth & Michael von Wolff Turner syndrome (TS) encompasses various chromosomal abnormalities, including monosomy X, mosaicism, and X chromosome structural issues, impacting fertility differently. Monosomy often leads to early ovarian insufficiency, while mosaicism and structural abnormalities may preserve it. Thus, genotype is crucial when considering fertility preservation (oocyte

or ovarian tissue freezing) in TS patients, especially given the increased pregnancy risks associated with the condition.

3. Existing work

Several methods are employed in the detection of Turner Syndrome (TS), they are :

Traditional TS Diagnosis Methods:

Karyotyping: The "Introduction" mentions karyotyping as a traditional method. Existing work should discuss the details of karyotyping, its strengths (gold standard for chromosomal analysis), and limitations (time-consuming, may not detect subtle variations).

Fluorescence in situ hybridization (FISH): FISH is another cytogenetic technique often used. Research should cover how FISH is used in TS diagnosis and its advantages/disadvantages.

Other Cytogenetic Techniques: Other methods like chromosomal microarray analysis (CMA) should also be discussed if relevant.

Clinical Features and Phenotype of TS:

The "Introduction" mentions short stature, ovarian insufficiency, and characteristic facial features. Existing literature

extensively documents the wide range of clinical manifestations in TS. Research should cite key papers describing these features and how they are used in clinical diagnosis.

Facial Features in TS: The calculations and the reference to facial recognition in previous examples suggest that facial features are being considered as a diagnostic tool. Existing work exploring the use of facial analysis or anthropometry in TS diagnosis needs to be reviewed.

Machine Learning in Medical Diagnosis:

The core of the research is applying a Random Forest algorithm. Existing work on the application of machine learning, particularly Random Forest, in the diagnosis of other genetic disorders or medical conditions should be included. This demonstrates the feasibility and potential benefits of such approaches.

Ensemble Learning: Since the research mentions ensemble learning, existing work on ensemble methods (like bagging, boosting, stacking) and their effectiveness in classification tasks should be reviewed.

Feature Selection: If feature selection was employed to choose the most informative features for the Random Forest model, existing feature selection techniques (e.g.,

filter methods, wrapper methods, embedded methods) and their applications in similar domains should be discussed.

Risk Assessment in Stock Market (If Applicable):

The calculations related to Bayes' Theorem and market trends seem out of context for a paper focused on TS diagnosis. If there's a connection to stock market risk assessment, then existing work in that area needs to be explained and justified in the context of TS. It's unclear how this is relevant to the core topic.

4. Proposed work

Turner Syndrome

Turner syndrome (TS) is a chromosomal disorder that affects females. It occurs when one of the two X chromosomes is missing or partially altered. This can lead to a variety of symptoms, which can vary in severity from person to person.

Common symptoms:

Short stature: This is one of the most common features of TS. Girls with TS tend to be shorter than their peers, and their adult height is often significantly less than what would be expected for a female member of their family.

Ovarian insufficiency: This means that the ovaries do not develop properly and do not produce eggs.




Fig. 4.1

This can lead to infertility and early menopause.

Characteristic facial features: These can include a webbed neck, low-set ears, a broad chest with widely spaced nipples, and a high, narrow roof of the mouth.

Other potential health problems:

Heart defects: These can range from mild to severe and may require surgery to correct.

Kidney problems: These are less common than heart defects but can still occur.

Hearing loss: This can be due to frequent ear infections or gradual loss of nerve function.

Vision problems: These can include strabismus (crossed eyes), near sightedness, and other issues.

Autoimmune disorders: Girls with TS have an increased risk of developing autoimmune disorders, such as hypothyroidism and diabetes.

Skeletal problems: These can include scoliosis (curvature of the spine) and osteoporosis (weak bones).

Learning disabilities: Girls with TS usually have normal intelligence, but they may have learning disabilities, particularly in math and spatial reasoning.

Mental health issues: Girls with TS may be more likely to experience anxiety, depression, and ADHD.

Random Forest Algorithm

The Random Forest algorithm is a powerful and popular machine learning technique that combines the predictions of multiple decision trees to make more accurate and robust predictions.




Fig. 4.2

Random Forest Algorithm in the detection of Turner Syndrome

The Random Forest algorithm offers a robust approach to detecting Turner Syndrome (TS) by leveraging its capacity to handle complex datasets and identify intricate patterns. Initially, relevant data encompassing clinical features (e.g., height, weight, facial measurements), genetic markers (e.g., karyotype data, specific chromosomal abnormalities), and potentially other relevant information is gathered and pre-processed. This data is then used to train the Random Forest model, which constructs an ensemble of decision trees, each trained on a random subset of the data and features. This ensemble approach enhances the model's ability to generalize and reduces the risk of overfitting. Following training, the model's performance is evaluated on unseen data using metrics like accuracy, precision, recall, and AUC. Furthermore, Random Forest provides insights into feature importance, highlighting the most influential factors in TS detection. Ultimately, this trained model can be deployed as a diagnostic tool, assisting clinicians in identifying individuals with TS more efficiently and potentially earlier.

Advantages of Random Forest in TS Detection:

Handles High-Dimensional Data: TS diagnosis may involve a large number of


clinical and genetic features, which Random Forest can effectively handle.

Robust to Outliers: Random Forest is less sensitive to outliers in the data, which is important in medical datasets where outliers are common.

Good Generalization Performance: Random Forest is less prone to overfitting, which means it can generalize well to unseen data.

Provides Feature Importance: Random Forest can identify the most important features for TS detection, which can be valuable for clinical understanding.

Random Forest Algorithm's results



The image shows a terminal window with a dark background and light-colored text. At the top, it says 'Accuracy: 1.0000'. Below that is a 'Classification Report' table. The table has columns for precision, recall, f1-score, and support. There are two rows for classes 0 and 1, all with values of 1.00. Below the table are summary statistics: accuracy (1.00), macro avg (1.00), and weighted avg (1.00). At the bottom is a 'Feature Importances (Sorted):' list with features and their importance scores.

Accuracy: 1.0000				
Classification Report:				
	precision	recall	f1-score	support
0	1.00	1.00	1.00	1
1	1.00	1.00	1.00	1
accuracy			1.00	2
macro avg	1.00	1.00	1.00	2
weighted avg	1.00	1.00	1.00	2
Feature Importances (Sorted):				
Feature	Importance			
Age (Years)	0.391970			
Height (cm)	0.383615			
Heart Murmur	0.097628			
Infertility	0.046691			
Eye Issues	0.031473			
Webbing	0.023588			
Hairline	0.012815			
Chest	0.010104			
Fingernails	0.002110			

Fig. 4.3

The provided output shows a machine learning model evaluation, likely for a binary classification task. The model achieved a perfect 100% accuracy (1.0000), which is immediately suspicious. A closer look reveals a classification report based on a test set containing only two data points (support of 1 for each class). This tiny

sample size makes the perfect accuracy meaningless and suggests severe overfitting. The report shows perfect precision, recall, and F1-scores for both classes, which is expected when dealing with such limited data. Additionally, the output lists feature importances, with "Age (Years)" and "Height (cm)" being the most influential. However, these importance values are also unreliable due to the small dataset. Essentially, the results are not trustworthy and highlight the critical need for larger datasets and proper evaluation techniques in machine learning. The perfect accuracy is a clear indicator of a flawed evaluation process and does not reflect the model's true capabilities.




Fig. 4.4

This image displays a bar graph of feature importances from a Random Forest model trained to predict Turner Syndrome. The horizontal axis, labeled "Importance Score," ranges from 0.00 to 0.40 and visually represents the degree to which each feature contributes to the prediction. The

vertical axis lists the features considered by the model, including "Age (Years)," "Height (cm)," "Heart Murmur," "Infertility," "Eye Issues," "Webbing," "Hairline," "Chest," and "Fingernails," ordered from most to least important. The length of each bar corresponds to the feature's importance score, with longer bars indicating greater influence on the prediction of Turner Syndrome. The graph clearly shows "Age (Years)" and "Height (cm)" as the dominant predictors, exhibiting substantially higher importance scores than all other features. "Heart Murmur" also appears to play a noticeable role, while "Infertility," "Eye Issues," and the remaining features show considerably lower importance, suggesting they contribute less to the model's ability to identify Turner Syndrome. This visualization provides a quick and intuitive understanding of which clinical features are most strongly associated with the presence of Turner Syndrome according to the trained Random Forest model. It suggests that age and height are particularly relevant factors in this specific predictive model.

Patient ID	Age (Years)	Height (cm)	Webbing (Yes/No)	Chest (Yes/No)	Hairline (Yes/No)	Fingernails (Yes/No)	Heart Murmur (Yes/No)	Eye Issues (Yes/No)	Infertility (Yes/No)	Syndrome (Yes/No)
1	10	120	No	No	No	No	No	No	N/A	No
2	15	145	Yes	Yes	Yes	Yes	No	No	N/A	Yes
3	8	105	No	No	No	No	Yes	No	N/A	Yes
4	12	130	No	No	No	No	No	Yes	N/A	Yes
5	18	162	No	No	No	No	No	No	Yes	Yes
6	11	125	No	No	No	No	No	No	N/A	No
7	16	140	Yes	Yes	Yes	Yes	Yes	No	N/A	Yes
8	9	110	No	No	No	No	No	No	N/A	No
9	14	138	Yes	Yes	Yes	Yes	No	Yes	N/A	Yes
10	17	150	No	No	No	No	No	No	Yes	Yes

Fig. 4.5

This table presents a dataset of ten patients and their associated clinical characteristics, potentially for the purpose of studying or predicting Turner Syndrome. The data includes demographic information like "Patient ID," "Age (Years)," and "Height (cm)." It also records the presence or absence of various clinical features indicated by "Yes" or "No," such as "Webbing," "Chest," "Hairline," "Fingernails," "Heart Murmur," and "Eye Issues." Additionally, the dataset notes "Infertility" as either "Yes," "No," or "N/A" (Not Applicable). The final column, "Turner Syndrome (Yes/No)," indicates whether each patient has been diagnosed with the condition, serving as a potential ground truth label for analysis or model training. This dataset, while small, provides a structured overview of patient characteristics and their Turner Syndrome status, suggesting its use in exploring potential relationships or building predictive models for the condition.

5. Conclusion

The research explores the application of a Random Forest algorithm to detect Turner Syndrome (TS) using clinical and demographic data. While the initial results show a perfect accuracy of 100. The feature importance analysis suggests that age and height are the most influential factors in the

model's predictions, followed by heart murmur. The research acknowledges the need for more data and proper evaluation techniques to build a robust and reliable diagnostic tool for TS. In its current state, the model's performance and feature importance findings should not be interpreted as clinically relevant or indicative of actual predictive power for Turner Syndrome. The study serves as a preliminary exploration of using Random Forest for TS detection. Early Turner Syndrome prediction allows for timely interventions like growth hormone therapy and proactive management of associated health issues, improving long-term outcomes. It also enables access to psychological support, educational planning, and fertility preservation discussions, enhancing quality of life.

6. Reference

- [1] Turner syndrome: narrative review of genetics and clinical aspects of management Elizabeth B. Fudge
- [2] An Automated Approach to Diagnose Turner Syndrome Using Ensemble Learning Methods by Qing Zhao; Guohong Yao; Faheem Akhtar; Jianqiang Li; Yan Pei
- [3] The Changing Face of Turner Syndrome by Claus H Gravholt, Mette

Viuff, Jesper Just, Kristian Sandahl, Sara Brun, Janielle van der Velden, Niels H Andersen, Anne Skakkebaek

[4] Turner Syndrome by Andreas Schüring, Frank Nawroth & Michael von Wolff

[5] Säwendahl L, Davenport ML. Delayed diagnoses of Turner's syndrome: proposed guidelines for change. *J Pediatr*. 2000 Oct;137(4):455-9. [[PubMed](#)]

[6] Barrenäs ML, Nylén O, Hanson C. The influence of karyotype on the auricle, otitis media and hearing in Turner syndrome. *Hear Res*. 1999 Dec;138(1-2):163-70. [[PubMed](#)]

[7] Adhikary HP. Ocular manifestations of Turner's syndrome. *Trans Ophthalmol Soc U K* (1962). 1981;101 (Pt 4):395-6. [[PubMed](#)]

[8] Lowenstein EJ, Kim KH, Glick SA. Turner's syndrome in dermatology. *J Am Acad Dermatol*. 2004 May;50(5):767-76. [[PubMed](#)]

[9] Lunding SA, Aksglaede L, Anderson RA, Main KM, Juul A, Hagen CP, Pedersen AT. AMH as Predictor of Premature Ovarian Insufficiency: A Longitudinal Study of 120 Turner Syndrome Patients. *J Clin Endocrinol Metab*. 2015 Jul;100(7):E1030- [[PubMed](#)]

[10] Kim HK, Gottliebson W, Hor K, Backeljauw P, Gutmark-Little I, Salisbury SR, Racadio JM, Helton-Skally K, Fleck R. Cardiovascular anomalies in Turner syndrome: spectrum, prevalence, and cardiac MRI findings in a pediatric and young adult population. *AJR Am J Roentgenol*. 2011 Feb;196(2):454-60. [[PubMed](#)]

[11] Cleemann L, Holm K, Fallentin E, Skouby SO, Smedegaard H, Møller N, Borch-Christensen H, Jeppesen EM, Wieslander SB, Andersson AM, Cohen A, Højbjerg Gravholt C: Uterus and ovaries in girls and young women with Turner syndrome evaluated by ultrasound and magnetic resonance imaging. *Clin Endocrinol (Oxf)* 2011; 74: 756–761.

[12] Marshall WA, Tanner JM: Variations in pattern of pubertal changes in girls. *Arch Dis Child* 1969; 44: 291–303.

[13] Negreiros LP, Bolina ER, Guimarães MM: Pubertal development profile in patients with Turner syndrome. *J Pediatr Endocrinol Metab* 2014; 27: 845–849.

[14] Tanaka T, Igarashi Y, Ozono K, Ohyama K, Ogawa M, Osada H, Onigata K, Kanzaki S, Kohno H, Seino Y, Takahashi H, Tajima T, Tachibana K, Tanaka H, Nishi Y, Hasegawa T, Fujita K, Yorifuji T, Horikawa R, Yokoya S: Frequencies of

spontaneous breast development and spontaneous menarche in Turner syndrome in Japan. *Clin Pediatr Endocrinol* 2015; 24: 167–173.

[15] Mortensen KH, Rohde MD, Uldbjerg N, Gravholt CH: Repeated spontaneous pregnancies in 45,X Turner syndrome. *Obstet Gynecol* 2010; 115(pt 2):446–449.

[16] Chrysis D, Spiliotis BE, Stene M, Cacciari E, Davenport ML: Gonadotropin secretion in girls with Turner syndrome measured by an ultrasensitive immunochemiluminometric assay. *Horm Res* 2006; 65: 261–266.

[17] Conte FA, Grumbach MM, Kaplan SL: A diphasic pattern of gonadotropin secretion in patients with the syndrome of gonadal dysgenesis. *J Clin Endocrinol Metab* 1975; 40: 670–674.

[18] Alves ST, Gallicchio CT, Guimarães MM, Santos M: Gonadotropin levels in Turner's syndrome: correlation with breast development and hormone replacement therapy. *Gynecol Endocrinol* 2003; 17: 295–301.

[19] Aso K, Koto S, Higuchi A, Ariyasu D, Izawa M, Miyamoto-Igaki J, Hasegawa Y: Serum FSH level below 10 mIU/mL at twelve years old is an index of spontaneous

and cyclical menstruation in Turner syndrome. *Endocr J* 2010; 57: 909–913.

[20] Messina MF, Aversa T, Salzano G, Costanzo D, Sferlazzas C, Mirabelli S, Zirilli G, Lombardo F: Inhibin B in adolescents and young adults with Turner syndrome. *J Pediatr Endocrinol Metab* 2015; 28: 1209–1214.