

## Assignment 1 Report

Entry Number: 2016CS10363

Name: Manish Tanwar

## 1 A

## 2 Locally Weighted Linear Regression:

## (a) Linear Regression (unweighted):

$$\theta = \begin{bmatrix} \theta_0 \\ \theta_1 \end{bmatrix} = \begin{bmatrix} 0.327680 \\ 0.175316 \end{bmatrix}$$




Figure 1: Linear Regression Plot (Underfitting)

Linear Regression(unweighted) is not a good fit for the data as shown in the Figure 1(underfitting).

## (b) Locally Weighted Linear Regression:

Weights :  $w^{(i)} = \exp\left(-\frac{(x - x^{(i)})^2}{2\tau^2}\right)$  (where  $\tau$  = Bandwidth Parameter)

Error Function:  $J(\theta) = \frac{1}{2m}(X\theta - Y)^T W (X\theta - Y)$  (where  $W = \text{diag}(w^{(i)})$ )

Minima:  $\nabla_{\theta} J(\theta) = 0 \Rightarrow \theta = (X^T W X)^{-1} X^T W Y$

**Plots:**




Figure 2: Locally Weighted Linear Regression Plot ( $\tau = 0.8$ )

**(c) Plots on Varving Bandwidth Parameter( $\tau$ )**




Figure 3:  $\tau = 0.1$




Figure 4:  $\tau = 0.3$




Figure 5:  $\tau = 2$




Figure 6:  $\tau = 10$

## Analysis:

- When Bandwidth Parameter( $\tau$ ) is:
  - **Too Small:** the model **overfits** the training data.
  - **Too Large:** The model tries to **underfits** the training data.
- $\tau = 0.8$  works the best.

## 3 Logistic Regression:

Log Likelihood: 
$$LL(\theta) = \sum_{i=1}^m y^{(i)} \log(h_{\theta}(x^{(i)})) + (1 - y^{(i)}) \log(1 - h_{\theta}(x^{(i)}))$$

$$\nabla_{\theta} LL(\theta) = X^T(Y - g(X\theta)) \quad (\text{where } g(x) = \frac{1}{1 + \exp(-x)})$$

Hessian Matrix: 
$$H = \nabla_{\theta}^2 LL(\theta) = -X^T D X$$
  
 (where  $D = \text{diag}(g(x^{(i)T}\theta)(1 - g(x^{(i)T}\theta)))$ )

### Newton's Method:

$$\theta^{(t+1)} = \theta^{(t)} - H^{-1} \nabla_{\theta} LL(\theta) \big|_{\theta_t}$$

### Convergence Condition:

$$|\theta_j^{(t+1)} - \theta_j^{(t)}| < \epsilon \quad \forall j \in 1, 2 \dots n \quad (\text{for a sufficiently small } \epsilon)$$

### Resulting Parameters:

$$\theta = \begin{bmatrix} \theta_0 \\ \theta_1 \\ \theta_2 \end{bmatrix} = \begin{bmatrix} 0.223295 \\ 1.962616 \\ -1.964861 \end{bmatrix}$$

Decision Boundary is the straight line boundary separating the region where  $h_{\theta}(x) \geq 0.5$  (class  $y = 1$ ) from where  $h_{\theta}(x) \leq 0.5$  (class  $y = 0$ ).

### Plot:




Figure 7: Logistic Regression : Given data and Linear Separator

## 4 Gaussian Discriminant Analysis:

(a):

$$\phi = \frac{1}{m} \sum_{i=1}^m \mathbb{1}\{y^{(i)} = 1\} \quad \mu_0 = \frac{\sum_{i=1}^m \mathbb{1}\{y^{(i)} = 0\} x^{(i)}}{\sum_{i=1}^m \mathbb{1}\{y^{(i)} = 0\}} \quad \mu_1 = \frac{\sum_{i=1}^m \mathbb{1}\{y^{(i)} = 1\} x^{(i)}}{\sum_{i=1}^m \mathbb{1}\{y^{(i)} = 1\}}$$

$$\Sigma = \frac{1}{m} \sum_{i=1}^m (x^{(i)} - \mu_{y^{(i)}})(x^{(i)} - \mu_{y^{(i)}})^T$$

which can be written as:  $\Sigma = \frac{1}{m} W^T W$  (where  $W = X - Y\mu_1^T - (1 - Y)\mu_0^T$ )

**Resulting Parameters:**

$$\phi = 0.5 \quad \mu_0 = \begin{bmatrix} 98.38 \\ 429.66 \end{bmatrix} \quad \mu_1 = \begin{bmatrix} 137.46 \\ 366.62 \end{bmatrix} \quad \Sigma = \begin{bmatrix} 287.482 & -26.748 \\ -26.748 & 1123.25 \end{bmatrix}$$

(b) & (c):

(i) Linear Boundary Equation:

$$2(\mu_1^T - \mu_0^T \Sigma^{-1})x + \mu_0^T \Sigma^{-1} \mu_0 - \mu_1^T \Sigma^{-1} \mu_1 - 2 \log \left( \frac{1 - \phi}{\phi} \right) = 0$$

(ii) Plot:




Figure 8: GDA : Given data and Linear Separator

(d):

$$\phi = \frac{1}{m} \sum_{i=1}^m \mathbb{1}\{y^{(i)} = 1\} \quad \mu_0 = \frac{\sum_{i=1}^m \mathbb{1}\{y^{(i)} = 0\} x^{(i)}}{\sum_{i=1}^m \mathbb{1}\{y^{(i)} = 0\}} \quad \mu_1 = \frac{\sum_{i=1}^m \mathbb{1}\{y^{(i)} = 1\} x^{(i)}}{\sum_{i=1}^m \mathbb{1}\{y^{(i)} = 1\}}$$

$$\Sigma_0 = (X - \mu_0^{ext})^T \text{diag}(y^{(i)}) (X - \mu_0^{ext}) \quad (\text{where } \mu_0^{ext} \text{ is } m \times n \text{ matrix with each row} = \mu_0^T)$$

**Resulting Parameters:**

$$\phi = 0.5 \quad \mu_0 = \begin{bmatrix} 98.38 \\ 429.66 \end{bmatrix} \quad \mu_1 = \begin{bmatrix} 137.46 \\ 366.62 \end{bmatrix}$$

$$\Sigma_0 = \begin{bmatrix} 255.3956 & -184.3308 \\ -184.3308 & 1371.1044 \end{bmatrix} \quad \Sigma_1 = \begin{bmatrix} 319.5684 & 130.8348 \\ 130.8348 & 875.3956 \end{bmatrix}$$

(e):

(i) Quadratic Boundary Equation:

$$(x - \mu_1)^T \Sigma_1^{-1} (x - \mu_1) - (x - \mu_0)^T \Sigma_0^{-1} (x - \mu_0) + \log \left( \frac{|\Sigma_1|}{|\Sigma_0|} \frac{(1 - \phi)^2}{\phi^2} \right) = 0$$

(ii) Plot:




Figure 9: GDA : Given data and Quadratic Separator

(f) Analysis: