COL774: Machine Learning **Date: February 12, 2019**

Assignment 1 Report

Entry Number: 2016CS10363 **Name: Manish Tanwar**

1 A

2 B

3 Logistic Regression:

Log Likelihood: $LL(\theta) = \sum_{i=1}^{m} y^{(i)} log(h_\theta(x^{(i)})) + (1 - y^{(i)}) log(1 - h_\theta(x^{(i)}))$

$$\nabla_\theta LL(\theta) = X^T(Y - g(X\theta)) \qquad \text{(where } g(x) = \frac{1}{1 + \exp(-x)}\text{)}$$

Hessian Matrix: $H = \nabla^2_\theta LL(\theta) = -X^T W X$

$$\text{(where } W = diag(\, g(x^{(i)T}\theta)(1 - g(x^{(i)T}\theta))\,)$$

Newton's Method:

$$\theta^{(t+1)} = \theta^{(t)} - H^{-1} \nabla_\theta LL(\theta)\big|_{\theta_t}$$

Convergence Condition:

$$\left|\theta_j^{(t+1)} - \theta_j^{(t)}\right| < \epsilon \qquad \text{(for a sufficiently small } \epsilon\text{)}$$

Resulting Parameters:

$$\theta = \begin{bmatrix} \theta_0 \\ \theta_1 \\ \theta_2 \end{bmatrix} = \begin{bmatrix} 0.223295 \\ 1.962616 \\ -1.964861 \end{bmatrix}$$

Decision Boundary is the straight line boundary separating the region where $h_\theta(x) \geq 0.5$ (class $y = 1$) from where $h_\theta(x) \leq 0.5$ (class $y = 0$).

Plot:



4 Gaussian Discrmimant Analysis: