

NCERT Solutions For Class 10 Chapter 8 Introduction to Trigonometry Exercise 8.3

Q1. Evaluate:

(i)
$$\frac{\sin 18^{\circ}}{\cos 72^{\circ}}$$

(ii)
$$\frac{\tan 26^{\circ}}{\cot 64^{\circ}}$$

Ans: (i)
$$\frac{\sin 18^{\circ}}{\cos 72^{\circ}} = \frac{\sin (90^{\circ} - 72^{\circ})}{\cos 72^{\circ}}$$

$$= \frac{\cos 72^{\circ}}{\cos 72^{\circ}} = 1$$

(ii)
$$\frac{\tan 26^\circ}{\cot 64^\circ} = \frac{\tan (90^\circ - 64^\circ)}{\cot 64^\circ}$$

$$=\frac{\cot 64^{\circ}}{\cot 64^{\circ}}=1$$

$$= \cos(90^{\circ} - 42^{\circ}) - \sin 42^{\circ}$$

$$= \sin 42^{\circ} - \sin 42^{\circ} = 0$$

$$=\cos ec \left(90^{\circ} - 59^{\circ}\right) - \sec 59^{\circ}$$

$$= \sec 59^{\circ} - \sec 59^{\circ} = 0$$

Q2. Show that:

(ii)
$$\cos 38^{\circ} \cos 52^{\circ} - \sin 38^{\circ} \sin 52^{\circ} = 0$$

Ans: (i)L.H.S. tan 48°tan 23°tan 42°tan 67°

=
$$\tan (90^{\circ} - 42^{\circ}) \tan (90^{\circ} - 67^{\circ}) \tan 42^{\circ} \tan 67^{\circ}$$

 $_{\pm}$ cot 42° cot 67° tan 42° tan 67°

$$=\frac{1}{\tan 42^{\circ}} \cdot \frac{1}{\tan 67^{\circ}} \cdot \tan 42^{\circ} \cdot \tan 67^{\circ} = 1 = \text{R.H.S.}$$

$$= \cos \left(90^{\circ} - 52^{\circ}\right) \cdot \cos \left(90^{\circ} - 38^{\circ}\right) - \sin 38^{\circ} \cdot \sin 52^{\circ}$$

$$= \sin 52^{\circ} \sin 38^{\circ} - \sin 38^{\circ} \sin 52^{\circ} = 0 = R.H.S.$$

Q3. If $\tan 2A = \cot(A-18^{\circ})$, where 2A is an acute angle, find the value of A.

Ans: Given: $\tan 2A = \cot (A-18^{\circ})$

$$\Rightarrow \cot(90^{\circ} - 2A) = \cot(A - 18^{\circ})$$

$$\Rightarrow$$
 90° - 2 $A = A - 18$ °

$$\Rightarrow$$
 $-2A-A=-18^{\circ}-90^{\circ}$

$$\Rightarrow$$
 $-3A = -108^{\circ}$

$$\Rightarrow$$
 A = 36°

Q4. If $\tan A = \cot B$, prove that $A + B = 90^\circ$.

Ans: Given: $\tan A = \cot B$

$$\Rightarrow \cot(90^{\circ} - A) = \cot B$$

$$\Rightarrow 90^{\circ} - A = B$$

$$\Rightarrow$$
 A + B = 90°

Q5. If $\sec 4A = \cos ec (A-20^{\circ})$, where 4A is an acute angle, find the value of A.

Ans: Given: $\sec 4A = \cos ec (A - 20^\circ)$

$$\Rightarrow \cos ec(90^{\circ} - 4A) = \cos ec(A - 20^{\circ})$$

$$\Rightarrow$$
 90° -4A = A - 20°

$$\Rightarrow$$
 $-4A-A=-20^{\circ}-90^{\circ}$

$$\Rightarrow$$
 $-5A = -110^{\circ}$

$$\Rightarrow$$
 A = 22°

Q6. If A, B and C are interior angles of a \triangle ABC, then show that $\sin\left(\frac{B+C}{2}\right) = \cos\frac{A}{2}$.

Ans: Given: A, B and C are interior angles of a \triangle ABC.

$$A + B + C = 180^{\circ}$$

$$\Rightarrow \frac{A+B+C}{2} = 90^{\circ}$$

$$\Rightarrow \frac{B+C}{2} = 90^{\circ} - \frac{A}{2}$$

$$\Rightarrow \sin\left(\frac{B+C}{2}\right) = \sin\left(90^{\circ} - \frac{A}{2}\right)$$

$$\Rightarrow \sin\left(\frac{B+C}{2}\right) = \cos\frac{A}{2}$$

Q7. Express $\sin 67^{\circ} + \cos 75^{\circ}$ in terms of trigonometric ratios of angles between 0° and 45° .

Ans:
$$\sin 67^{\circ} + \cos 75^{\circ}$$

= $\sin (90^{\circ} - 23^{\circ}) + \cos (90^{\circ} - 15^{\circ})$
= $\cos 23^{\circ} + \sin 15^{\circ}$