



Functions Ex 2.3 Q7

We are given that f is a real function and g is a function given by $g(x) = 2x$

To prove; $g \circ f = f + f$.

L.H.S

$$\begin{aligned} g \circ f(x) &= g(f(x)) = 2f(x) \\ &= f(x) + f(x) = \text{R.H.S} \end{aligned}$$

$$\Rightarrow g \circ f = f + f$$

Functions Ex 2.3 Q8

$$f(x) = \sqrt{1-x}, \quad g(x) = \log_e^x$$

Domain of f and g are \mathbb{R} .

$$\text{Range of } f = (-\infty, 1)$$

$$\text{Range of } g = (0, e)$$

Clearly $\text{Range } f \subset \text{Domain } g \Rightarrow g \circ f$ exists

$\text{Range } g \subset \text{Domain } f \Rightarrow f \circ g$ exists

$$\therefore g \circ f(x) = g(f(x)) = g(\sqrt{1-x})$$

$$g \circ f(x) = \log_e^{\sqrt{1-x}}$$

Again

$$f \circ g(x) = f(g(x)) = f(\log_e^x)$$

$$f \circ g(x) = \sqrt{1 - \log_e^x}$$

Functions Ex 2.3 Q9

$f : \left(-\frac{\pi}{2}, \frac{\pi}{2}\right) \rightarrow \mathbb{R}$ and $g : [-1, 1] \rightarrow \mathbb{R}$ defined as $f(x) = \tan x$ and $g(x) = \sqrt{1-x^2}$

Range of f : let $y = f(x) \Rightarrow y = \tan x$
 $\Rightarrow x = \tan^{-1} y$

Since $x \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right), y \in (-\infty, \infty)$

\therefore Range of $f \subset$ domain of $g = [-1, 1]$

$\therefore g \circ f$ exists.

By similar argument $f \circ g$ exists.

Now,

$$f \circ g(x) = f(g(x)) = f(\sqrt{1-x^2})$$

$$f \circ g(x) = \tan \sqrt{1-x^2}$$

Again

$$g \circ f(x) = g(f(x))$$

$$= g(\tan x)$$

$$g \circ f(x) = \sqrt{1-\tan^2 x}$$

Functions Ex 2.3 Q10

$$f(x) = \sqrt{x+3} \text{ and } g(x) = x^2 + 1$$

Now,

$$\text{Range of } f = [-3, \infty] \text{ and}$$

$$\text{Range of } g = (1, \infty)$$

Then, Range of $f \subset$ Domain g and

Range of $g \subset$ Domain f

$\therefore f \circ g$ and $g \circ f$ exist.

Now,

$$f \circ g(x) = f(g(x)) = f(x^2 + 1)$$

$$f \circ g(x) = \sqrt{x^2 + 4}$$

Again,

$$g \circ f(x) = g(f(x)) = g(\sqrt{x+3})$$

$$= (\sqrt{x+3})^2 + 1$$

$$g \circ f(x) = x + 4$$

Functions Ex 2.3 Q11(i)

We have, $f(x) = \sqrt{x-2}$

Clearly, $\text{Domain}(f) = [2, \infty)$ and $\text{Range}(f) = [0, \infty)$.

We observe that $\text{range}(f)$ is not a subset of $\text{domain}(f)$.

$$\begin{aligned}\therefore \text{Domain of } (f \circ f) &= \{x : x \in \text{Domain}(f) \text{ and } f(x) \in \text{Domain}(f)\} \\ &= \{x : x \in [2, \infty) \text{ and } \sqrt{x-2} \in [2, \infty)\} \\ &= \{x : x \in [2, \infty) \text{ and } \sqrt{x-2} \geq 2\} \\ &= \{x : x \in [2, \infty) \text{ and } x-2 \geq 4\} \\ &= \{x : x \in [2, \infty) \text{ and } x \geq 6\} \\ &= [6, \infty)\end{aligned}$$

Now,

$$(f \circ f)(x) = f(f(x)) = f(\sqrt{x-2}) = \sqrt{\sqrt{x-2}-2}$$

$\therefore f \circ f : [6, \infty) \rightarrow \mathbb{R}$ defined as

$$(f \circ f)(x) = \sqrt{\sqrt{x-2}-2}$$

***** END *****