NCERT Solutions For Class 10 Maths Polynomials Exercise 2.2

Q 1. Find the zeroes of the following quadratic polynomials and verify the relationship between the zeroes and the coefficients.

$(\text{i})\, x^2 - 2x - 8 \; (\text{ii})\, 4s^2 - 4s + 1 \; (\text{iii})\, 6x^2 - 3 - 7x$

$(\text{iv})\, 4u^2 + 8u \; (\text{v})\, t^2 - 15 \; (\text{vi})\, 3x^2 - x - 4$

Answer :

(i) $x^2 - 2x - 8 = (x-4)(x+2)$

The value of $x^2 - 2x - 8$ is zero when x - 4 = 0 or x + 2 = 0, i.e., when x = 4 or x = - 2

Therefore, the zeroes of $x^2 - 2x - 8$ are 4 and - 2.

Sum of zeroes =

$$4 - 2 = 2 = \frac{-(-2)}{1} = \frac{-(\text{Coefficient of } x)}{\text{Coefficient of } x^2}$$

Product of zeroes

$$= 4 \times (-2) = -8 = \frac{(-8)}{1} = \frac{\text{Constant term}}{\text{Coefficient of } x^2}$$

(ii) $4s^2 - 4s + 1 = (2s-1)^2$

The value of 4s2 - 4s + 1 is zero when 2s - 1 = 0, i.e., $s = \frac{1}{2}$

Therefore, the zeroes of 4s2 - 4s + 1 are $\frac{1}{2}$ and $\frac{1}{2}$

.

Sum of zeroes =

$$\frac{1}{2} + \frac{1}{2} = 1 = \frac{-(-4)}{4} = \frac{-(\text{Coefficient of } s)}{(\text{Coefficient of } s^2)}$$

Product of zeroes $= \frac{1}{2} \times \frac{1}{2} = \frac{1}{4} = \frac{\text{Constant term}}{\text{Coefficient of } s^2}$

Therefore, the zeroes of $4s^2 - 4s + 1$ are $\frac{1}{2}$ and $\frac{1}{2}$

.

Sum of zeroes =

$$\frac{1}{2}+\frac{1}{2}=1=\frac{-(-4)}{4}=\frac{-(\text{Coefficient of } s)}{(\text{Coefficient of } s^2)}$$

Product of zeroes $=\frac{1}{2}\times\frac{1}{2}=\frac{1}{4}=\frac{\text{Constant term}}{\text{Coefficient of } s^2}$

(iii) $6x^2-3-7x=6x^2-7x-3=(3x+1)(2x-3)$

The value of $6x^2 - 3 - 7x$ is zero when $3x + 1 = 0$

or $2x - 3 = 0$, i.e., $x=\frac{-1}{3}$ or $x=\frac{3}{2}$

Therefore, the zeroes of $6x^2 - 3 - 7x$ are

$\frac{-1}{3}$ and $\frac{3}{2}$.

Sum of zeroes =

$$\frac{-1}{3}+\frac{3}{2}=\frac{7}{6}=\frac{-(-7)}{6}=\frac{-(\text{Coefficient of } x)}{\text{Coefficient of } x^2}$$

Product of zeroes =

$$\frac{-1}{3}\times\frac{3}{2}=\frac{-1}{2}=\frac{-3}{6}=\frac{\text{Constant term}}{\text{Coefficient of } x^2}$$

(iv) $4u^2+8u=4u^2+8u+0$

$=4u(u+2)$

The value of $4u2 + 8u$ is zero when $4u = 0$ or $u + 2 = 0$, i.e., $u = 0$ or $u = - 2$

Therefore, the zeroes of $4u2 + 8u$ are 0 and - 2.

Sum of zeroes =

$$0+(-2)=-2=\frac{-(8)}{4}=\frac{-(\text{Coefficient of } u)}{\text{Coefficient of } u^2}$$

Product of zeroes =

$$0\times(-2)=0=\frac{0}{4}=\frac{\text{Constant term}}{\text{Coefficient of } u^2}$$

(v) t^2-15

$=t^2-0t-15$

$=\left(t-\sqrt{15}\right)\left(t+\sqrt{15}\right)$

The value of $t2 - 15$ is zero when $t-\sqrt{15}=0$ or $t+\sqrt{15}=0$, i.e., when

Q 2 . Find a quadratic polynomial each with the given numbers as the sum and product of its zeroes respectively.

(i) $\frac{1}{4},-1$ (ii) $\sqrt{2},\frac{1}{3}$ (iii) $0,\sqrt{5}$

(iv) $1,1$ (v) $-\frac{1}{4},\frac{1}{4}$ (vi) $4,1$

Answer :

(i) $\frac{1}{4},-1$

Let the polynomial be ax^2+bx+c, and its zeroes be α and β.

$$\alpha+\beta=\frac{1}{4}=\frac{-b}{a}$$

$$\alpha\beta=-1=\frac{-4}{4}=\frac{c}{a}$$

If $a=4,$ then $b=-1,\ c=-4$

Therefore, the quadratic polynomial is $4x^2 - x - 4$.

(ii) $\sqrt{2},\frac{1}{3}$

Let the polynomial be ax^2+bx+c, and its zeroes be α and β.

$$\alpha+\beta=\sqrt{2}=\frac{3\sqrt{2}}{3}=\frac{-b}{a}$$

$$\alpha\beta=\frac{1}{3}=\frac{c}{a}$$

If $a=3,$ then $b=-3\sqrt{2},\ c=1$

Therefore, the quadratic polynomial is $3x^2 - 3\sqrt{2}x + 1$.

(iii) $0,\sqrt{5}$

Let the polynomial be ax^2+bx+c, and its zeroes be α and β.

$$\alpha+\beta=0=\frac{0}{1}=\frac{-b}{a}$$

$$\alpha\times\beta=\sqrt{5}=\frac{\sqrt{5}}{1}=\frac{c}{a}$$

If $a=1,$ then $b=0,\ c=\sqrt{5}$

Therefore, the quadratic polynomial is $x^2+\sqrt{5}$.

(iv) 1, 1

Let the polynomial be ax^2+bx+c, and its zeroes be α and β.

$$\alpha+\beta=1=\frac{1}{1}=\frac{-b}{a}$$

$$\alpha\times\beta=1=\frac{1}{1}=\frac{c}{a}$$

If $a=1$, then $b=-1$, $c=1$

Therefore, the quadratic polynomial is x^2-x+1.

(v) $-\frac{1}{4},\frac{1}{4}$

Let the polynomial be ax^2+bx+c, and its zeroes be α and

********** END **********