




Exercise 10A

Question 21:

Given: ABCD is a parallelogram and P, Q, R and S are the midpoints of AB, BC, CD and DA respectively.



To Prove: PQRS is a parallelogram and $\text{ar}(\text{||gm PQRS})$

$$= \frac{1}{2} \text{ar}(\text{||gm ABCD})$$

Construction: Join AC, BD and SQ.

Proof: As S and R are the midpoints of AD and CD. So, in $\triangle ADC$,

$$SR \parallel AC \quad [\text{By mid point theorem}]$$

Also, as P and Q are the midpoints of AB and BC. So, in $\triangle ABC$,

$$PQ \parallel AC$$

$$\therefore PQ \parallel AC \parallel SR$$

$$\therefore PQ \parallel SR$$

Similarly, we can prove $SP \parallel RQ$.

Thus PQRS is a parallelogram as its opposite sides are parallel since diagonals of a parallelogram bisect each other.

So in $\triangle ABD$,

O is the midpoint of AC and S is the midpoint of AD.

$$\therefore OS \parallel AB \quad [\text{By midpoint theorem}]$$

Similarly in $\triangle ABC$, we can prove that,

$$OQ \parallel AB$$

$$\text{i.e. } SQ \parallel AB$$

Thus, ABQS is a parallelogram.

$$\text{Now, } \text{ar}(\triangle SPQ) = \frac{1}{2} \text{ar}(\text{||gm ABQS}) \quad \dots (i)$$

$\left[\because \triangle SPQ \text{ and ||gm ABQS have the same base and lie between same parallel lines} \right]$

Similarly, we can prove that;

$$\text{ar}(\triangle SRQ) = \frac{1}{2} \text{ar}(\text{||gm SQCD}) \quad \dots (ii)$$

Adding (i) and (ii) we get

$$\text{ar}(\triangle SPQ) + \text{ar}(\triangle SRQ) = \frac{1}{2} [\text{ar}(\text{||gm ABQS}) + \text{ar}(\text{||gm SQCD})]$$

$$\therefore \text{ar}(\text{||gm PQRS}) = \frac{1}{2} \text{ar}(\text{||gm ABCD})$$

***** END *****