

## Trigonometric Identities Ex 6.2 Q8

## Answer:

Given:  $\cot \theta = \sqrt{3}$ 

We have to find the value of the expression  $\frac{\csc^2\theta + \cot^2\theta}{\csc^2\theta - \sec^2\theta}$ 

We know that,

$$\cot \theta = \sqrt{3} \Rightarrow \cot^2 \theta = 3$$

$$\csc^2 \theta = 1 + \cot^2 \theta = 1 + \left(\sqrt{3}\right)^2 = 4$$

$$\sec^2 \theta = \frac{1}{\cos^2 \theta} = \frac{1}{1 - \sin^2 \theta} = \frac{1}{1 - \frac{1}{\csc^2 \theta}} = \frac{1}{1 - \frac{1}{4}} = \frac{4}{3}$$

Therefore,

$$\frac{\csc^2\theta + \cot^2\theta}{\csc^2\theta - \sec^2\theta} = \frac{4+3}{4-\frac{4}{3}}$$
$$= \frac{21}{8}$$

Hence, the value of the given expression is 218.

Trigonometric Identities Ex 6.2 Q9

## Answer:

Given:  $3\cos\theta = 1$ 

We have to find the value of the expression  $\frac{6\sin^2\theta + \tan^2\theta}{4\cos\theta}$ 

We have.

$$3\cos\theta = 1$$

$$\Rightarrow \cos \theta = \frac{1}{3}$$

$$\sin\theta = \sqrt{1 - \cos^2\theta} = \sqrt{1 - \left(\frac{1}{3}\right)^2} = \frac{\sqrt{8}}{3}$$

$$\tan \theta = \frac{\sin \theta}{\cos \theta} = \frac{\frac{\sqrt{8}}{3}}{\frac{1}{3}} = \sqrt{8}$$

Therefore,

$$\frac{6\sin^2\theta + \tan^2\theta}{4\cos\theta} = \frac{6\times\left(\frac{\sqrt{8}}{3}\right)^2 + \left(\sqrt{8}\right)^2}{4\times\frac{1}{3}}$$

$$= 10$$

Hence, the value of the expression is 10.

Trigonometric Identities Ex 6.2 Q10

Answer:

Given:  $\sqrt{3} \tan \theta = 3 \sin \theta$ 

We have to find the value of  $\sin^2 \theta - \cos^2 \theta$ 

$$\sqrt{3} \tan \theta = 3 \sin \theta$$

$$\Rightarrow \sqrt{3} \frac{\sin \theta}{\cos \theta} = 3 \sin \theta$$

$$\Rightarrow \cos \theta = \frac{\sqrt{3}}{3} = \frac{1}{\sqrt{3}}$$

Therefore,

$$\sin^2 \theta - \cos^2 \theta = 1 - \cos^2 \theta - \cos^2 \theta$$
 (since,  $\sin^2 \theta + \cos^2 \theta = 1$ )  

$$= 1 - 2\cos^2 \theta$$
  

$$= 1 - 2 \times \left(\frac{1}{\sqrt{3}}\right)^2$$
  

$$= \frac{1}{3}$$

Hence, the value of the expression is  $\frac{1}{3}$ .

\*\*\*\*\*\*\*\*\* END \*\*\*\*\*\*\*