

Exercise 2A

Question 7:

We have

$$f(x) = x^2 - 5 = (x)^2 - (\sqrt{5})^2$$

$$= (x - \sqrt{5})(x + \sqrt{5})$$

$$[:: a^2 - b^2 = (a - b)(a + b)]$$

$$f(x) = 0 \Rightarrow (x - \sqrt{5})(x + \sqrt{5}) = 0$$

$$:: x - \sqrt{5} = 0 \text{ or } x + \sqrt{5} = 0$$

$$\Rightarrow x = \sqrt{5} \text{ or } x = -\sqrt{5}$$
So the zeros of  $f(x)$  are  $\sqrt{5}$  and  $-\sqrt{5}$ 

Sum of zeros =  $\sqrt{5} + (-\sqrt{5}) = 0 = -\frac{\text{Coeff. of } x}{\text{Coeff. of } x^2}$ 

Product of zeros =  $(\sqrt{5})(-\sqrt{5}) = -5 = \frac{-5}{1} = \frac{\text{Cons tant term}}{\text{Coeff. of } x^2}$ 

Question 8:

Let

$$f(x) = 8x^{2} - 4 = 4(2x^{2} - 1) = 4\left[\left(\sqrt{2}x\right)^{2} - 1^{2}\right]$$

$$= 4\left(\sqrt{2}x - 1\right)\left(\sqrt{2}x + 1\right)$$

$$f(x) = 0 \Rightarrow \left(\sqrt{2}x - 1\right)\left(\sqrt{2}x + 1\right) = 0$$

$$\therefore \sqrt{2}x - 1 = 0 \text{ or } \sqrt{2}x + 1 = 0$$

$$\therefore x = \frac{1}{\sqrt{2}} \text{ or } x = -\frac{1}{\sqrt{2}}$$
So, the zeros of  $f(x)$  are  $\frac{1}{\sqrt{2}}$  and  $-\frac{1}{\sqrt{2}}$ 

$$\text{Sum of zeros} = \left(\frac{1}{\sqrt{2}}\right) + \left(-\frac{1}{\sqrt{2}}\right) = 0 = \frac{0}{8} = \frac{\text{Coeff. of } x}{\text{Coeff. of } x^{2}}$$

$$\text{Product of zeros} = \left(\frac{1}{\sqrt{2}}\right)x\left(-\frac{1}{\sqrt{2}}\right) = -\frac{1}{2} = \frac{-4}{8}$$

$$= \frac{\text{constant term}}{\text{Coeff. of } x^{2}}$$

Question 9:

Let, 
$$f(x) = 5u^2 + 10u = 5u(u + 2)$$
  
 $f(x) = 0 \Rightarrow 5u(u + 2) = 0$   
∴  $u = 0$  or  $u + 2 = 0$   
⇒  $u = 0$  or  $u = -2$   
Sum of zeros =  $0 + (-2) = -2 = -\frac{10}{5}$   
 $= -\frac{Coeff. \text{ of } x}{Coeff. \text{ of } x^2}$   
Product of zero =  $0 \times (-2) = 0 = \frac{0}{5} = \frac{Cons \text{ tan t term}}{Coeff. \text{ of } x^2}$ 

\*\*\*\*\*\* END \*\*\*\*\*\*