



Arithmetic Progressions Ex 9.4 Q6

Answer :

Here, we are given three terms,

$$\text{First term } (a_1) = 8x + 4$$

$$\text{Second term } (a_2) = 6x - 2$$

$$\text{Third term } (a_3) = 2x + 7$$

We need to find the value of x for which these terms are in A.P. So, in an A.P. the difference of two adjacent terms is always constant. So, we get,

$$d = a_2 - a_1$$

$$d = (6x - 2) - (8x + 4)$$

$$d = 6x - 8x - 2 - 4$$

$$d = -2x - 6 \quad \dots\dots (1)$$

Also,

$$d = a_3 - a_2$$

$$d = (2x + 7) - (6x - 2)$$

$$d = 2x - 6x + 7 + 2$$

$$d = -4x + 9 \quad \dots\dots (2)$$

Now, on equating (1) and (2), we get,

$$-2x - 6 = -4x + 9$$

$$4x - 2x = 9 + 6$$

$$2x = 15$$

$$x = \frac{15}{2}$$

Therefore, for $x = \frac{15}{2}$, these three terms will form an A.P.

Arithmetic Progressions Ex 9.4 Q7

Answer :

Here, we are given three terms which are in A.P.,

$$\text{First term } (a_1) = x + 1$$

$$\text{Second term } (a_2) = 3x$$

$$\text{Third term } (a_3) = 4x + 2$$

We need to find the value of x . So, in an A.P. the difference of two adjacent terms is always constant.

So, we get,

$$d = a_2 - a_1$$

$$d = (3x) - (x + 1)$$

$$d = 3x - x - 1$$

$$d = 2x - 1 \quad \dots\dots (1)$$

Also,

$$d = a_3 - a_2$$

$$d = (4x + 2) - (3x)$$

$$d = 4x - 3x + 2$$

$$d = x + 2 \quad \dots\dots (2)$$

Now, on equating (1) and (2), we get,

$$2x - 1 = x + 2$$

$$2x - x = 2 + 1$$

$$x = 3$$

Therefore, for $x = 3$, these three terms will form an A.P.

Arithmetic Progressions Ex 9.4 Q8

Answer :

Here, we are given three terms and we need to show that they are in A.P.,

$$\text{First term } (a_1) = (a-b)^2$$

$$\text{Second term } (a_2) = (a^2 + b^2)$$

$$\text{Third term } (a_3) = (a+b)^2$$

So, in an A.P. the difference of two adjacent terms is always constant. So, to prove that these terms are in A.P. we find the common difference, we get,

$$d = a_2 - a_1$$

$$d = (a^2 + b^2) - (a-b)^2$$

$$d = a^2 + b^2 - (a^2 + b^2 - 2ab)$$

$$d = a^2 + b^2 - a^2 - b^2 + 2ab$$

$$d = 2ab \quad \dots\dots (1)$$

Also,

$$d = a_3 - a_2$$

$$d = (a+b)^2 - (a^2 + b^2)$$

$$d = a^2 + b^2 + 2ab - a^2 - b^2$$

$$d = 2ab \quad \dots\dots (2)$$

Now, since in equations (1) and (2) the values of d are equal, we can say that these terms are in A.P. with $2ab$ as the common difference.

Hence proved

***** END *****