

## Binomial Theorem Ex 18.2 Q34

We have, 
$$(1+2a)^4(2-a)^5$$
 Now, 
$$(1+2a)^4 = {}^4C_0 + {}^4C_12a + {}^4C_2(2a)^2 + {}^4C_3(2a)^3 + {}^4C_4(2a)^4$$
 and, 
$$(2-5)^5 = {}^5C_0 \times 2^5 + {}^5C_2 \times 2^4(-a) + {}^5C_2 \times 2^3(-a)^3 + {}^5C_3 \times 2^2(-a)^3 + {}^5C_4 \times 2(-a)^4 + {}^5C_5(-a)^5$$
 
$$= {}^5C_0 \times 2^5 - {}^5C_1 \times 2^4 \times a + {}^5C_2 \times 2^3 \times a^2 - {}^5C_3 \times 2^2 \times a^3 + {}^5C_4 \times 2 \times a^4 - {}^5C_5 \times a^5$$
 
$$(1+2a)^4(2-a)^5 = \left[ {}^4C_0 + {}^4C_12a + {}^4C_2(2a)^2 + {}^4C_3(2a)^3 + {}^4C_4(2a)^4 \right] \left[ {}^5C_0 \times 2^5 - {}^5C_1 \times 2^4 \times a + {}^5C_2 \times 2^3 \times a^2 - {}^5C_3 \times 2^2 \times a^3 + {}^5C_4 \times 2 \times a^4 - {}^5C_5 \times 2^5 \times a^5 \right]$$
 
$$(2-a)^5 \times (2-a)^5 \times (2$$

∴ Coefficients of a<sup>4</sup> = -438.

## Binomial Theorem Ex 18.2 Q35

$$\left(\sqrt{x} - \frac{k}{x^2}\right)^{10}$$

$$\binom{10}{2} \left(\sqrt{x}\right)^8 \left(-\frac{k}{x^2}\right)^2 = 405$$

$$45k^2 = 405$$

$$k^2 = 9$$

$$k = 3$$

Binomial Theorem Ex 18.2 Q36

$$(y^{1/2} + x^{1/3})^n$$

$$\binom{n}{n-2} (y^{1/2})^2 (x^{1/3})^{n-2}$$

$$\binom{n}{n-2} = 45$$

$$n(n-1) = 90$$

$$n^2 - 10n + 9n - 90$$

$$n(n-10) + 9(n-10) = 0$$

$$n = -9 \text{ or } 10$$

$$n \text{ cannot be negative. So, } n = 10$$

$$6thterm \binom{10}{5} (y^{1/2})^5 (x^{1/3})^5 = 252y^{\frac{5}{3}}x^{\frac{5}{3}}$$