



### Trigonometric Ratios Ex 5.1 Q32

**Answer :**

Given:

$$\operatorname{cosec} A = 2 \quad \dots\dots (1)$$

To find:

$$\frac{1}{\tan A} + \frac{\sin A}{1 + \cos A}$$

Now we know  $\operatorname{cosec} A$  is defined as below

$$\operatorname{cosec} A = \frac{1}{\sin A}$$

Therefore,

$$\sin A = \frac{1}{\operatorname{cosec} A}$$

Now by substituting the value of  $\operatorname{cosec} A$  from equation (1)

We get,

$$\sin A = \frac{1}{2} \quad \dots\dots (2)$$

Now by substituting the value of  $\sin A$  in the following identity of trigonometry

$$\sin^2 A + \cos^2 A = 1$$

We get,

$$\cos^2 A = 1 - \sin^2 A$$

$$= 1 - \left(\frac{1}{2}\right)^2$$

$$= 1 - \frac{1}{4}$$

Now by taking L.C.M we get

$$\cos^2 A = \frac{4-1}{4}$$

$$= \frac{3}{4}$$

Now by taking square root on both sides

We get,

$$\cos A = \sqrt{\frac{3}{4}}$$

$$= \frac{\sqrt{3}}{\sqrt{4}}$$

$$= \frac{\sqrt{3}}{2}$$

Therefore,

$$\cos A = \frac{\sqrt{3}}{2} \quad \dots\dots (3)$$

Now  $\tan A$  is defined as follows

$$\tan A = \frac{\sin A}{\cos A}$$

Now by substituting the value of  $\sin A$  and  $\cos A$  from equation (2) and (3) respectively we get,

$$\begin{aligned}\tan A &= \frac{\frac{1}{2}}{\frac{\sqrt{3}}{2}} \\ &= \frac{1}{2} \times \frac{2}{\sqrt{3}} \\ &= \frac{1}{\sqrt{3}}\end{aligned}$$

Therefore,

$$\tan A = \frac{1}{\sqrt{3}} \dots\dots (4)$$

Now by substituting the value of  $\sin A$ ,  $\cos A$  and  $\tan A$  from equation (2), (3) and (4) respectively we get,

$$\begin{aligned}\frac{1}{\tan A} + \frac{\sin A}{1 + \cos A} &= \frac{1}{\frac{1}{\sqrt{3}}} + \frac{\frac{1}{2}}{1 + \frac{\sqrt{3}}{2}} \\ &= \frac{\sqrt{3}}{1} + \frac{1}{2\left(1 + \frac{\sqrt{3}}{2}\right)}\end{aligned}$$

Now by taking L.C.M we get

$$\frac{1}{\tan A} + \frac{\sin A}{1 + \cos A} = \frac{\sqrt{3}}{1} + \frac{1}{2\left(\frac{2 + \sqrt{3}}{2}\right)}$$

Now 2 gets cancelled and we get

$$\frac{1}{\tan A} + \frac{\sin A}{1 + \cos A} = \frac{\sqrt{3}}{1} + \frac{1}{(2 + \sqrt{3})}$$

Now by taking L.C.M, we get,

$$\begin{aligned}\frac{1}{\tan A} + \frac{\sin A}{1 + \cos A} &= \frac{\sqrt{3} \times (2 + \sqrt{3})}{1 \times (2 + \sqrt{3})} + \frac{1}{(2 + \sqrt{3})} \\ &= \frac{\sqrt{3} \times (2 + \sqrt{3}) + 1}{(2 + \sqrt{3})}\end{aligned}$$

Now by opening the brackets in the numerator

We get,

$$\frac{1}{\tan A} + \frac{\sin A}{1 + \cos A} = \frac{2\sqrt{3} + \sqrt{3}\sqrt{3} + 1}{(2 + \sqrt{3})}$$

Since  $\sqrt{3}\sqrt{3} = 3$

Therefore,

$$\begin{aligned}\frac{1}{\tan A} + \frac{\sin A}{1 + \cos A} &= \frac{2\sqrt{3} + 3 + 1}{(2 + \sqrt{3})} \\ &= \frac{2\sqrt{3} + 4}{(2 + \sqrt{3})}\end{aligned}$$

Now by taking 2 common

We get,

$$\begin{aligned}\frac{1}{\tan A} + \frac{\sin A}{1 + \cos A} &= \frac{2(\sqrt{3} + 2)}{(2 + \sqrt{3})} \\ &= \frac{2(2 + \sqrt{3})}{(2 + \sqrt{3})}\end{aligned}$$

Now as  $(2 + \sqrt{3})$  is present in both numerator as well as denominator, it gets cancelled

Therefore,

$$\frac{1}{\tan A} + \frac{\sin A}{1 + \cos A} = 2$$

\*\*\*\*\* END \*\*\*\*\*