



Exercise 11A

Question 15:

Two circles with centres A and B, having radii 5 cm and 3 cm touch each other internally.

The perpendicular bisector of AB meets the bigger circle in P and Q.


Join AP.

Let PQ intersect AB at L.

Then, $AB = (5 - 3) \text{ cm} = 2 \text{ cm}$

Since PQ is the perpendicular bisector of AB, we have

$$\begin{aligned} AL &= \frac{1}{2} \times AB \\ &= \left(\frac{1}{2} \times 2 \right) \text{ cm} = 1 \text{ cm} \end{aligned}$$



Now, in right angle $\triangle PLA$

$$\therefore AP^2 = AL^2 + PL^2$$


$$\begin{aligned} \Rightarrow PL &= \sqrt{AP^2 - AL^2} \text{ cm} \\ &= \sqrt{(25 - 1)} \text{ cm} = \sqrt{24} \text{ cm} = 2\sqrt{6} \text{ cm} \end{aligned}$$

$$\therefore PQ = (2 \times PL) = (2 \times 2\sqrt{6}) \text{ cm} = 4\sqrt{6} \text{ cm}$$

$$\therefore \text{the length of } PQ = 4\sqrt{6} \text{ cm}$$

Question 16:

Given: AB is a chord of a circle with centre O. AB is produced to C such that $BC = OB$. Also, CO is joined to meet the circle in D. $\angle ACD = y^\circ$ and $\angle AOD = x^\circ$.



To Prove : $x = 3y$
 Proof : $OB = BC$ [Given]
 $\therefore \angle BOC = \angle BCO = y^\circ$ [isosceles triangle]
 Ext. $\angle OBA = \angle BOC + \angle BCO = (2y)^\circ$
 Again, $OA = OB$ [radii of same circle]
 $\therefore \angle OAB = \angle OBA = (2y)^\circ$ [isosceles triangle]
 Ext. $\angle AOD = \angle OAC + \angle ACO$
 $= \angle OAB + \angle BCO = 3y^\circ$
 $\therefore x^\circ = 3y^\circ$ [$\because \angle AOD = x$ (given)]

***** END *****