

Exercise 9B

## Question 9:

Let  $f_1$  and  $f_2$  be the frequencies of class intervals 0 - 10 and 40 - 50

$$f_1 + 5 + 9 + 12 + f_2 + 3 + 2 = 40$$
  
 $\Rightarrow f_1 + f_2 = 9$ 

Median is 32.5 which lies in 30 - 40, so the median class is 30 - 40

$$I = 30, h = 10, f = 12, N = 40 \text{ and } C = f_1 + 5 + 9 = (f_1 + 14)$$

Now, median = I + 
$$\left[h \times \frac{\left(\frac{N}{2} - c\right)}{f}\right]$$
  
 $\Rightarrow 32.5 = \left[30 + \left(10 \times \frac{20 - f_1 - 14}{12}\right)\right]$   
 $\Rightarrow = \left[30 + \left(10 \times \frac{6 - f_1}{12}\right)\right]$   
 $\Rightarrow = \left[30 + \left(\frac{30 - 5f_1}{6}\right)\right]$   
 $\frac{30 - 5f_1}{6} = 2.5$   
 $30 - 5f_1 = 15$   
 $5f_1 = 15 \Rightarrow f_1 = 3$   
 $f_1 = 3$  and  $f_2 = (9 - 3) = 6$ 

Question 10: The given series is of inclusive form. Converting it into exclusive form and preparing the cumulative frequency table, we get

Class	Frequency	C.F
	fi	
18.5 - 25.5	35	35
25.5 - 32.5	96	131
32.5 - 39.5	68	199
39.5 - 46.5	102	301
46.5 - 53.5	35	336
53.5 - 60.5	4	340
	$\sum f_i = N = 340$	

$$N = 340 \Rightarrow \frac{N}{2} = 170$$

The cumulative frequency just greater than 170 is 199 and the corresponding class is 32.5 - 39.5.

.. Median class is 32.5 - 39.5

 $_{\odot}$  I = 32.5, h = 7, f = 68, c = C.F. of preceding median class = 131

Median 
$$m_e = I + \left[ h \times \frac{\left(\frac{N}{2} - c\right)}{f} \right] = 32.5 + \left[ 7 \times \frac{(170 - 131)}{68} \right]$$
  
= 32.5 + 4 = 36.5

Hence median is 36.5 years

\*\*\*\*\*\*\* END \*\*\*\*\*\*\*