



### Trigonometric Identities Ex 6.1 Q45

**Answer :**

In the given question, we need to prove  $\frac{\tan^2 A}{1 + \tan^2 A} + \frac{\cot^2 A}{1 + \cot^2 A} = 1$ .

Here, we will first solve the LHS.

Now, using  $\tan \theta = \frac{\sin \theta}{\cos \theta}$  and  $\cot \theta = \frac{\cos \theta}{\sin \theta}$ , we get

$$\begin{aligned} \frac{\tan^2 A}{1 + \tan^2 A} + \frac{\cot^2 A}{1 + \cot^2 A} &= \frac{\left(\frac{\sin^2 A}{\cos^2 A}\right)}{\left(1 + \frac{\sin^2 A}{\cos^2 A}\right)} + \frac{\left(\frac{\cos^2 A}{\sin^2 A}\right)}{\left(1 + \frac{\cos^2 A}{\sin^2 A}\right)} \\ &= \frac{\left(\frac{\sin^2 A}{\cos^2 A}\right)}{\left(\frac{\cos^2 A + \sin^2 A}{\cos^2 A}\right)} + \frac{\left(\frac{\cos^2 A}{\sin^2 A}\right)}{\left(\frac{\sin^2 A + \cos^2 A}{\sin^2 A}\right)} \\ &= \frac{\left(\frac{\sin^2 A}{\cos^2 A}\right)}{\left(\frac{1}{\cos^2 A}\right)} + \frac{\left(\frac{\cos^2 A}{\sin^2 A}\right)}{\left(\frac{1}{\sin^2 A}\right)} \quad \left(\text{Using } \sin^2 \theta + \cos^2 \theta = 1\right) \end{aligned}$$

On further solving by taking the reciprocal of the denominator, we get,

$$\begin{aligned} \frac{\left(\frac{\sin^2 A}{\cos^2 A}\right)}{\left(\frac{1}{\cos^2 A}\right)} + \frac{\left(\frac{\cos^2 A}{\sin^2 A}\right)}{\left(\frac{1}{\sin^2 A}\right)} &= \left(\frac{\sin^2 A}{\cos^2 A}\right)\left(\frac{\cos^2 A}{1}\right) + \left(\frac{\cos^2 A}{\sin^2 A}\right)\left(\frac{\sin^2 A}{1}\right) \\ &= \sin^2 A + \cos^2 A \quad \left(\text{Using } \sin^2 \theta + \cos^2 \theta = 1\right) \\ &= 1 \end{aligned}$$

Hence proved.

### Trigonometric Identities Ex 6.1 Q46

**Answer :**

In the given question, we need to prove  $\frac{\cot A - \cos A}{\cot A + \cos A} = \frac{\operatorname{cosec} A - 1}{\operatorname{cosec} A + 1}$

Here, we will first solve the LHS.

Now, using  $\cot \theta = \frac{\cos \theta}{\sin \theta}$ , we get

$$\begin{aligned}\frac{\cot A - \cos A}{\cot A + \cos A} &= \frac{\left(\frac{\cos A}{\sin A} - \cos A\right)}{\left(\frac{\cos A}{\sin A} + \cos A\right)} \\ &= \frac{\left(\frac{\cos A - \cos A \sin A}{\sin A}\right)}{\left(\frac{\cos A + \cos A \sin A}{\sin A}\right)}\end{aligned}$$

On further solving by taking the reciprocal of the denominator, we get,

$$\begin{aligned}\frac{\left(\frac{\cos A - \cos A \sin A}{\sin A}\right)}{\left(\frac{\cos A + \cos A \sin A}{\sin A}\right)} &= \left(\frac{\cos A - \cos A \sin A}{\sin A}\right) \left(\frac{\sin A}{\cos A + \cos A \sin A}\right) \\ &= \left(\frac{\cos A - \cos A \sin A}{\cos A + \cos A \sin A}\right)\end{aligned}$$

Now, taking  $\cos A \sin A$  common from both the numerator and the denominator, we get

$$\begin{aligned}\left(\frac{\cos A - \cos A \sin A}{\cos A + \cos A \sin A}\right) &= \frac{\cos A \sin A \left(\frac{1}{\sin A} - 1\right)}{\cos A \sin A \left(\frac{1}{\sin A} + 1\right)} \\ &= \frac{\left(\frac{1}{\sin A} - 1\right)}{\left(\frac{1}{\sin A} + 1\right)} \\ &= \frac{\operatorname{cosec} A - 1}{\operatorname{cosec} A + 1} \quad \left(\text{Using } \frac{1}{\sin \theta} = \operatorname{cosec} \theta\right)\end{aligned}$$

Hence proved.

\*\*\*\*\* END \*\*\*\*\*