

EXERCISE - 2.2

Question-1

Let $A = \{1, 2, 3, \dots, 14\}$. Define a relation R from A to A by $B = \{(x, y): 3x - y = 0, \text{ where } x, y \in A\}$. Write down its domain, codomain and range.

Ans

The relation R from A to A is given as

$$R = \{(x, y): 3x - y = 0, \text{ where } x, y \in A\}$$

i.e.,
$$R = \{(x, y): 3x = y, \text{ where } x, y \in A\}$$

$$R = \{(1, 3), (2, 6), (3, 9), (4, 12)\}$$

The domain of R is the set of all first elements of the ordered pairs in the relation.

 \therefore Domain of R = {1, 2, 3, 4}

The whole set A is the codomain of the relation R.

 \therefore Codomain of $R = A = \{1, 2, 3, ..., 14\}$

The range of R is the set of all second elements of the ordered pairs in the relation.

 \therefore Range of R = {3, 6, 9, 12}

Question-2

Define a relation R on the set **N** of natural numbers by $R = \{(x, y): y = x + 5, x \text{ is a natural number less than 4}; x, y \in \mathbf{N}\}$. Depict this relationship using roster form. Write down the domain and the range.

Ans.

 $R = \{(x, y): y = x + 5, x \text{ is a natural number less than } 4, x, y \in \mathbb{N}\}$

The natural numbers less than 4 are 1, 2, and 3.

$$R = \{(1, 6), (2, 7), (3, 8)\}$$

The domain of R is the set of all first elements of the ordered pairs in the relation.

... Domain of $R = \{1, 2, 3\}$

The range of R is the set of all second elements of the ordered pairs in the relation.

 \therefore Range of R = {6, 7, 8}

Question-3

 $A = \{1, 2, 3, 5\}$ and $B = \{4, 6, 9\}$. Define a relation R from A to B by $B = \{(x, y): \text{ the difference between } x \text{ and } y \text{ is odd}; x \in A, y \in B\}$. Write R in roster form.

Ans.

$$A = \{1, 2, 3, 5\}$$
 and $B = \{4, 6, 9\}$

 $R = \{(x, y): \text{ the difference between } x \text{ and } y \text{ is odd}; x \in A, y \in B\}$


$$R = \{(1, 4), (1, 6), (2, 9), (3, 4), (3, 6), (5, 4), (5, 6)\}$$

Question-4

The given figure shows a relationship between the sets P and Q. write this relation

(i) in set-builder form (ii) in roster form.

What is its domain and range?



Ans

According to the given figure, $P = \{5, 6, 7\}, Q = \{3, 4, 5\}$

(i)
$$R = \{(x, y): y = x - 2; x \in P\}$$
 or $R = \{(x, y): y = x - 2 \text{ for } x = 5, 6, 7\}$

(ii)
$$R = \{(5, 3), (6, 4), (7, 5)\}$$

Domain of $R = \{5, 6, 7\}$

Range of $R = \{3, 4, 5\}$

Question-5

Let A = {1, 2, 3, 4, 6}. Let R be the relation on A defined by

 $\{(a, b): a, b \in A, b \text{ is exactly divisible by } a\}.$

- (i) Write R in roster form
- (ii) Find the domain of R
- (iii) Find the range of R.

Ans.

 $A = \{1, 2, 3, 4, 6\}, R = \{(a, b): a, b \in A, b \text{ is exactly divisible by } a\}$

$$(i) \ R = \{(1, 1), (1, 2), (1, 3), (1, 4), (1, 6), (2, 2), (2, 4), (2, 6), (3, 3), (3, 6), (4, 4), (6, 6)\}$$

- (ii) Domain of $R = \{1, 2, 3, 4, 6\}$
- (iii) Range of R = {1, 2, 3, 4, 6}

Question-6

Determine the domain and range of the relation R defined by $R = \{(x, x + 5): x \in \{0, 1, 2, 3, 4, 5\}\}.$

Ans.

$$R = \{(x, x + 5): x \in \{0, 1, 2, 3, 4, 5\}\}$$

$$\therefore$$
 R = {(0, 5), (1, 6), (2, 7), (3, 8), (4, 9), (5, 10)}

 \therefore Domain of R = {0, 1, 2, 3, 4, 5}

Range of $R = \{5, 6, 7, 8, 9, 10\}$

Question-7

Write the relation R = $\{(x, x^3): x \text{ is a prime number less than 10}\}$ in roster form. Ans.

 $R = \{(x, x^3): x \text{ is a prime number less than } 10\}$

The prime numbers less than 10 are 2, 3, 5, and 7.

$$R = \{(2, 8), (3, 27), (5, 125), (7, 343)\}$$

Question-8

Let $A = \{x, y, z\}$ and $B = \{1, 2\}$. Find the number of relations from A to B

It is given that $A = \{x, y, z\}$ and $B = \{1, 2\}$.

$$\therefore$$
 A × B = {(x, 1), (x, 2), (y, 1), (y, 2), (z, 1), (z, 2)}

Since $n(A \times B) = 6$, the number of subsets of $A \times B$ is 2^6 .

Therefore, the number of relations from A to B is 26.

Question-9

Let R be the relation on **Z** defined by $R = \{(a, b): a, b \in \mathbf{Z}, a - b \text{ is an integer}\}$. Find the domain and range of R.

Ans.

 $R = \{(a, b): a, b \in \mathbb{Z}, a - b \text{ is an integer}\}$

It is known that the difference between any two integers is always an integer.

 \therefore Domain of R = Z

Range of R = Z

******* END ******