

Differentiation Ex 11.2 Q52 Consider

$$y = \frac{x^2 \left(1 - x^2\right)^3}{\cos 2x}$$

Differentiating it with respect to x and applying the chain and product rule, we get

$$\frac{dy}{dx} = \frac{\cos 2x \frac{d}{dx} x^2 (1-x^2)^3 - x^2 (1-x^2)^3 \frac{d}{dx} \cos 2x}{\cos^2 2x}$$

$$= \frac{\cos 2x \left[x^2 \frac{d}{dx} (1-x^2)^3 + (1-x^2)^3 \frac{d}{dx} x^2 - x^2 (1-x^2)^3 (-2\sin 2x) \right]}{\cos^2 2x}$$

$$= \frac{\cos 2x \left[-6x^3 (1-x^2)^2 + (1-x^2)^3 2x + 2x^2 (1-x^2)^3 \sin 2x \right]}{\cos^2 2x}$$

$$= \frac{2x (1-x^2)^2}{\cos 2x} - \frac{6x^3 (1-x^2)^2}{\cos 2x} + \frac{2x^2 (1-x^2)^3 \sin 2x}{\cos^2 2x}$$

$$= 2x (1-x^2) \sec 2x \left\{ 1 - 4x^2 + x (1-x^2) \tan 2x \right\}$$
Therefore,

$$\frac{dy}{dx} = 2x(1-x^2)\sec 2x\{1-4x^2+x(1-x^2)\tan 2x\}$$

Differentiation Ex 11.2 Q53

Consider

$$y = \log(3x+2) - x^2 \log(2x-1)$$

Differentiating it with respect to x and applying the chain and product rule, we get

$$\frac{dy}{dx} = \frac{d}{dx} \left[\log(3x+2) - x^2 \log(2x-1) \right]$$

$$\frac{dy}{dx} = \frac{3}{3x+2} - \left[x^2 \frac{d}{dx} \log(2x-1) + \log(2x-1) \frac{d}{dx} x^2 \right]$$

$$\frac{dy}{dx} = \frac{3}{3x+2} - \left(\frac{2x^2}{2x-1} + 2x \log(2x-1) \right)$$

$$\frac{dy}{dx} = \frac{3}{3x+2} - \frac{2x^2}{2x-1} - 2x \log(2x-1)$$

Therefore.

$$\frac{dy}{dx} = \frac{3}{3x+2} - \frac{2x^2}{2x-1} - 2x \log(2x-1)$$

Differentiation Ex 11.2 Q54

Consider

$$y = e^{ax} \sec x \tan 2x$$

Differentiating it with respect to x and applying the chain and product rule, we get

$$\frac{dy}{dx} = \frac{d}{dx} \left(e^{xx} \sec x \tan 2x \right)$$

$$= e^{ax} \frac{d}{dx} \sec x \tan 2x + \sec x \tan 2x \frac{d}{dx} e^{xx}$$

$$= e^{ax} \left[\sec x \tan x \tan 2x + \left(2 + 2 \tan^2 2x \right) \sec x \right] + ae^{xx} \sec x \tan 2x$$

$$= e^{ax} \left[\sec x \tan x \tan 2x + 2 \sec x + 2 \tan^2 2x \sec x \right] + ae^{xx} \sec x \tan 2x$$

$$= ae^{ax} \sec x \tan 2x + e^{xx} \sec x \tan x \tan 2x + e^{xx} \sec x \left(2 + 2 \tan^2 2x \right)$$

$$\frac{dy}{dx} = e^{ax} \sec x \left\{ a \tan 2x + \tan x \tan 2x + 2 \sec^2 2x \right\}$$
Therefore,
$$\frac{dy}{dx} = e^{ax} \sec x \left\{ a \tan 2x + \tan x \tan 2x + 2 \sec^2 2x \right\}$$

Consider

$$y = \log(\cos x^2)$$

Differentiating it with respect to x and applying the chain and product rule, we get

$$\frac{dy}{dx} = \frac{d}{dx} \log(\cos x^2)$$

$$= \frac{-2x \sin x^2}{\cos x^2}$$

$$\frac{dy}{dx} = -2x \tan x^2$$

Therefore,

$$\frac{dy}{dx} = -2x \tan x^2$$

Differentiation Ex 11.2 Q56

Consider

$$y = \cos(\log x)^2$$

Differentiating it with respect to x and applying the chain and product rule, we get

$$\frac{dy}{dx} = \frac{d}{dx}\cos(\log x)^2$$

$$= -\sin(\log x)^2 \frac{d}{dx}(\log x)^2$$

$$= -\sin(\log x)^2 \frac{2\log x}{x}$$

$$= \frac{dy}{dx} = \frac{-2\log x \sin(\log x)^2}{x}$$
efore,

Therefore.

$$\frac{dy}{dx} = \frac{-2 \log x \sin (\log x)^2}{x}$$

********** END ********