



Chapter 9 Continuity Ex 9.2 Q10

Let $f(x) = \sin|x|$

This function f is defined for every real number and f can be written as the composition of two functions as,

$f = g \circ h$, where $g(x) = |x|$ and $h(x) = \sin x$

$$[\because (g \circ h)(x) = g(h(x)) = g(\sin x) = |\sin x| = f(x)]$$

It has to be proved first that $g(x) = |x|$ and $h(x) = \sin x$ are continuous functions.

$g(x) = |x|$ can be written as

$$g(x) = \begin{cases} -x, & \text{if } x < 0 \\ x, & \text{if } x \geq 0 \end{cases}$$

Clearly, g is defined for all real numbers.

Let c be a real number.

Case I:

If $c < 0$, then $g(c) = -c$ and $\lim_{x \rightarrow c} g(x) = \lim_{x \rightarrow c} (-x) = -c$

$$\therefore \lim_{x \rightarrow c} g(x) = g(c)$$

Therefore, g is continuous at all points x , such that $x < 0$

Case II:

If $c > 0$, then $g(c) = c$ and $\lim_{x \rightarrow c} g(x) = \lim_{x \rightarrow c} x = c$

$$\therefore \lim_{x \rightarrow c} g(x) = g(c)$$

Therefore, g is continuous at all points x , such that $x > 0$

Case III:

If $c = 0$, then $g(c) = g(0) = 0$

$$\lim_{x \rightarrow 0^-} g(x) = \lim_{x \rightarrow 0^-} (-x) = 0$$

$$\lim_{x \rightarrow 0^+} g(x) = \lim_{x \rightarrow 0^+} (x) = 0$$

$$\therefore \lim_{x \rightarrow 0^-} g(x) = \lim_{x \rightarrow 0^+} g(x) = g(0)$$

Therefore, g is continuous at $x = 0$

From the above three observations, it can be concluded that g is continuous at all points.

$$h(x) = \sin x$$

It is evident that $h(x) = \sin x$ is defined for every real number.

Let c be a real number. Put $x = c + k$

If $x \rightarrow c$, then $k \rightarrow 0$

$$h(c) = \sin c$$

$$h(c) = \sin c$$

$$\lim_{x \rightarrow c} h(x) = \lim_{x \rightarrow c} \sin x$$

$$= \lim_{k \rightarrow 0} \sin(c + k)$$

$$= \lim_{k \rightarrow 0} [\sin c \cos k + \cos c \sin k]$$

$$= \lim_{k \rightarrow 0} (\sin c \cos k) + \lim_{k \rightarrow 0} (\cos c \sin k)$$

$$= \sin c \cos 0 + \cos c \sin 0$$

$$= \sin c + 0$$

$$= \sin c$$

$$\therefore \lim_{x \rightarrow c} h(x) = g(c)$$

Therefore, h is a continuous function.

It is known that for real valued functions g and h , such that $(g \circ h)$ is defined at c , if g is continuous at c and if h is continuous at $g(c)$, then $(g \circ h)$ is continuous at c .

Therefore, $f(x) = (g \circ h)(x) = g(h(x)) = g(\sin x) = |\sin x|$ is a continuous function.

Chapter 9 Continuity Ex 9.2 Q11

When $x < 0$, we have,

$$f(x) = \frac{\sin x}{x}$$

We know that the $\sin x$ and the identity function x are continuous for $x < 0$.

So, the quotient function $f(x) = \frac{\sin x}{x}$ is continuous for $x < 0$.

When $x > 0$, we have,

$f(x) = x + 1$, which is a polynomial of degree 1. So, $f(x)$ is continuous for $x > 0$

Now, consider the point $x = 0$.

$$f(0) = 0 + 1 = 1.$$

$$\text{LHL} = \lim_{x \rightarrow 0^-} f(x) = \lim_{h \rightarrow 0} f(0 - h) = \lim_{h \rightarrow 0} \frac{\sin(-h)}{-h} = \lim_{h \rightarrow 0} \frac{-\sin h}{-h} = 1$$

$$\text{RHL} = \lim_{x \rightarrow 0^+} f(x) = \lim_{h \rightarrow 0} f(0 + h) = \lim_{h \rightarrow 0} h + 1 = 1$$

$$\text{Thus, LHL} = \text{RHL} = f(0) = 1$$

So, $f(x)$ is continuous at $x = 0$.

Hence, $f(x)$ is continuous everywhere

Chapter 9 Continuity Ex 9.2 Q12

The given function is $g(x) = x - [x]$

It is evident that g is defined at all integral points.

Let n be an integer.

Then,

$$g(n) = n - [n] = n - n = 0$$

The left hand limit of f at $x = n$ is,

$$\lim_{x \rightarrow n^-} g(x) = \lim_{x \rightarrow n^-} (x - [x]) = \lim_{x \rightarrow n^-} (x) - \lim_{x \rightarrow n^-} [x] = n - (n-1) = 1$$

The right hand limit of f at $x = n$ is,

$$\lim_{x \rightarrow n^+} g(x) = \lim_{x \rightarrow n^+} (x - [x]) = \lim_{x \rightarrow n^+} (x) - \lim_{x \rightarrow n^+} [x] = n - n = 0$$

It is observed that the left and right hand limits of f at $x = n$ do not coincide.

Therefore, f is not continuous at $x = n$

Hence, g is discontinuous at all integral points

Chapter 9 Continuity Ex 9.2 Q13

It is known that if g and h are two continuous functions, then

$g + h$, $g - h$, and $g.h$ are also continuous.

It has to be proved first that $g(x) = \sin x$ and $h(x) = \cos x$ are continuous functions.

Let $g(x) = \sin x$

It is evident that $g(x) = \sin x$ is defined for every real number.

Let c be a real number. Put $x = c + h$

If $x \rightarrow c$, then $h \rightarrow 0$

$$g(c) = \sin c$$

$$\begin{aligned}\lim_{x \rightarrow c} g(x) &= \lim_{x \rightarrow c} \sin x \\ &= \lim_{h \rightarrow 0} \sin(c + h) \\ &= \lim_{h \rightarrow 0} [\sin c \cos h + \cos c \sin h] \\ &= \lim_{h \rightarrow 0} (\sin c \cos h) + \lim_{h \rightarrow 0} (\cos c \sin h) \\ &= \sin c \cos 0 + \cos c \sin 0 \\ &= \sin c + 0 \\ &= \sin c\end{aligned}$$

$$\therefore \lim_{x \rightarrow c} g(x) = g(c)$$

Therefore, g is a continuous function.

Let $h(x) = \cos x$

It is evident that $h(x) = \cos x$ is defined for every real number.

Let c be a real number. Put $x = c + h$

If $x \rightarrow c$, then $h \rightarrow 0$

$h(c) = \cos c$

$$\begin{aligned}\lim_{x \rightarrow c} h(x) &= \lim_{x \rightarrow c} \cos x \\ &= \lim_{h \rightarrow 0} \cos(c + h) \\ &= \lim_{h \rightarrow 0} [\cos c \cos h - \sin c \sin h] \\ &= \lim_{h \rightarrow 0} \cos c \cos h - \lim_{h \rightarrow 0} \sin c \sin h \\ &= \cos c \cos 0 - \sin c \sin 0 \\ &= \cos c \times 1 - \sin c \times 0 \\ &= \cos c\end{aligned}$$

$$\therefore \lim_{x \rightarrow c} h(x) = h(c)$$

Therefore, h is a continuous function.

Therefore, it can be concluded that

- (a) $f(x) = g(x) + h(x) = \sin x + \cos x$ is a continuous function
- (b) $f(x) = g(x) - h(x) = \sin x - \cos x$ is a continuous function
- (c) $f(x) = g(x) \times h(x) = \sin x \times \cos x$ is a continuous function

***** END *****