

Exercise 14A

Q1

Answer:

Given: l | m

t is a transversal.

$$\angle 5 = 70^{\circ}$$

$$\angle 5 = \angle 3 = 70^{\circ}$$
 (alternate interior angles)

$$\angle 5 + \angle 8 = 180^{\circ}$$
 (linear pair)


or
$$70^{\circ} + \angle 8 = 180^{\circ}$$

$$\angle 8 = 110^{\circ}$$

$$\angle 1 = \angle 3 = 70^{\circ}$$
 (vertically opposite angles)


$$\angle 3 + \angle 4 = 180^{\circ}$$
 (linear pair)

or
$$\angle 4 = 180 - \angle 3 = 180 - 70 = 110^{\circ}$$



Q2

Answer:



Given: $l \parallel m$

t is a transversal.

$$\angle 1: \angle 2 = 5:7$$

Let the angles measure 5x and 7x.

$$\angle 1 + \angle 2 = 180^{\circ}$$

(linear pair)

$$\therefore 5x + 7x = 180$$

or
$$12x = 180$$

or
$$x = 15$$

$$\therefore \angle 1 = 5x = 5(15) = 75^{\circ}$$

and
$$\angle 2 = 7x = 7(15) = 105^{\circ}$$

$$\angle 2 + \angle 3 = 180^{\circ}$$

(linear pair)

$$\angle 3 = 180 - 105 = 75^{\circ}$$

 $\angle 3 + \angle 6 = 180$ supplementary)

(interior angles on the same side of the transversal are

$$\angle 6 = 180 - \angle 3 = 105^{\circ}$$

and $\angle 6 = \angle 8 = 105^{\circ}$ (vertically opposite angles)
 $\therefore \angle 1 = 75^{\circ}$
 $\angle 2 = 105^{\circ}$
 $\angle 3 = 75^{\circ}$
 $\angle 8 = 105^{\circ}$
Q3
Answer:

Given: l | m t is a transversal. Let:

$$\angle 1 = (2x - 8)^{\circ}$$

 $\angle 2 = (3x - 7)^{\circ}$

We know that the consecutive interior angles are supplementary.

$$\therefore \angle 1 + \angle 2 = 180^{\circ}$$
or $(2x-8) + (3x-7) = 180$
or $5x - 15 = 180$
or $5x = 195$
or $x = 39$

$$\angle 1 = (2x-8) = (2 \times 39 - 8) = 70^{\circ}$$

$$\angle 2 = (3x-7) = (3 \times 39 - 7) = 110^{\circ}$$

********** END ********