



## EXERCISE - 2.3

### Question-1

Which of the following relations are functions? Give reasons. If it is a function, determine its domain and range.

(i)  $\{(2, 1), (5, 1), (8, 1), (11, 1), (14, 1), (17, 1)\}$

(ii)  $\{(2, 1), (4, 2), (6, 3), (8, 4), (10, 5), (12, 6), (14, 7)\}$

(iii)  $\{(1, 3), (1, 5), (2, 5)\}$

Ans.

(i)  $\{(2, 1), (5, 1), (8, 1), (11, 1), (14, 1), (17, 1)\}$

Since 2, 5, 8, 11, 14, and 17 are the elements of the domain of the given relation having their unique images, this relation is a function.

Here, domain =  $\{2, 5, 8, 11, 14, 17\}$  and range =  $\{1\}$

(ii)  $\{(2, 1), (4, 2), (6, 3), (8, 4), (10, 5), (12, 6), (14, 7)\}$

Since 2, 4, 6, 8, 10, 12, and 14 are the elements of the domain of the given relation having their unique images, this relation is a function.

Here, domain =  $\{2, 4, 6, 8, 10, 12, 14\}$  and range =  $\{1, 2, 3, 4, 5, 6, 7\}$

(iii)  $\{(1, 3), (1, 5), (2, 5)\}$

Since the same first element i.e., 1 corresponds to two different images i.e., 3 and 5, this relation is not a function.

### Question-2

Find the domain and range of the following real function:

(i)  $f(x) = -|x|$  (ii)  $f(x) = \sqrt{9 - x^2}$

Ans.

(i)  $f(x) = -|x|, x \in \mathbf{R}$

We know that  $|x| = \begin{cases} x, & x \geq 0 \\ -x, & x < 0 \end{cases}$

$\therefore f(x) = -|x| = \begin{cases} -x, & x \geq 0 \\ x, & x < 0 \end{cases}$

Since  $f(x)$  is defined for  $x \in \mathbf{R}$ , the domain of  $f$  is  $\mathbf{R}$ .

It can be observed that the range of  $f(x) = -|x|$  is all real numbers except positive real numbers.

$\therefore$  The range of  $f$  is  $(-\infty, 0]$ .

(ii)  $f(x) = \sqrt{9 - x^2}$

Since  $\sqrt{9 - x^2}$  is defined for all real numbers that are greater than or equal to  $-3$  and less than or equal to  $3$ , the domain of  $f(x)$  is  $\{x : -3 \leq x \leq 3\}$  or  $[-3, 3]$ .

For any value of  $x$  such that  $-3 \leq x \leq 3$ , the value of  $f(x)$  will lie between 0 and 3.

$\therefore$  The range of  $f(x)$  is  $\{x : 0 \leq x \leq 3\}$  or  $[0, 3]$ .

Question-3

A function  $f$  is defined by  $f(x) = 2x - 5$ . Write down the values of

(i)  $f(0)$ , (ii)  $f(7)$ , (iii)  $f(-3)$

Ans.

The given function is  $f(x) = 2x - 5$ .

Therefore,

$$(i) f(0) = 2 \times 0 - 5 = 0 - 5 = -5$$

$$(ii) f(7) = 2 \times 7 - 5 = 14 - 5 = 9$$

$$(iii) f(-3) = 2 \times (-3) - 5 = -6 - 5 = -11$$

Question-4

The function ' $t$ ' which maps temperature in degree Celsius into temperature in degree Fahrenheit is defined by  $t(C) = \frac{9C}{5} + 32$ .

Find (i)  $t(0)$  (ii)  $t(28)$  (iii)  $t(-10)$  (iv) The value of  $C$ , when  $t(C) = 212$

Ans.

The given function is  $t(C) = \frac{9C}{5} + 32$ .

Therefore,

$$(i) t(0) = \frac{9 \times 0}{5} + 32 = 0 + 32 = 32$$

$$(ii) t(28) = \frac{9 \times 28}{5} + 32 = \frac{252 + 160}{5} = \frac{412}{5}$$

$$(iii) t(-10) = \frac{9 \times (-10)}{5} + 32 = 9 \times (-2) + 32 = -18 + 32 = 14$$

(iv) It is given that  $t(C) = 212$

$$\therefore 212 = \frac{9C}{5} + 32$$

$$\Rightarrow \frac{9C}{5} = 212 - 32$$

$$\Rightarrow \frac{9C}{5} = 180$$

$$\Rightarrow 9C = 180 \times 5$$

$$\Rightarrow C = \frac{180 \times 5}{9} = 100$$

Thus, the value of  $t$ , when  $t(C) = 212$ , is 100.

Question-5

Find the range of each of the following functions.

(i)  $f(x) = 2 - 3x, x \in \mathbf{R}, x > 0.$

(ii)  $f(x) = x^2 + 2, x, \text{ is a real number.}$

(iii)  $f(x) = x, x \text{ is a real number}$

Ans.

(i)  $f(x) = 2 - 3x, x \in \mathbf{R}, x > 0$

The values of  $f(x)$  for various values of real numbers  $x > 0$  can be written in the tabular form as

$x$	0.01	0.1	0.9	1	2	2.5	4	5	...
$f(x)$	1.97	1.7	-0.7	-1	-4	-5.5	-10	-13	...

Thus, it can be clearly observed that the range of  $f$  is the set of all real numbers less than 2.

i.e., range of  $f = (-\infty, 2)$

**Alter:**

Let  $x > 0$

$$\Rightarrow 3x > 0$$

$$\Rightarrow 2 - 3x < 2$$

$$\Rightarrow f(x) < 2$$

$\therefore$  Range of  $f = (-\infty, 2)$

(ii)  $f(x) = x^2 + 2, x, \text{ is a real number}$

The values of  $f(x)$  for various values of real numbers  $x$  can be written in the tabular form as

$x$	0	$\pm 0.3$	$\pm 0.8$	$\pm 1$	$\pm 2$	$\pm 3$	...
$f(x)$	2	2.09	2.64	3	6	11	.....

Thus, it can be clearly observed that the range of  $f$  is the set of all real numbers greater than 2.

i.e., range of  $f = [2, \infty)$

**Alter:**

Let  $x$  be any real number.

Accordingly,

$$x^2 \geq 0$$

$$\Rightarrow x^2 + 2 \geq 0 + 2$$

$$\Rightarrow x^2 + 2 \geq 2$$

$$\Rightarrow f(x) \geq 2$$

$\therefore$  Range of  $f = [2, \infty)$

(iii)  $f(x) = x, x \text{ is a real number}$

It is clear that the range of  $f$  is the set of all real numbers.

$\therefore$  Range of  $f = \mathbf{R}$

\*\*\*\*\* END \*\*\*\*\*