



WIKIPEDIA  
The Free Encyclopedia

Main page  
Contents  
Featured content  
Current events  
Random article  
Donate to Wikipedia  
Wikipedia store

Interaction  
Help  
About Wikipedia  
Community portal  
Recent changes  
Contact page

Tools  
What links here  
Related changes  
Upload file  
Special pages  
Permanent link  
Page information  
Wikidata item  
Cite this page

Print/export  
Create a book  
Download as PDF  
Printable version

Languages  
العربية  
Català  
Čeština  
Deutsch  
Español  
Français  
한국어  
Italiano  
עברית  
Nederlands  
日本語  
Polski  
Português  
Русский  
Српски / srpski  
Українська  
Tiếng Việt  
中文

Edit links

[Create account](#) [Log in](#)

Article [Talk](#)

[Read](#) [Edit](#) [View history](#)

# Miller–Rabin primality test

From Wikipedia, the free encyclopedia

The **Miller–Rabin primality test** or **Rabin–Miller primality test** is a [primality test](#): an [algorithm](#) which determines whether a given number is [prime](#), similar to the [Fermat primality test](#) and the [Solovay–Strassen primality test](#). Its original version, due to [Gary L. Miller](#), is [deterministic](#), but the determinism relies on the unproven [Extended Riemann hypothesis](#);<sup>[1]</sup> [Michael O. Rabin](#) modified it to obtain an unconditional [probabilistic algorithm](#).<sup>[2]</sup>

## Contents

- [Concepts](#)
- [Example](#)
- [Algorithm and running time](#)
- [Accuracy of the test](#)
- [Deterministic variants of the test](#)
- [Notes](#)
- [External links](#)

## Concepts [\[ edit \]](#)

Just like the Fermat and Solovay–Strassen tests, the Miller–Rabin test relies on an equality or set of equalities that hold true for prime values, then checks whether or not they hold for a number that we want to test for primality.

First, a [lemma](#) about square [roots of unity](#) in the [finite field](#)  $\mathbf{Z}/p\mathbf{Z}$ , where  $p$  is prime and  $p > 2$ . Certainly 1 and  $-1$  always yield 1 when squared modulo  $p$ ; call these [trivial square roots](#) of 1. There are no *nontrivial* square roots of 1 modulo  $p$  (a special case of the result that, in a field, a [polynomial](#) has no more zeroes than its degree). To show this, suppose that  $x$  is a square root of 1 modulo  $p$ . Then:

$$\begin{aligned}x^2 &\equiv 1 \pmod{p} \\(x-1)(x+1) &\equiv 0 \pmod{p}.\end{aligned}$$

In other words, prime  $p$  divides the product  $(x-1)(x+1)$ . By [Euclid's lemma](#) it divides one of the factors  $x-1$  or  $x+1$ , implying that  $x$  is congruent to either 1 or  $-1$  modulo  $p$ .

Now, let  $n$  be prime with  $n > 2$ . It follows that  $n-1$  is even and we can write it as  $2^s \cdot d$ , where  $s$  and  $d$  are positive integers ( $d$  is odd). For each  $a$  in  $(\mathbf{Z}/n\mathbf{Z})^*$ , either

$$a^d \equiv 1 \pmod{n}$$

or

$$a^{2^r \cdot d} \equiv -1 \pmod{n}$$

for some  $0 \leq r \leq s-1$ .

To show that one of these must be true, recall [Fermat's little theorem](#), that for a prime number  $n$ :

$$a^{n-1} \equiv 1 \pmod{n}.$$

By the lemma above, if we keep taking square roots of  $a^{n-1}$ , we will get either 1 or  $-1$ . If we get  $-1$  then the second equality holds and it is done. If we never get  $-1$ , then when we have taken out every power of 2, we are left with the first equality.

The Miller–Rabin primality test is based on the [contrapositive](#) of the above claim. That is, if we can find an  $a$  such that

$$a^d \not\equiv 1 \pmod{n}$$

and

$$a^{2^r d} \not\equiv -1 \pmod{n}$$

for all  $0 \leq r \leq s-1$ , then  $n$  is not prime. We call  $a$  a [witness](#) for the compositeness of  $n$  (sometimes misleadingly

called a *strong witness*, although it is a certain proof of this fact). Otherwise  $a$  is called a *strong liar*, and  $n$  is a [strong probable prime](#) to base  $a$ . The term "strong liar" refers to the case where  $n$  is composite but nevertheless the equations hold as they would for a prime.

Every odd composite  $n$  has many witnesses  $a$ , however, no simple way of generating such an  $a$  is known. The solution is to make the test [probabilistic](#): we choose a non-zero  $a$  in  $\mathbb{Z}/n\mathbb{Z}$  randomly, and check whether or not it is a witness for the compositeness of  $n$ . If  $n$  is composite, most of the choices for  $a$  will be witnesses, and the test will detect  $n$  as composite with high probability. There is, nevertheless, a small chance that we are unlucky and hit an  $a$  which is a strong liar for  $n$ . We may reduce the probability of such error by repeating the test for several independently chosen  $a$ .

For testing large numbers, it is common to choose random bases  $a$ , as, a priori, we don't know the distribution of witnesses and liars among the numbers  $1, 2, \dots, n-1$ . In particular, Arnault [3] gave a 397-digit composite number for which all bases  $a$  less than 307 are strong liars. As expected this number was reported to be prime by the [Maple](#) `isprime()` function, which implemented the Miller–Rabin test by checking the specific bases 2,3,5,7, and 11. However, selection of a few specific small bases can guarantee identification of composites for  $n$  less than some maximum determined by said bases. This maximum is generally quite large compared to the bases. As random bases lack such determinism for small  $n$ , specific bases are better in some circumstances.

## Example [\[ edit \]](#)

Suppose we wish to determine if  $n = 221$  is prime. We write  $n - 1 = 220$  as  $2^2 \cdot 55$ , so that we have  $s = 2$  and  $d = 55$ . We randomly select a number  $a$  such that  $0 < a < n$ , say  $a = 174$ . We proceed to compute:

- $a^{2^0 \cdot d} \bmod n = 174^{55} \bmod 221 = 47 \neq 1, n-1$
- $a^{2^1 \cdot d} \bmod n = 174^{110} \bmod 221 = 220 = n-1$ .

Since  $220 \equiv -1 \bmod n$ , either 221 is prime, or 174 is a strong liar for 221. We try another random  $a$ , this time choosing  $a = 137$ :

- $a^{2^0 \cdot d} \bmod n = 137^{55} \bmod 221 = 188 \neq 1, n-1$
- $a^{2^1 \cdot d} \bmod n = 137^{110} \bmod 221 = 205 \neq n-1$ .

Hence 137 is a witness for the compositeness of 221, and 174 was in fact a strong liar. Note that this tells us nothing about the factors of 221 (which are 13 and 17). However, the example with 341 in the next section shows how these calculations can sometimes produce a factor of  $n$ .

## Algorithm and running time [\[ edit \]](#)

The algorithm can be written in [pseudocode](#) as follows:

```
Input:  $n > 3$ , an odd integer to be tested for primality;
Input:  $k$ , a parameter that determines the accuracy of the test
Output: composite if  $n$  is composite, otherwise probably prime
write  $n - 1$  as  $2^s \cdot d$  with  $d$  odd by factoring powers of 2 from  $n - 1$ 
WitnessLoop: repeat  $k$  times:
    pick a random integer  $a$  in the range  $[2, n - 2]$ 
     $x \leftarrow a^d \bmod n$ 
    if  $x = 1$  or  $x = n - 1$  then do next WitnessLoop
    repeat  $s - 1$  times:
         $x \leftarrow x^2 \bmod n$ 
        if  $x = 1$  then return composite
        if  $x = n - 1$  then do next WitnessLoop
    return composite
return probably prime
```

Using [modular exponentiation](#) by [repeated squaring](#), the running time of this algorithm is  $O(k \log^3 n)$ , where  $k$  is the number of different values of  $a$  that we test; thus this is an efficient, polynomial-time algorithm. [FFT](#)-based multiplication can push the running time down to  $O(k \log^2 n \log \log n \log \log \log n) = \tilde{O}(k \log^2 n)$ .

If we insert [Greatest common divisor](#) (GCD) calculations into the above algorithm, we can sometimes obtain a factor of  $n$  instead of merely determining that  $n$  is composite. In particular, if  $n$  is a probable prime base  $a$  but not a strong probable prime base  $a$ , then either  $\text{GCD}((a^d \bmod n) - 1, n)$  or (for some  $r$  in the above range)  $\text{GCD}((a^{2^r \cdot d} \bmod n) - 1, n)$  will produce a (not necessarily prime) factor of  $n$ ; see page 1402 of [4]. If factoring is a goal, these GCDs can be inserted into the above algorithm at little additional computational cost.

For example, consider  $n = 341$ . We have  $n - 1 = 85 \cdot 4$ . Then  $2^{85} \bmod 341 = 32$ . This tells us that  $n$  is not a

strong probable prime base 2, so we know  $n$  is composite. If we take a GCD at this stage, we can get a factor of 341:  $\text{GCD}((2^{85} \bmod 341) - 1, 341) = 31$ . This works because 341 is a pseudoprime base 2, but is not a [strong pseudoprime](#) base 2.

In the case that the algorithm returns "composite" because  $x = 1$ , it has also discovered that  $d2^r$  is (an odd multiple of) the [order](#) of  $a$ —a fact which can (as in [Shor's algorithm](#)) be used to factorize  $n$ , since  $n$  then divides

$$a^{d2^r} - 1 = (a^{d2^{r-1}} - 1)(a^{d2^{r-1}} + 1)$$

but not either factor by itself. The reason Miller–Rabin does *not* yield a probabilistic [factorization](#) algorithm is that if

$$a^{n-1} \not\equiv 1 \pmod{n}$$

(i.e.,  $n$  is not a [pseudoprime](#) to base  $a$ ) then no such information is obtained about the period of  $a$ , and the second "return composite" is taken.

## Accuracy of the test [\[edit\]](#)

The more bases  $a$  we test, the better the accuracy of the test. It can be shown that for any odd composite  $n$ , at least 3/4 of the bases  $a$  are witnesses for the compositeness of  $n$ .<sup>[2][5]</sup> The Miller–Rabin test is strictly stronger than the Solovay–Strassen primality test in the sense that for every composite  $n$ , the set of strong liars for  $n$  is a subset of the set of [Euler liars](#) for  $n$ , and for many  $n$ , the subset is proper. If  $n$  is composite then the Miller–Rabin primality test declares  $n$  probably prime with a probability at most  $4^{-k}$ . On the other hand, the Solovay–Strassen primality test declares  $n$  probably prime with a probability at most  $2^{-k}$ .

It is important to note that in many common applications of this algorithm, we are not interested in the error bound described above. The above error bound is the probability of a composite number being declared as a probable prime after  $k$  rounds of testing. We are often instead interested in the probability that, after passing  $k$  rounds of testing, the number being tested is actually a composite number. Formally, if we call the event of declaring  $n$  a probable prime after  $k$  rounds of Miller–Rabin  $Y_k$ , and we call the event that  $n$  is composite  $X$  (and denote the event that  $n$  is prime  $\overline{X}$ ), then the above bound gives us  $P(Y_k|X)$ , whereas we are interested in  $P(X|Y_k)$ . [Bayes' theorem](#) gives us a way to relate these two conditional probabilities, namely

$$P(X|Y_k) = \frac{P(Y_k|X)P(X)}{P(Y_k|X)P(X) + P(Y_k|\overline{X})P(\overline{X})}$$

This tells us that the probability that we are often interested in is related not just to the  $4^{-k}$  bound above, but also probabilities related to the density of prime numbers in the region near  $n$ .

In addition, for large values of  $n$ , on average the probability that a composite number is declared *probably prime* is significantly smaller than  $4^{-k}$ . [Damgård](#), Landrock and [Pomerance](#)<sup>[6]</sup> compute some explicit bounds and provide a method to make a reasonable selection for  $k$  for a desired error bound. Such bounds can, for example, be used to *generate* probable primes; however, they should not be used to *verify* primes with unknown origin, since in [cryptographic](#) applications an adversary might try to send you a [pseudoprime](#) in a place where a prime number is required. In such cases, only the error bound of  $4^{-k}$  can be relied upon.

## Deterministic variants of the test [\[edit\]](#)

The Miller–Rabin algorithm can be made deterministic by trying all possible  $a$  below a certain limit. The problem in general is to set the limit so that the test is still reliable.

If the tested number  $n$  is composite, the strong liars  $a$  coprime to  $n$  are contained in a proper [subgroup](#) of the group  $(\mathbf{Z}/n\mathbf{Z})^*$ , which means that if we test all  $a$  from a set which [generates](#)  $(\mathbf{Z}/n\mathbf{Z})^*$ , one of them must be a witness for the compositeness of  $n$ . Assuming the truth of the [generalized Riemann hypothesis](#) (GRH), it is known that the group is generated by its elements smaller than  $O((\log n)^2)$ , which was already noted by Miller.<sup>[1]</sup> The constant involved in the [Big O notation](#) was reduced to 2 by [Eric Bach](#).<sup>[7]</sup> This leads to the following conditional primality testing algorithm:

```
Input:  $n > 1$ , an odd integer to test for primality.
Output: composite if  $n$  is composite, otherwise prime
write  $n-1$  as  $2^s \cdot d$  by factoring powers of 2 from  $n-1$ 
repeat for all  $a \in [2, \min(n-1, \lceil 2(\ln n)^2 \rceil)]$ :
    if  $a^d \not\equiv 1 \pmod{n}$  and  $a^{2^r \cdot d} \not\equiv -1 \pmod{n}$  for all  $r \in [0, s-1]$ 
        then return composite
return prime
```

The running time of the algorithm is  $\tilde{O}((\log n)^4)$  (with FFT-based multiplication). The full power of the generalized Riemann hypothesis is not needed to ensure the correctness of the test: as we deal with subgroups of even [index](#), it suffices to assume the validity of GRH for [quadratic Dirichlet characters](#).<sup>[5]</sup>

The Miller test (the algorithm above) is not used in practice. For most purposes, proper use of the probabilistic Miller-Rabin test or the [Baillie-PSW primality test](#) gives sufficient confidence while running much faster. It is also slower in practice than commonly used proof methods such as [APR-CL](#) and [ECPP](#) which give results that do not rely on unproven assumptions. For theoretical purposes requiring a deterministic polynomial time algorithm, it was superseded by the [AKS primality test](#), which also does not rely on unproven assumptions.

Note that Miller-Rabin pseudoprimes are called [strong pseudoprimes](#).

When the number  $n$  to be tested is small, trying all  $a < 2(\ln n)^2$  is not necessary, as much smaller sets of potential witnesses are known to suffice. For example, Pomerance, Selfridge and Wagstaff<sup>[8]</sup> and Jaeschke<sup>[9]</sup> have verified that

- if  $n < 2,047$ , it is enough to test  $a = 2$ ;
- if  $n < 1,373,653$ , it is enough to test  $a = 2$  and  $3$ ;
- if  $n < 9,080,191$ , it is enough to test  $a = 31$  and  $73$ ;
- if  $n < 25,326,001$ , it is enough to test  $a = 2, 3$ , and  $5$ ;
- if  $n < 4,759,123,141$ , it is enough to test  $a = 2, 7$ , and  $61$ ;
- if  $n < 1,122,004,669,633$ , it is enough to test  $a = 2, 13, 23$ , and  $1662803$ ;
- if  $n < 2,152,302,898,747$ , it is enough to test  $a = 2, 3, 5, 7$ , and  $11$ ;
- if  $n < 3,474,749,660,383$ , it is enough to test  $a = 2, 3, 5, 7, 11$ , and  $13$ ;
- if  $n < 341,550,071,728,321$ , it is enough to test  $a = 2, 3, 5, 7, 11, 13$ , and  $17$ .

Using the work of Feitsma and Galway enumerating all base 2 pseudoprimes in 2010, this was extended (see [A014233](#)), with the first result later shown using different methods in Jiang and Deng.<sup>[10]</sup>

- if  $n < 3,825,123,056,546,413,051$ , it is enough to test  $a = 2, 3, 5, 7, 11, 13, 17, 19$ , and  $23$ .
- if  $n < 2^{64}$ , it is enough to test  $a = 2, 3, 5, 7, 11, 13, 17, 19, 23, 29, 31$ , and  $37$ .

Sorenson and Webster<sup>[11]</sup> verify the above and calculate precise results for these larger than 64-bit results:

- if  $n < 318,665,857,834,031,151,167,461$ , it is enough to test  $a = 2, 3, 5, 7, 11, 13, 17, 19, 23, 29, 31$ , and  $37$ .
- if  $n < 3,317,044,064,679,887,385,961,981$ , it is enough to test  $a = 2, 3, 5, 7, 11, 13, 17, 19, 23, 29, 31, 37$ , and  $41$ .

Other criteria of this sort, often more efficient (fewer bases required) than those shown above, exist<sup>[12][13][14][15]</sup> and these results give very fast deterministic primality tests for numbers in the appropriate range without any assumptions.

There is a small list of potential witnesses for every possible input size (at most  $n$  values for  $n$ -bit numbers). However, no finite set of bases is sufficient for all composite numbers. Alford, Granville, and Pomerance have shown that there exist infinitely many composite numbers  $n$  whose smallest compositeness witness is at least  $(\ln n)^{1/(3 \ln \ln n)}$ .<sup>[16]</sup> They also argue heuristically that the smallest number  $w$  such that every composite number below  $n$  has a compositeness witness less than  $w$  should be of order  $\Theta(\log n \log \log n)$ .


## Notes [\[edit\]](#)

- <sup>a b</sup> Miller, Gary L. (1976), "Riemann's Hypothesis and Tests for Primality", *Journal of Computer and System Sciences* **13** (3): 300–317, doi:[10.1145/800116.803773](#)
- <sup>a b</sup> Rabin, Michael O. (1980), "Probabilistic algorithm for testing primality", *Journal of Number Theory* **12** (1): 128–138, doi:[10.1016/0022-314X\(80\)90084-0](#)
- <sup>a</sup> F. Amault (August 1995). "Constructing Carmichael Numbers Which Are Strong Pseudoprimes to Several Bases". *Journal of Symbolic Computation* **20** (2): 151–161. doi:[10.1006/jsc.1995.1042](#).
- <sup>a</sup> Robert Baillie; Samuel S. Wagstaff, Jr. (October 1980). "Lucas Pseudoprimes" (PDF). *Mathematics of Computation* **35** (152): 1391–1417. doi:[10.1090/S0025-5718-1980-0583518-6](#). MR [583518](#).
- <sup>a b</sup> Schoof, René (2004), "Four primality testing algorithms", *Algorithmic Number Theory: Lattices, Number Fields, Curves and Cryptography* (PDF), Cambridge University Press, ISBN 0-521-80854-5
- <sup>a</sup> Damgård, I.; Landrock, P. & Pomerance, C. (1993), "Average case error estimates for the strong probable prime test" (PDF), *Mathematics of Computation* **61** (203): 177–194, doi:[10.2307/2152945](#)
- <sup>a</sup> Bach, Eric (1990), "Explicit bounds for primality testing and related problems", *Mathematics of Computation* **55** (191): 355–380, doi:[10.2307/2008811](#)
- <sup>a</sup> Carl Pomerance; John L. Selfridge, Samuel S. Wagstaff, Jr. (July 1980). "The pseudoprimes to  $25 \cdot 10^9$ " (PDF). *Mathematics of Computation* **35** (151): 1003–1026. doi:[10.1090/S0025-5718-1980-0572872-7](#).

9. <sup>^</sup> Jaeschke, Gerhard (1993), "On strong pseudoprimes to several bases", *Mathematics of Computation* **61** (204): 915–926, doi:10.2307/2153262
10. <sup>^</sup> Jiang, Yupeng; Deng, Yingpu (2014). "Strong pseudoprimes to the first eight prime bases". *Mathematics of Computation* **83** (290): 2915–2924. doi:10.1090/S0025-5718-2014-02830-5
11. <sup>^</sup> Sorenson, Jonathan; Webster, Jonathan (2015). "Strong Pseudoprimes to Twelve Prime Bases". *arXiv:1509.00864*  [math.NT] .
12. <sup>^</sup> *The Primes Page*
13. <sup>^</sup> Zhang, Zhenxiang & Tang, Min (2003), "Finding strong pseudoprimes to several bases. II", *Mathematics of Computation* **72** (44): 2085–2097, doi:10.1090/S0025-5718-03-01545-X
14. <sup>^</sup> "Sloane's A014233 : Smallest odd number for which Miller-Rabin primality test on bases  $\leq n$ -th prime does not reveal compositeness ", *The On-Line Encyclopedia of Integer Sequences*. OEIS Foundation.
15. <sup>^</sup> *SPRP Records*
16. <sup>^</sup> Alford, W. R.; Granville, A.; Pomerance, C. (1994), "On the difficulty of finding reliable witnesses"  (PDF), *Lecture Notes in Computer Science* (Springer-Verlag) **877**: 1–16, doi:10.1007/3-540-58691-1\_36

## External links [ edit ]

- Weisstein, Eric W., "Rabin-Miller Strong Pseudoprime Test" , *MathWorld*.
- Interactive Online Implementation of the Deterministic Variant  (stepping through the algorithm step-by-step)
- Applet (German)
- Miller-Rabin primality test in C#
- Miller-Rabin primality test in JavaScript using arbitrary precision arithmetic



The Wikibook *Algorithm Implementation* has a page on the topic of: **Primality testing**

<span>v</span> · <span>t</span> · <span>e</span>	Number-theoretic algorithms
<b>Primality tests</b>	AKS test · APR test · Baillie–PSW · ECPP test · Elliptic curve · Pocklington · Fermat · Lucas · Lucas–Lehmer · Lucas–Lehmer–Riesel · Proth's theorem · Pépin's · Quadratic Frobenius test · Solovay–Strassen · <b>Miller–Rabin</b>
<b>Prime-generating</b>	Sieve of Atkin · Sieve of Eratosthenes · Sieve of Sundaram · Wheel factorization
<b>Integer factorization</b>	Continued fraction (CFRAC) · Dixon's · Lenstra elliptic curve (ECM) · Euler's · Pollard's rho · <i>p</i> − 1 · <i>p</i> + 1 · Quadratic sieve (QS) · General number field sieve (GNFS) · <i>Special number field sieve</i> (SNFS) · Rational sieve · Fermat's · Shanks' square forms · Trial division · Shor's
<b>Multiplication</b>	Ancient Egyptian · Long · Karatsuba · Toom–Cook · Schönhage–Strassen · Fürer's
<b>Discrete logarithm</b>	Baby-step giant-step · Pollard rho · Pollard kangaroo · Pohlig–Hellman · Index calculus · Function field sieve
<b>Greatest common divisor</b>	Binary · Euclidean · Extended Euclidean · Lehmer's
<b>Modular square root</b>	Cipolla · Pocklington's · Tonelli–Shanks
<b>Other algorithms</b>	Chakravala · Cornacchia · Integer relation · Integer square root · Modular exponentiation · Schoof's
<i>Italics</i> indicate that algorithm is for numbers of special forms · Smallcaps indicate a deterministic algorithm	

Categories: Primality tests | Finite fields

This page was last modified on 4 September 2015, at 14:57.

Text is available under the  Creative Commons Attribution-ShareAlike License; additional terms may apply. By using this site, you agree to the  Terms of Use and  Privacy Policy. Wikipedia® is a registered trademark of the  Wikimedia Foundation, Inc., a non-profit organization.

Privacy policy  About Wikipedia  Disclaimers  Contact Wikipedia  Developers  Mobile view

