





WIKIPEDIA  
The Free Encyclopedia

Main page  
Contents  
Featured content  
Current events  
Random article  
Donate to Wikipedia  
Wikipedia store

Interaction  
Help  
About Wikipedia  
Community portal  
Recent changes  
Contact page

Tools  
What links here  
Related changes  
Upload file  
Special pages  
Permanent link  
Page information  
Wikidata item  
Cite this page

Print/export  
Create a book  
Download as PDF  
Printable version

Languages   
Deutsch  
Français  
Polski  Edit links

[Create account](#) [Log in](#)

Article [Talk](#)

[Read](#)

[Edit](#)

[More](#) ▾



# Neville's algorithm

From Wikipedia, the free encyclopedia

In mathematics, **Neville's algorithm** is an algorithm used for [polynomial interpolation](#) that was derived by the mathematician [Eric Harold Neville](#). Given  $n + 1$  points, there is a unique polynomial of degree  $\leq n$  which goes through the given points. Neville's algorithm evaluates this polynomial.

Neville's algorithm is based on the [Newton form](#) of the interpolating polynomial and the recursion relation for the [divided differences](#). It is similar to Aitken's algorithm (named after [Alexander Aitken](#)), which is nowadays not used.

## Contents

[\[hide\]](#)

- 1 The algorithm
- 2 Application to numerical differentiation
- 3 References
- 4 External links

## The algorithm [\[edit\]](#)

Given a set of  $n+1$  data points  $(x_i, y_i)$  where no two  $x_i$  are the same, the interpolating polynomial is the polynomial  $p$  of degree at most  $n$  with the property

$$p(x_i) = y_i \text{ for all } i = 0, \dots, n$$

This polynomial exists and it is unique. Neville's algorithm evaluates the polynomial at some point  $x$ .

Let  $p_{i,j}$  denote the polynomial of degree  $j - i$  which goes through the points  $(x_k, y_k)$  for  $k = i, i + 1, \dots, j$ . The  $p_{i,j}$  satisfy the recurrence relation

$$\begin{aligned} p_{i,i}(x) &= y_i, & 0 \leq i \leq n, \\ p_{i,j}(x) &= \frac{(x_j - x)p_{i,j-1}(x) + (x - x_i)p_{i+1,j}(x)}{x_j - x_i}, & 0 \leq i < j \leq n. \end{aligned}$$

This recurrence can calculate  $p_{0,n}(x)$ , which is the value being sought. This is Neville's algorithm.

For instance, for  $n = 4$ , one can use the recurrence to fill the triangular tableau below from the left to the right.

$$\begin{array}{ccccccc} p_{0,0}(x) & = & y_0 & & & & \\ & & & p_{0,1}(x) & & & \\ p_{1,1}(x) & = & y_1 & & p_{0,2}(x) & & \\ & & & p_{1,2}(x) & & p_{0,3}(x) & \\ p_{2,2}(x) & = & y_2 & & p_{1,3}(x) & & p_{0,4}(x) \\ & & & p_{2,3}(x) & & p_{1,4}(x) & \\ p_{3,3}(x) & = & y_3 & & p_{2,4}(x) & & \\ & & & p_{3,4}(x) & & & \\ p_{4,4}(x) & = & y_4 & & & & \end{array}$$

This process yields  $p_{0,4}(x)$ , the value of the polynomial going through the  $n + 1$  data points  $(x_i, y_i)$  at the point  $x$ .

This algorithm needs  $O(n^2)$  floating point operations.

The derivative of the polynomial can be obtained in the same manner, i.e:




$$\begin{aligned} p'_{i,i}(x) &= 0, & 0 \leq i \leq n, \\ p'_{i,j}(x) &= \frac{(x_j - x)p'_{i,j-1}(x) - p_{i,j-1}(x) + (x - x_i)p'_{i+1,j}(x) + p_{i+1,j}(x)}{x_j - x_i}, & 0 \leq i < j \leq n. \end{aligned}$$

## Application to numerical differentiation [\[edit\]](#)

Lyness and Moler showed in 1966 that using undetermined coefficients for the polynomials in Neville's algorithm,

one can compute the Maclaurin expansion of the final interpolating polynomial, which yields numerical approximations for the derivatives of the function at the origin. While "this process requires more arithmetic operations than is required in finite difference methods", "the choice of points for function evaluation is not restricted in any way". They also show that their method can be applied directly to the solution of linear systems of the Vandermonde type.

## References [\[edit\]](#)

- Press, William; Saul Teukolsky; William Vetterling; Brian Flannery (1992). "[§3.1 Polynomial Interpolation and Extrapolation \(encrypted\)](#)"  (PDF). *Numerical Recipes in C. The Art of Scientific Computing* (2nd edition ed.). Cambridge University Press. doi:[10.2277/0521431085](#) . ISBN 978-0-521-43108-8. (link is bad)
- J. N. Lyness and C.B. Moler, Van Der Monde Systems and Numerical Differentiation, *Numerische Mathematik* 8 (1966) 458-464 (doi: [10.1007/BF02166671](#) )

## External links [\[edit\]](#)

- Weisstein, Eric W., "Neville's Algorithm" , *MathWorld*.
- Module for Neville Interpolation by John H. Mathews 
- Java Code by Behzad Torkian 

Categories: [Polynomials](#) | [Interpolation](#)

This page was last modified on 14 July 2015, at 14:45.

Text is available under the [Creative Commons Attribution-ShareAlike License](#); additional terms may apply. By using this site, you agree to the [Terms of Use](#) and [Privacy Policy](#). Wikipedia® is a registered trademark of the [Wikimedia Foundation, Inc.](#), a non-profit organization.

[Privacy policy](#) [About Wikipedia](#) [Disclaimers](#) [Contact Wikipedia](#) [Developers](#) [Mobile view](#)

