



WIKIPEDIA
The Free Encyclopedia

[Main page](#)
[Contents](#)
[Featured content](#)
[Current events](#)
[Random article](#)
[Donate to Wikipedia](#)
[Wikipedia store](#)

Interaction
[Help](#)
[About Wikipedia](#)
[Community portal](#)
[Recent changes](#)
[Contact page](#)

Tools
[What links here](#)
[Related changes](#)
[Upload file](#)
[Special pages](#)
[Permanent link](#)
[Page information](#)
[Wikidata item](#)
[Cite this page](#)

Print/export
[Create a book](#)
[Download as PDF](#)
[Printable version](#)


Languages 
[Français](#)
[Nederlands](#)
[Português](#)
[Русский](#)
 [Edit links](#)

[Create account](#) [Log in](#)

Article [Talk](#)

[Read](#) [Edit](#)

[More](#) ▾



Muller's method

From Wikipedia, the free encyclopedia

Muller's method is a [root-finding algorithm](#), a [numerical](#) method for solving equations of the form $f(x) = 0$. It was first presented by [David E. Muller](#) in 1956.

Muller's method is based on the [secant method](#), which constructs at every iteration a line through two points on the graph of f . Instead, Muller's method uses three points, constructs the [parabola](#) through these three points, and takes the intersection of the [x-axis](#) with the parabola to be the next approximation.

Contents [hide]

- [1 Recurrence relation](#)
- [2 Speed of convergence](#)
- [3 Generalizations and related methods](#)
- [4 References](#)
- [5 External links](#)

Recurrence relation [\[edit\]](#)

Muller's method is a recursive method which generates an approximation of the [root](#) ξ of f at each iteration. Starting with the three initial values x_0 , x_1 and x_2 , the first iteration calculates the first approximation x_1 , the second iteration calculates the second approximation x_2 , the third iteration calculates the third approximation x_3 , etc. Hence the k^{th} iteration generates approximation x_k . Each iteration takes as input the last three generated approximations and the value of f at these approximations. Hence the k^{th} iteration takes as input the values x_{k-1} , x_{k-2} and x_{k-3} and the function values $f(x_{k-1})$, $f(x_{k-2})$ and $f(x_{k-3})$. The approximation x_k is calculated as follows.

A parabola $y_k(x)$ is constructed which goes through the three points $(x_{k-1}, f(x_{k-1}))$, $(x_{k-2}, f(x_{k-2}))$ and $(x_{k-3}, f(x_{k-3}))$. When written in the [Newton form](#), $y_k(x)$ is

$$y_k(x) = f(x_{k-1}) + (x - x_{k-1})f[x_{k-1}, x_{k-2}] + (x - x_{k-1})(x - x_{k-2})f[x_{k-1}, x_{k-2}, x_{k-3}],$$

where $f[x_{k-1}, x_{k-2}]$ and $f[x_{k-1}, x_{k-2}, x_{k-3}]$ denote [divided differences](#). This can be rewritten as

$$y_k(x) = f(x_{k-1}) + w(x - x_{k-1}) + f[x_{k-1}, x_{k-2}, x_{k-3}](x - x_{k-1})^2$$

where

$$w = f[x_{k-1}, x_{k-2}] + f[x_{k-1}, x_{k-3}] - f[x_{k-2}, x_{k-3}].$$

The next iterate x_k is now given as the solution closest to x_{k-1} of the quadratic equation $y_k(x) = 0$. This yields the [recurrence relation](#)

$$x_k = x_{k-1} - \frac{2f(x_{k-1})}{w \pm \sqrt{w^2 - 4f(x_{k-1})f[x_{k-1}, x_{k-2}, x_{k-3}]}.$$

In this formula, the sign should be chosen such that the denominator is as large as possible in magnitude. We do not use the standard formula for solving [quadratic equations](#) because that may lead to [loss of significance](#).

Note that x_k can be complex, even if the previous iterates were all real. This is in contrast with other root-finding algorithms like the [secant method](#), [Sidi's generalized secant method](#) or [Newton's method](#), whose iterates will remain real if one starts with real numbers. Having complex iterates can be an advantage (if one is looking for complex roots) or a disadvantage (if it is known that all roots are real), depending on the problem.

Speed of convergence [\[edit\]](#)

The [order of convergence](#) of Muller's method is approximately 1.84. This can be compared with 1.62 for the [secant method](#) and 2 for [Newton's method](#). So, the secant method makes less progress per iteration than Muller's method and Newton's method makes more progress.

More precisely, if ξ denotes a single root of f (so $f(\xi) = 0$ and $f'(\xi) \neq 0$), f is three times continuously differentiable, and the initial guesses x_0 , x_1 , and x_2 are taken sufficiently close to ξ , then the iterates satisfy

$$\lim_{k \rightarrow \infty} \frac{|x_k - \xi|}{|x_{k-1} - \xi|^\mu} = \left| \frac{f'''(\xi)}{6f'(\xi)} \right|^{(\mu-1)/2},$$

where $\mu \approx 1.84$ is the positive solution of $x^3 - x^2 - x - 1 = 0$.

Generalizations and related methods [edit]

Muller's method fits a parabola, i.e. a second-order [polynomial](#), to the last three obtained points $f(x_{k-1})$, $f(x_{k-2})$ and $f(x_{k-3})$ in each iteration. One can generalize this and fit a polynomial $p_{k,m}(x)$ of [degree](#) m to the last $m+1$ points in the k^{th} iteration. Our parabola y_k is written as $p_{k,2}$ in this notation. The degree m must be 1 or larger. The next approximation x_k is now one of the roots of the $p_{k,m}$, i.e. one of the solutions of $p_{k,m}(x)=0$. Taking $m=1$ we obtain the secant method whereas $m=2$ gives Muller's method.

Muller calculated that the sequence $\{x_k\}$ generated this way converges to the root ξ with an order μ_m where μ_m is the positive solution of $x^{m+1} - x^m - x^{m-1} - \dots - x - 1 = 0$.

The method is much more difficult though for $m>2$ than it is for $m=1$ or $m=2$ because it is much harder to determine the roots of a polynomial of degree 3 or higher. Another problem is that there seems no prescription of which of the roots of $p_{k,m}$ to pick as the next approximation x_k for $m>2$.

These difficulties are overcome by [Sidi's generalized secant method](#) which also employs the polynomial $p_{k,m}$. Instead of trying to solve $p_{k,m}(x)=0$, the next approximation x_k is calculated with the aid of the derivative of $p_{k,m}$ at x_{k-1} in this method.

References [edit]

- Muller, David E., "A Method for Solving Algebraic Equations Using an Automatic Computer," *Mathematical Tables and Other Aids to Computation*, 10 (1956), 208-215. [JSTOR 2001916](#) [↗]
- Atkinson, Kendall E. (1989). *An Introduction to Numerical Analysis*, 2nd edition, Section 2.4. John Wiley & Sons, New York. [ISBN 0-471-50023-2](#).
- Burden, R. L. and Faires, J. D. *Numerical Analysis*, 4th edition, pages 77ff.
- Press, WH; Teukolsky, SA; Vetterling, WT; Flannery, BP (2007). "[Section 9.5.2. Muller's Method](#)" [↗]. *Numerical Recipes: The Art of Scientific Computing* (3rd ed.). New York: Cambridge University Press. [ISBN 978-0-521-88068-8](#).

External links [edit]

- [Module for Muller's Method by John H. Mathews](#) [↗]

Categories: [Root-finding algorithms](#)

This page was last modified on 15 February 2015, at 11:19.

Text is available under the [Creative Commons Attribution-ShareAlike License](#); additional terms may apply. By using this site, you agree to the [Terms of Use](#) and [Privacy Policy](#). Wikipedia® is a registered trademark of the [Wikimedia Foundation, Inc.](#), a non-profit organization.

[Privacy policy](#) [About Wikipedia](#) [Disclaimers](#) [Contact Wikipedia](#) [Developers](#) [Mobile view](#)

