



WIKIPEDIA  
The Free Encyclopedia

[Main page](#)  
[Contents](#)  
[Featured content](#)  
[Current events](#)  
[Random article](#)  
[Donate to Wikipedia](#)  
[Wikipedia store](#)

Interaction

[Help](#)  
[About Wikipedia](#)  
[Community portal](#)  
[Recent changes](#)  
[Contact page](#)

Tools

[What links here](#)  
[Related changes](#)  
[Upload file](#)  
[Special pages](#)  
[Permanent link](#)  
[Page information](#)  
[Wikidata item](#)  
[Cite this page](#)

Print/export

[Create a book](#)  
[Download as PDF](#)  
[Printable version](#)

Languages

[Português](#)  
[Edit links](#)

[Create account](#) [Log in](#)

[Article](#) [Talk](#)

[Read](#) [Edit](#) [View history](#)

Search

# Lax–Wendroff method

From Wikipedia, the free encyclopedia  
(Redirected from [Lax-Wendroff method](#))

The **Lax–Wendroff method**, named after [Peter Lax](#) and [Burton Wendroff](#), is a [numerical](#) method for the solution of [hyperbolic partial differential equations](#), based on [finite differences](#). It is second-order accurate in both space and time. This method is an example of [explicit time integration](#) where the function that defines governing equation is evaluated at the current time.

Suppose one has an equation of the following form:

$$\frac{\partial u(x,t)}{\partial t} + \frac{\partial f(u(x,t))}{\partial x} = 0$$

where *x* and *t* are independent variables, and the initial state, *u*(*x*, 0) is given.

In the linear case, where *f*(*u*) = *Au* , and *A* is a constant,<sup>[1]</sup>

$$u_i^{n+1} = u_i^n - \frac{\Delta t}{2\Delta x} A \left[ u_{i+1}^n - u_{i-1}^n \right] + \frac{\Delta t^2}{2\Delta x^2} A^2 \left[ u_{i+1}^n - 2u_i^n + u_{i-1}^n \right].$$

This linear scheme can be extended to the general non-linear case in different ways. One of them is letting

$$A(u) = f'(u) = \frac{\partial f}{\partial u}$$

The conservative form of Lax-Wendroff for a general non-linear equation is then

$$u_i^{n+1} = u_i^n - \frac{\Delta t}{2\Delta x} \left[ f(u_{i+1}^n) - f(u_{i-1}^n) \right] + \frac{\Delta t^2}{2\Delta x^2} \left[ A_{i+1/2} \left( f(u_{i+1}^n) - f(u_i^n) \right) - A_{i-1/2} \left( f(u_i^n) - f(u_{i-1}^n) \right) \right].$$

where *A*<sub>*i*±1/2</sub> is the Jacobian matrix evaluated at  $\frac{1}{2}(U_i^n + U_{i\pm1/2}^n)$

To avoid the Jacobian evaluation, use a two-step procedure. What follows is the Richtmyer two-step Lax–Wendroff method. The first step in the Richtmyer two-step Lax–Wendroff method calculates values for *f*(*u*(*x*, *t*)) at half time steps, *t*<sub>*n* + 1/2</sub> and half grid points, *x*<sub>*i* + 1/2</sub>. In the second step values at *t*<sub>*n* + 1</sub> are calculated using the data for *t*<sub>*n*</sub> and *t*<sub>*n* + 1/2</sub>.

First (Lax) steps:

$$\begin{aligned} u_{i+1/2}^{n+1/2} &= \frac{1}{2}(u_{i+1}^n + u_i^n) - \frac{\Delta t}{2\Delta x} (f(u_{i+1}^n) - f(u_i^n)), \\ u_{i-1/2}^{n+1/2} &= \frac{1}{2}(u_i^n + u_{i-1}^n) - \frac{\Delta t}{2\Delta x} (f(u_i^n) - f(u_{i-1}^n)). \end{aligned}$$

Second step:

$$u_i^{n+1} = u_i^n - \frac{\Delta t}{\Delta x} \left[ f(u_{i+1/2}^{n+1/2}) - f(u_{i-1/2}^{n+1/2}) \right].$$

Another method of this same type was proposed by MacCormack. MacCormack's method uses first forward differencing and then backward differencing:

First step:

$$u_i^* = u_i^n - \frac{\Delta t}{\Delta x} (f(u_{i+1}^n) - f(u_i^n)).$$

Second step:

$$u_i^{n+1} = \frac{1}{2}(u_i^n + u_i^*) - \frac{\Delta t}{2\Delta x} \left[ f(u_i^*) - f(u_{i-1}^*) \right].$$

Alternatively, First step:

$$u_i^* = u_i^n - \frac{\Delta t}{\Delta x} (f(u_i^n) - f(u_{i-1}^n)).$$

Second step:

$$u_i^{n+1} = \frac{1}{2}(u_i^n + u_i^*) - \frac{\Delta t}{2\Delta x} \left[ f(u_{i+1}^*) - f(u_i^*) \right].$$

## References <sup>[edit]</sup>

- ↑ LeVeque, Randy J. *Numerical Methods for Conservation Laws*, Birkhauser Verlag, 1992, p. 125.
- ↑ P.D Lax; B. Wendroff (1960). "Systems of conservation laws". *Commun. Pure Appl Math.* **13** (2): 217–237. doi:10.1002/cpa.3160130205 .
- ↑ Michael J. Thompson, *An Introduction to Astrophysical Fluid Dynamics*, Imperial College Press, London, 2006.
- ↑ Press, WH; Teukolsky, SA; Vetterling, WT; Flannery, BP (2007). "Section 20.1. Flux Conservative Initial Value Problems" . *Numerical Recipes: The Art of Scientific Computing* (3rd ed.). New York: Cambridge University Press. p. 1040. ISBN 978-0-521-88068-8.

<span>v</span> · <span>t</span> · <span>e</span>	<b>Numerical partial differential equations by method</b> <span>[hide]</span>	
<b>Finite difference</b>	Parabolic	Forward-time central-space (FTCS) · Crank–Nicolson
	Hyperbolic	Lax–Friedrichs · <b>Lax–Wendroff</b> · MacCormack · Upwind · Method of characteristics
	Others	Alternating direction-implicit (ADI) · Finite-difference time-domain (FDTD)
<b>Finite volume</b>	Godunov · High-resolution · Monotonic upstream-centered (MUSCL) · Advection upstream-splitting (AUSM) · Riemann solver	
<b>Finite element</b>	hp-FEM · Extended (XFEM) · Discontinuous Galerkin (DG) · Spectral element (SEM) · Mortar	
<b>Meshless/Meshfree</b>	Smoothed-particle hydrodynamics (SPH) · Material point method (MPM)	
<b>Domain decomposition</b>	Schur complement · Fictitious domain · Schwarz alternating (additive · abstract additive) · Neumann–Dirichlet · Neumann–Neumann · Poincaré–Steklov operator · Balancing (BDD) · Balancing by constraints (BDDC) · Tearing and interconnect (FETI) · FETI-DP	
<b>Others</b>	Spectral · Pseudospectral (DVR) · Method of lines · Multigrid · Collocation · Level set · Boundary element · Immersed boundary · Analytic element · Particle-in-cell · Isogeometric analysis	

✚✚ *This **applied mathematics**-related article is a **stub**. You can help Wikipedia by **expanding it**.*

Categories: Numerical differential equations | Computational fluid dynamics | Applied mathematics stubs