



 Edit links

$$\begin{aligned}\sin \sigma &= \sqrt{(\cos U_2 \sin \lambda)^2 + (\cos U_1 \sin U_2 - \sin U_1 \cos U_2 \cos \lambda)^2} \\ \cos \sigma &= \sin U_1 \sin U_2 + \cos U_1 \cos U_2 \cos \lambda\end{aligned}$$

$$\sigma = \arctan \frac{\sin \sigma}{\cos \sigma}^{[1][2]}$$

$$\sin \alpha = \frac{\cos U_1 \cos U_2 \sin \lambda}{\sin \sigma}^{[3]}$$

$$\cos^2 \alpha = 1 - \sin^2 \alpha$$

$$\cos(2\sigma_m) = \cos \sigma - \frac{2 \sin U_1 \sin U_2}{\cos^2 \alpha}^{[4]}$$

$$C = \frac{f}{16} \cos^2 \alpha [4 + f(4 - 3 \cos^2 \alpha)]$$

$$\lambda = L + (1 - C)f \sin \alpha \left\{ \sigma + C \sin \sigma \left[ \cos(2\sigma_m) + C \cos \sigma (-1 + 2 \cos^2(2\sigma_m)) \right] \right\}$$

When  $\lambda$  has converged to the desired degree of accuracy ( $10^{-12}$  corresponds to approximately 0.06mm), evaluate the following:

$$u^2 = \cos^2 \alpha \frac{a^2 - b^2}{b^2}$$

$$A = 1 + \frac{u^2}{16384} \left\{ 4096 + u^2 \left[ -768 + u^2(320 - 175u^2) \right] \right\}$$

$$B = \frac{u^2}{1024} \left\{ 256 + u^2 \left[ -128 + u^2(74 - 47u^2) \right] \right\}$$

$$\Delta\sigma = B \sin \sigma \left\{ \cos(2\sigma_m) + \frac{1}{4}B \left[ \cos \sigma (-1 + 2 \cos^2(2\sigma_m)) - \frac{1}{6}B \cos(2\sigma_m) (-3 + 4 \sin^2 \sigma) (-3 + 4 \cos^2(2\sigma_m)) \right] \right\}$$

$$s = bA(\sigma - \Delta\sigma)$$

$$\alpha_1 = \arctan \left( \frac{\cos U_2 \sin \lambda}{\cos U_1 \sin U_2 - \sin U_1 \cos U_2 \cos \lambda} \right)^{[2]}$$

$$\alpha_2 = \arctan \left( \frac{\cos U_1 \sin \lambda}{-\sin U_1 \cos U_2 + \cos U_1 \sin U_2 \cos \lambda} \right)^{[2]}$$

Between two nearly antipodal points, the iterative formula may fail to converge; this will occur when the first guess at  $\lambda$  as computed by the equation above is greater than  $\pi$  in absolute value.

## Direct Problem [\[edit\]](#)

Given an initial point  $(\Phi_1, L_1)$  and initial azimuth,  $\alpha_1$ , and a distance,  $s$ , along the geodesic the problem is to find the end point  $(\Phi_2, L_2)$  and azimuth,  $\alpha_2$ .

Start by calculating the following:

$$U_1 = \arctan((1 - f) \tan \phi_1)$$

$$\sigma_1 = \arctan \left( \frac{\tan U_1}{\cos \alpha_1} \right)^{[2]}$$

$$\sin \alpha = \cos U_1 \sin \alpha_1$$

$$\cos^2 \alpha = 1 - \sin^2 \alpha$$

$$u^2 = \cos^2 \alpha \frac{a^2 - b^2}{b^2}$$

$$A = 1 + \frac{u^2}{16384} \left\{ 4096 + u^2 \left[ -768 + u^2(320 - 175u^2) \right] \right\}$$

$$B = \frac{u^2}{1024} \left\{ 256 + u^2 \left[ -128 + u^2(74 - 47u^2) \right] \right\}$$

Then, using an initial value  $\sigma = \frac{s}{bA}$ , iterate the following equations until there is no significant change in  $\sigma$ :

$$2\sigma_m = 2\sigma_1 + \sigma$$

$$\Delta\sigma = B \sin \sigma \left\{ \cos(2\sigma_m) + \frac{1}{4}B \left[ \cos \sigma (-1 + 2 \cos^2(2\sigma_m)) - \frac{1}{6}B \cos(2\sigma_m) (-3 + 4 \sin^2 \sigma) (-3 + 4 \cos^2(2\sigma_m)) \right] \right\}$$

$$\sigma = \frac{s}{bA} + \Delta\sigma$$

Once  $\sigma$  is obtained to sufficient accuracy evaluate:

$$\phi_2 = \arctan \left( \frac{\sin U_1 \cos \sigma + \cos U_1 \sin \sigma \cos \alpha_1}{(1 - f) \sqrt{\sin^2 \alpha + (\sin U_1 \sin \sigma - \cos U_1 \cos \sigma \cos \alpha_1)^2}} \right)^{[2]}$$

$$\lambda = \arctan \left( \frac{\sin \sigma \sin \alpha_1}{\cos U_1 \cos \sigma - \sin U_1 \sin \sigma \cos \alpha_1} \right)^{[2]}$$

$$C = \frac{f}{16} \cos^2 \alpha [4 + f(4 - 3 \cos^2 \alpha)]$$

$$L = \lambda - (1 - C)f \sin \alpha \left\{ \sigma + C \sin \sigma \left[ \cos(2\sigma_m) + C \cos \sigma (-1 + 2 \cos^2(2\sigma_m)) \right] \right\}$$

$$L_2 = L + L_1$$

$$\alpha_2 = \arctan \left( \frac{\sin \alpha}{-\sin U_1 \sin \sigma + \cos U_1 \cos \sigma \cos \alpha_1} \right)^{[2]}$$

If the initial point is at the North or South pole then the first equation is indeterminate. If the initial azimuth is due East or West then the second equation is indeterminate. If a double valued *atan2* type function is used then these values are usually handled correctly.

## Vincenty's modification [edit]

In his letter to Survey Review in 1976, Vincenty suggested replacing his series expressions for *A* and *B* with simpler formulas using Helmert's expansion parameter *k*<sub>1</sub>:

$$A = \frac{1 + \frac{1}{4}(k_1)^2}{1 - k_1}$$
$$B = k_1(1 - \frac{3}{8}(k_1)^2)$$

where  $k_1 = \frac{\sqrt{(1+u^2)} - 1}{\sqrt{(1+u^2)} + 1}$

## Nearly antipodal points [edit]

As noted above, the iterative solution to the inverse problem fails to converge or converges slowly for nearly antipodal points. An example of slow convergence is (*Φ*<sub>1</sub>, *L*<sub>1</sub>) = (0°, 0°) and (*Φ*<sub>2</sub>, *L*<sub>2</sub>) = (0.5°, 179.5°) for the WGS84 ellipsoid. This requires about 130 iterations to give a result accurate to 1 mm. Depending on how the inverse method is implemented, the algorithm might return the correct result (19936288.579 m), an incorrect result, or an error indicator. An example of an incorrect result is provided by the [NGS online utility](#) [↗] which returns a distance which is about 5 km too long. Vincenty suggested a method of accelerating the convergence in such cases (Rapp, 1973).

An example of a failure of the inverse method to converge is (*Φ*<sub>1</sub>, *L*<sub>1</sub>) = (0°, 0°) and (*Φ*<sub>2</sub>, *L*<sub>2</sub>) = (0.5°, 179.7°) for the WGS84 ellipsoid. In an unpublished report, Vincenty (1975b) gave an alternative iterative scheme to handle such cases. This converges to the correct result 19944127.421 m after about 60 iterations; however, in other cases many thousands of iterations are required.

Newton's method has been successfully used to give rapid convergence for all pairs of input points (Karney, 2013).

## See also [edit]

- Geographical distance
- Great-circle distance
- Meridian arc
- Geodesics on an ellipsoid
- Thaddeus Vincenty
- Geodesy

## Notes [edit]

- ↑ *σ* isn't evaluated directly from sin *σ* or cos *σ* to preserve numerical accuracy near the poles and equator
- ↑ ***a b c d e f g*** The arctan quantity should be evaluated using a two argument *atan2* type function.
- ↑ If sin *σ* = 0 the value of sin *α* is indeterminate. It represents an end point equal to, or diametrically opposite the start point.
- ↑ Start and end point are on the equator. In this case *C* = 0 so the value of cos(*2σ<sub>m</sub>*) is not used. The limiting value is cos(*2σ<sub>m</sub>*) = −1.

## References [edit]

- Bessel, F. W. (2010). "The calculation of longitude and latitude from geodesic measurements (1825)". *Astron. Nachr.* **331** (8): 852–861. arXiv:0908.1824 [↗]. doi:10.1002/asna.201011352 [↗]. English translation of *Astron. Nachr.* **4**, 241–254 (1825).
- Helmert, F. R. (1964). *Mathematical and Physical Theories of Higher Geodesy, Part 1 (1880)* [↗]. St. Louis: Aeronautical Chart and Information Center. Retrieved 2011-07-30. English translation of *Die Mathematischen und Physikalischen Theorieen der Höheren Geodäsie*, Vol. 1 (Teubner, Leipzig, 1880).
- Karney, C. F. F. (2013). "Algorithms for geodesics". *Journal of Geodesy* **87** (1): 43–42. arXiv:1109.4448 [↗]. Bibcode:2013JGeod..87...43K [↗]. doi:10.1007/s00190-012-0578-z [↗] (open access). Addenda [↗].
- Legendre, A. M. (1806). "Analyse des triangles tracés sur la surface d'un sphéroïde" [↗]. *Mém. de l'Inst. Nat. de France* (1st sem.): 130–161. Retrieved 2011-07-30.
- Rainsford, H. F. (1955). "Long geodesics on the ellipsoid". *Bulletin géodésique* **37**: 12–22. Bibcode:1955BGeod..29...12R [↗]. doi:10.1007/BF02527187 [↗].
- Rapp, R. H. (March 1993). *Geometric Geodesy, Part II* [↗] (Technical report). Ohio State University. Retrieved 2011-08-01.
- Vincenty, T. (April 1975a). "Direct and Inverse Solutions of Geodesics on the Ellipsoid with application of nested equations" [↗] (PDF). *Survey Review*. XXIII (misprinted as XXII) (176): 88–93. Retrieved 2009-07-11.
- Vincenty, T. (April 1976). "Correspondence". *Survey Review*. XXIII (180): 294.
- Vincenty, T. (August 1975b). *Geodetic inverse solution between antipodal points* [↗] (PDF) (Technical report). DMAAC Geodetic Survey Squadron. Retrieved 2011-07-28.
- Geocentric Datum of Australia (GDA) Reference Manual* [↗] (PDF). Intergovernmental committee on survey and mapping (ICSM). February 2006. ISBN 0-9579951-0-5. Retrieved 2009-07-11.

## External links [edit]

- Online calculators from [Geoscience Australia](#):
  - [Vincenty Direct](#) [↗] (destination point)
  - [Vincenty Inverse](#) [↗] (distance between points)
- Calculators from the [U.S. National Geodetic Survey](#):
  - [Online and downloadable PC-executable calculation utilities](#) [↗], including forward (direct) and inverse problems, in both two and three dimensions (accessed 2011-08-01).

- Online calculators with JavaScript source code by Chris Veness (Creative Commons Attribution license):
  - [Vincenty Direct](#) (destination point)
  - [Vincenty Inverse](#) (distance between points)
- [GeographicLib](#) provides a utility GeodSolve (with MIT/X11 licensed source code) for solving direct and inverse geodesic problems. Compared to Vincenty, this is about 1000 times more accurate (error = 15 nm) and the inverse solution is complete. Here is an [online version of GeodSolve](#).

Categories: [Geodesy](#)

This page was last modified on 2 September 2015, at 11:42.

Text is available under the [Creative Commons Attribution-ShareAlike License](#); additional terms may apply. By using this site, you agree to the [Terms of Use](#) and [Privacy Policy](#). Wikipedia® is a registered trademark of the [Wikimedia Foundation, Inc.](#), a non-profit organization.

[Privacy policy](#) [About Wikipedia](#) [Disclaimers](#) [Contact Wikipedia](#) [Developers](#) [Mobile view](#)

