




WIKIPEDIA  
The Free Encyclopedia

Main page  
Contents  
Featured content  
Current events  
Random article  
Donate to Wikipedia  
Wikipedia store

Interaction  
Help  
About Wikipedia  
Community portal  
Recent changes  
Contact page

Tools  
What links here  
Related changes  
Upload file  
Special pages  
Permanent link  
Page information  
Wikidata item  
Cite this page

Print/export  
Create a book  
Download as PDF  
Printable version


Languages   
العربية  
Català  
Deutsch  
Español  
Français  
עברית  
Nederlands  
Русский  
Suomi  
Svenska  
Українська  
Tiếng Việt

 Edit links

Create account Log in

Article [Talk](#)

[Read](#) [Edit](#) [More](#) ▾



# Lucas primality test

From Wikipedia, the free encyclopedia

*"Lucas–Lehmer test" redirects here. For the test for Mersenne numbers, see [Lucas–Lehmer primality test](#). For the Lucas–Lehmer–Riesel test, see [Lucas–Lehmer–Riesel test](#). For the Lucas probable prime test, see [Lucas pseudoprime](#).*

In [computational number theory](#), the **Lucas test** is a [primality test](#) for a natural number *n*; it requires that the [prime factors](#) of *n* − 1 be already known.<sup>[1][2]</sup> It is the basis of the [Pratt certificate](#) that gives a concise verification that *n* is prime.

## Contents

[\[hide\]](#)

- 1 Concepts
- 2 Example
- 3 Algorithm
- 4 See also
- 5 Notes

## Concepts

[\[edit\]](#)

Let *n* be a positive integer. If there exists an integer 1 < *a* < *n* such that

$$a^{n-1} \equiv 1 \pmod{n}$$

and for every prime factor *q* of *n* − 1

$$a^{(n-1)/q} \not\equiv 1 \pmod{n}$$

then *n* is prime. If no such number *a* exists, then *n* is either 1 or [composite](#).

The reason for the correctness of this claim is as follows: if the first equivalence holds for *a*, we can deduce that *a* and *n* are [coprime](#). If *a* also survives the second step, then the [order](#) of *a* in the [group](#) (**Z**/*n***Z**)<sup>\*</sup> is equal to *n*−1, which means that the order of that group is *n*−1 (because the order of every element of a group divides the order of the group), implying that *n* is [prime](#). Conversely, if *n* is prime, then there exists a [primitive root modulo](#) *n*, or [generator](#) of the group (**Z**/*n***Z**)<sup>\*</sup>. Such a generator has order |(bZ/nZ)<sup>\*</sup>| = *n*−1 and both equivalences will hold for any such primitive root.

Note that if there exists an *a* < *n* such that the first equivalence fails, *a* is called a [Fermat witness](#) for the compositeness of *n*.

## Example

[\[edit\]](#)

For example, take *n* = 71. Then *n* − 1 = 70 and the prime factors of 70 are 2, 5 and 7. We randomly select an *a*=17 < *n*. Now we compute:

$$17^{70} \equiv 1 \pmod{71}.$$

For all integers *a* it is known that

$$a^{n-1} \equiv 1 \pmod{n} \text{ if and only if } \text{ord}(a)|(n-1).$$

Therefore, the multiplicative order of 17 (mod 71) is not necessarily 70 because some factor of 70 may also work above. So check 70 divided by its prime factors:

$$17^{35} \equiv 70 \not\equiv 1 \pmod{71}$$

$$17^{14} \equiv 25 \not\equiv 1 \pmod{71}$$

$$17^{10} \equiv 1 \equiv 1 \pmod{71}.$$

Unfortunately, we get that 17<sup>10</sup>≡1 (mod 71). So we still don't know if 71 is prime or not.

We try another random *a*, this time choosing *a* = 11. Now we compute:

$$11^{70} \equiv 1 \pmod{71}.$$

Again, this does not show that the multiplicative order of 11 (mod 71) is 70 because some factor of 70 may also work. So check 70 divided by its prime factors:

$$11^{35} \equiv 70 \not\equiv 1 \pmod{71}$$
$$11^{14} \equiv 54 \not\equiv 1 \pmod{71}$$
$$11^{10} \equiv 32 \not\equiv 1 \pmod{71}.$$

So the multiplicative order of 11 (mod 71) is 70, and thus 71 is prime.

(To carry out these [modular exponentiations](#), one could use a fast exponentiation algorithm like [binary](#) or [addition-chain exponentiation](#)).

## Algorithm [edit]

The algorithm can be written in [pseudocode](#) as follows:

```
Input:  $n > 2$ , an odd integer to be tested for primality;  $k$ , a parameter that determines the accuracy of the test
Output: prime if  $n$  is prime, otherwise composite or possibly composite;
determine the prime factors of  $n-1$ .
LOOP1: repeat  $k$  times:
    pick  $a$  randomly in the range  $[2, n - 1]$ 
    if  $a^{n-1} \not\equiv 1 \pmod{n}$  then return composite
    otherwise
        LOOP2: for all prime factors  $q$  of  $n-1$ :
            if  $a^{(n-1)/q} \not\equiv 1 \pmod{n}$ 
                if we did not check this equality for all prime factors of  $n-1$ 
                    then do next LOOP2
                otherwise return prime
            otherwise do next LOOP1
return possibly composite.
```

## See also [edit]

- Édouard Lucas
- Fermat's little theorem

## Notes [edit]

1.

^

Crandall, Richard; Pomerance, Carl (2005). *Prime Numbers: a Computational Perspective (2nd edition)*. Springer. p. 173. ISBN 0-387-25282-7.

2.

^

Křížek, Michal; Luca, Florian; Somer, Lawrence (2001). *17 Lectures on Fermat Numbers: From Number Theory to Geometry*. CMS Books in Mathematics 9. Canadian Mathematical Society/Springer. p. 41. ISBN 0-387-95332-9.

<div>V · T · E</div>	Number-theoretic algorithms	<span>[hide]</span>
<b>Primality tests</b>	AKS test · APR test · Baillie–PSW · ECPP test · Elliptic curve · Pocklington · Fermat · Lucas · Lucas–Lehmer · Lucas–Lehmer–Riesel · Proth's theorem · Pépin's · Quadratic Frobenius test · Solovay–Strassen · Miller–Rabin	
<b>Prime-generating</b>	Sieve of Atkin · Sieve of Eratosthenes · Sieve of Sundaram · Wheel factorization	
<b>Integer factorization</b>	Continued fraction (CFRAC) · Dixon's · Lenstra elliptic curve (ECM) · Euler's · Pollard's rho · $p - 1$ · $p + 1$ · Quadratic sieve (QS) · General number field sieve (GNFS) · Special number field sieve (SNFS) · Rational sieve · Fermat's · Shanks' square forms · Trial division · Shor's	
<b>Multiplication</b>	Ancient Egyptian · Long · Karatsuba · Toom–Cook · Schönhage–Strassen · Fürer's	
<b>Discrete logarithm</b>	Baby-step giant-step · Pollard rho · Pollard kangaroo · Pohlig–Hellman · Index calculus · Function field sieve	
<b>Greatest common divisor</b>	Binary · Euclidean · Extended Euclidean · Lehmer's	
<b>Modular square root</b>	Cipolla · Pocklington's · Tonelli–Shanks	
<b>Other algorithms</b>	Chakravala · Cornacchia · Integer relation · Integer square root · Modular exponentiation · Schoof's	
<i>Italics</i> indicate that algorithm is for numbers of special forms · <small>SMALLCAPS</small> indicate a deterministic algorithm		

This page was last modified on 24 May 2015, at 21:26.

Text is available under the [Creative Commons Attribution-ShareAlike License](#); additional terms may apply. By using this site, you agree to the [Terms of Use](#) and [Privacy Policy](#). Wikipedia® is a registered trademark of the [Wikimedia Foundation, Inc.](#), a non-profit organization.

[Privacy policy](#) [About Wikipedia](#) [Disclaimers](#) [Contact Wikipedia](#) [Developers](#) [Mobile view](#)

