

Main page
Contents
Featured content
Current events
Random article
Donate to Wkipedia
Wkipedia store

Interaction

Help About Wikipedia Community portal Recent changes Contact page

Tools

What links here Related changes Upload file Special pages Permanent link Page information Wkidata item Cite this page

Print/export

Create a book Download as PDF Printable version

Languages

Italiano

Українська

0

Article Talk Read Edit View histon

Petrick's method

From Wikipedia, the free encyclopedia

In Boolean algebra, **Petrick's method** (also known as the *branch-and-bound* method) is a technique for determining all minimum sum-of-products solutions from a prime implicant chart. Petrick's method is very tedious for large charts, but it is easy to implement on a computer.

- 1. Reduce the prime implicant chart by eliminating the essential prime implicant rows and the corresponding columns.
- 2. Label the rows of the reduced prime implicant chart P_1 , P_2 , P_3 , P_4 , etc.
- 3. Form a logical function P which is true when all the columns are covered. P consists of a product of sums where each sum term has the form $(P_{i0} + P_{i1} + \cdots + P_{iN})$, where each P_{ij} represents a row covering column i.
- 4. Reduce P to a minimum sum of products by multiplying out and applying X+XY=X
- 5. Each term in the result represents a solution, that is, a set of rows which covers all of the minterms in the table. To determine the minimum solutions, first find those terms which contain a minimum number of prime implicants.
- 6. Next, for each of the terms found in step five, count the number of literals in each prime implicant and find the total number of literals.
- 7. Choose the term or terms composed of the minimum total number of literals, and write out the corresponding sums of prime implicants.

Example of Petrick's method (copied from http://www.mrc.uidaho.edu/mrc/people/jff/349/lect.10 년)

Following is the function we want to reduce:

$$f(A,B,C) = \sum m(0,1,2,5,6,7)$$

The prime implicant chart from the Quine-McCluskey algorithm is as follows:

Based on the X marks in the table above, build a product of sums of the rows where each row is added, and columns are multiplied together:

```
(K+L) (K+M) (L+N) (M+P) (N+Q) (P+Q)
```

Use the distributive law to turn that expression into a sum of products. Also use the following equivalences to simplify the final expression: X + XY = X and XX = X and X+X=X

```
= (K+L) (K+M) (L+N) (M+P) (N+Q) (P+Q)
= (K+LM) (N+LQ) (P+MQ)
= (KN+KLQ+LMN+LMQ) (P+MQ)
= KNP + KLPQ + LMNP + LMPQ + KMNQ + KLMQ + LMNQ + LMQ
```

Now use again the following equivalence to further reduce the equation: X + XY = X

```
= \  \, \mathsf{KNP} \ + \  \, \mathsf{KLPQ} \ + \  \, \mathsf{LMNP} \ + \  \, \mathsf{LMQ} \ + \  \, \mathsf{KMNQ}
```

Choose products with fewest terms, in our example, there are two products with three terms:

KNP LMQ

Choose term or terms with fewest total literals. In our example, the two products both expand to 6 literals total each:

```
KNP expands to a'b'+ bc'+ ac

LMQ expands to a'c'+ b'c + ab
```

So either one can be used. In general, application of Petricks method is tedious for large charts, but it is easy to implement on a computer.

External links [edit]

• [1] & Tutorial on Quine-McCluskey and Petrick's method (pdf).

Categories: Boolean algebra

This page was last modified on 15 March 2013, at 06:41.

Text is available under the Creative Commons Attribution-ShareAlike License; additional terms may apply. By using this site, you agree to the Terms of Use and Privacy Policy. Wikipedia® is a registered trademark of the Wikimedia Foundation, Inc., a non-profit organization.

Privacy policy About Wikipedia Disclaimers Contact Wikipedia Developers Mobile view



