

# **OPEN** Generation of macroscopic Schrödinger cat state in diamond mechanical resonator

Received: 18 August 2016 Accepted: 27 October 2016 Published: 23 November 2016 Qizhe Hou<sup>1</sup>, Wanli Yanq<sup>2</sup>, Changyong Chen<sup>3</sup> & Zhangqi Yin<sup>4</sup>

We propose a scheme to generate macroscopic Schrödinger cat state (SCS) in diamond mechanical resonator (DMR) via the dynamical strain-mediated coupling mechanism. In our model, the direct coupling between the nitrogen-vacancy (NV) center and lattice strain field enables coherent spinphonon interactions in the quantum regime. Based on a cyclic  $\Delta$ -type transition structure of the NV center constructed by combining the quantized mechanical strain field and a pair of external microwave fields, the populations of the different energy levels can be selectively transferred by controlling microwave fields, and the SCS can be created by adjusting the controllable parameters of the system. Furthermore, we demonstrate the nonclassicality of the mechanical SCS both in non-dissipative case and dissipative case. The experimental feasibility and challenge are justified using currently available technology.

Recently, the nitrogen-vacancy (NV) centers formed by a substitutional nitrogen atom and an adjacent lattice vacancy in diamond become the most promising solid-state platform for quantum information processing (QIP)1 and nanoscale sensors<sup>2</sup>, due to its easy controllability and fast manipulation<sup>3</sup>, as well as long coherence properties (electron spin  $\sim 1 \, ms$  and nuclear spin  $> 1 \, s$ ) in a wide temperature range, even at room-temperature  $^{4-8}$ . Additionally, the NV center could couple to both optical and microwave fields simultaneously, which makes it possible be used as an excellent quantum interface between optical and solid-state systems<sup>9,10</sup>.

Meantime, the hybrid system consisting of a high-Q single-crystal diamond mechanical resonator (DMR) and embedded NV centers may provide a promising platform and open up a new perspective towards achieving quantum control and studying significant quantum optics or novel quantum phenomena<sup>11-22</sup>. In these systems, the embedded NV centers are highly susceptible to deformations of the surrounding lattice, where the strain field is robust against the dephasing or heating of the environment<sup>12,13</sup>. Applying this direct strain coupling mechanism, previous investigations have been focused on the mechanical spin driving<sup>12-15</sup>, enhancement of the coherence time of the NV center<sup>16</sup>, spin squeezing<sup>17</sup>, phonon cooling and lasing<sup>18</sup>.

On the other hand, based on the quantum superposition principle, the generation of the macroscopic quantum superposition state such as Schrödinger cat state (SCS) has attracted abundant attention because SCS plays an important role in the study of the foundations of quantum theory<sup>23–31</sup>, and brings many potentially applications, such as testing the wave function collapse<sup>32</sup> and uncertainty relation<sup>33</sup>. Experimentally, the macroscopic SCS has been observed in various physical systems including superconducting qubit system<sup>34</sup>, ion trap<sup>35</sup>, cavity QED36, and linear optical system37. However, generating coherent states with macroscopically distinct amplitudes in the phase space is a difficult task in nanomechanical systems<sup>38–40</sup>, it remains a challenge to realize SCS in such

In the present work, we introduce a scheme to generate macroscopic SCS in DMR via the dynamical strain-mediated coupling mechanism. In our model, the direct coupling between the lattice strain field and the NV center enables coherent spin-phonon interactions in the quantum regime, and the cyclic  $\Delta$ -type transition structure of the NV center system could be constructed by combining the quantized mechanical strain field and a pair of external microwave fields, where one-photon and two-photon processes coexist. Using this cyclic

<sup>1</sup>National Laboratory of Solid State Microstructures and School of Physics, Nanjing University, Nanjing 210093, China. 2State Key Laboratory of Magnetic Resonance and Atomic and Molecular Physics, Wuhan Institute of Physics and Mathematics, Chinese Academy of Sciences, Wuhan 430071, China. <sup>3</sup>Department of Physics, Shaoguan University, Shaoquan, Guangdong 512005, China. <sup>4</sup>Center for Quantum Information, Institute for Interdisciplinary Information Sciences, Tsinghua University, Beijing 100084, China. Correspondence and requests for materials should be addressed to W.Y. (email: ywl@wipm.ac.cn) or Z.Y. (email: yinzhanqqi@mail.tsinqhua.edu.cn)




Figure 1. (a) Schematic of the hybrid system, where the NV electron spin (red row) is embedded in the DMR. L, w, and  $h(L \gg w, h)$  represent the length, width, and height of the DMR, respectively. (b) Levels of  $\Delta$ -type transition ground-state electronic spin NV center, consisting of the levels of  $|0\rangle$  and  $|\pm 1\rangle$ , where, two classical microwave fields with frequencies  $\omega_1$  and  $\omega_2$  induce the transitions  $|0\rangle \leftrightarrow |\pm 1\rangle$  with Rabi frequencies  $G_1$  and  $G_2$ , respectively. In addition, the transition  $|+1\rangle \leftrightarrow |-1\rangle$  is coupled to the strain field (the frequency  $\omega_m$ ) with the coupling strength  $\lambda$ .  $D_{gs}$  is the zero-field splitting between the state  $|0\rangle$  and the nearly degenerated states  $|\pm 1\rangle$  which can be split by  $\Delta_p$  in a magnetic field.  $\Delta_{1(2)}$  are the detunings between the frequencies of the transition  $|0\rangle \leftrightarrow |\pm 1\rangle$  and the frequencies of the microwave fields. (c) Level structure of the effective  $\Lambda$ -type transition structure. The transitions  $|\pm\rangle \leftrightarrow |+1\rangle$  are coupled to the displaced quantized fields with the coupling strengths  $\lambda_1$  and  $\lambda_2$ , respectively. The energy spacing between energy levels  $|+\rangle$  and  $|-\rangle$  is  $2\omega$ .

population transfer, the population of the different energy levels of the NV center can be selectively transferred by controlling classical fields, and the SCS can be created by adjusting the controllable parameters of the system, such as the detuning between the NV center and the driving fields, along with their Rabi frequency. Here the mechanical SCS is described by the phase-space quasiprobability distribution, such as Husimi Q function and Winger function. Additionally, comparing with the non-dissipative case with the dissipative case through the superoperator method<sup>41,42</sup>, we find that the dissipation effect decreases the generation efficiency of SCS. Furthermore, we demonstrate the nonclassicality of the SCS in both non-dissipative case and dissipative case.

### Results

**System and Model.** As shown in Fig. 1(a), the system under consideration is a monolithic hybrid quantum device DMR including a diamond cantilever with an embedded NV center. The NV center consists of a substitutional nitrogen atom and a vacancy in an adjacent lattice site with  $C_{3\nu}$  symmetry<sup>43</sup>. It is negatively charged with two unpaired electrons located at the vacancy, and is treated as electronic spin S=1. The ground state is spin triplet and labeled as  $|m_s\rangle$  with  $m_s=0$ ,  $\pm 1$ . The zero-field splitting between the spin sublevels  $|0\rangle$  and  $|\pm 1\rangle$  is  $D_{gs}/2\pi \simeq 2.87\,\mathrm{GHz}$ , originating from the nonaveraged electronic spin-spin interactions<sup>44</sup>. In general, the quantum spin control of the NV center can be achieved with magnetic field<sup>45</sup> or optical field<sup>46</sup>.

In our work, we mainly study individual NV center embedded in DMR, here the coherent strain driving of the NV center is based on the sensitive response of the spin to strain in the diamond host lattice. More specifically, when the DMR vibrates, it changes the local strain and induces a strain field at the position of the NV center. The strain field (behaved as an effective local electric field<sup>44</sup>) breaks the symmetry of the NV center and causes the energy shifts as well as a mixing of the states  $|+1\rangle$  and  $|-1\rangle$ , which results in the direct coupling of the DMR and the transition  $|+1\rangle \leftrightarrow |-1\rangle$  of the NV center<sup>43,44</sup>. It offers a new approach to mechanical driving the NV spin. The Hamiltonian of the NV center coupling to the magnetic field and the strain field can be written as 17,44,47 (in units of the Planck constant, i.e.,  $\hbar = 1$ , hereafter)

$$H_{NV} = (D_{gs} + \epsilon_{\parallel} E_z) S_z^2 + g_s \mu_B \vec{B} \cdot \vec{S} - \epsilon_{\perp} [E_x (S_x S_y + S_y S_x) + E_y (S_x^2 - S_y^2)], \tag{1}$$

where  $\vec{B} = B_x \vec{e}_x + B_y \vec{e}_y + B_z \vec{e}_z$  and  $\vec{S} = S_x \vec{e}_x + S_y \vec{e}_y + S_z \vec{e}_z$  are the external magnetic field and spin-1 operator with  $B_i$  and  $S_i$  (i=x,y,z) the corresponding components, respectively.  $\vec{e}_x$ ,  $\vec{e}_y$ , and  $\vec{e}_z$  are unit vectors along the x,y,z direction.  $e_{\parallel(\perp)}$  are the stress coupling constants along the direction parallel (perpendicular) to the NV symmetry axis (here we assume that the z-axis is aligned with the NV symmetry axis).  $g_z \simeq 2$ ,  $\mu_B$  is the Bohr magneton.  $E_i$  (i=x,y,z) are the i-axis components of the strain field. In the  $S_z$  basis ( $S_z|m_s\rangle = m_s|m_s\rangle$ ), the spin operator can be expressed as  $S_z = \sigma_{+1,+1} - \sigma_{-1,-1}$ ,  $S_z = (\sigma_{+1,0} + \sigma_{0,+1} + \sigma_{-1,0} + \sigma_{0,-1})/\sqrt{2}$ , and  $S_y = (\sigma_{+1,0} - \sigma_{0,+1} - \sigma_{-1,0} + \sigma_{0,-1})/\sqrt{2}$  is with  $\sigma_{j,k} = |j\rangle \langle k|$  ( $j,k=0,\pm 1$ ) the raising or lowering operators (for  $j\neq k$ ) and energy level populations (for j=k) of the NV center. Then, the Hamiltonian  $H_{NV}$  becomes

$$H_{1} = (D_{gs} + \epsilon_{\parallel}E_{z} + g_{s}\mu_{B}B_{z})\sigma_{+1,+1} + (D_{gs} + \epsilon_{\parallel}E_{z} - g_{s}\mu_{B}B_{z})\sigma_{-1,-1} + g_{s}\mu_{B}(B_{m}\sigma_{0,+1} + B_{n}\sigma_{0,-1} + H.c.) - \epsilon_{\perp}(E_{+}S_{+}^{2} + E_{-}S_{-}^{2})/2,$$
(2)

where  $E_{\pm} = E_x \pm i E_y$ ,  $S_{\pm} = S_x \pm i S_y$ ,  $B_m = (B_x + i B_y)/\sqrt{2}$ ,  $B_n = (B_x - i B_y)/\sqrt{2}$ , and the state  $|0\rangle$  is set as the energy zero point.

The DMR can be treated as a harmonic oscillator with the Hamiltonian  $H_{DMR} = \omega_m a^\dagger a$ , where a ( $a^\dagger$ ) the annihilation (creation) operators of the quantized mode, and  $\omega_m$  is the frequency of DMR. In addition, for small beam displacements, the perpendicular strain field can be quantized as  $E_+ = E_0 a$  and  $E_- = E_0 a^\dagger$  with  $E_0$  the strain on account of the zero point motion of the resonant mode<sup>17</sup>. The NV center in diamond is highly susceptible to deformations of the surrounding lattice, therefore, the vibration due to the ground mechanical mode of the nanoresonator changes the local strain where the NV center is located, and gives rising to an effective, strain-induced electric field  $E_0$ , which results into the magnetically-forbidden transition  $|+1\rangle \leftrightarrow |-1\rangle$  with the related spin-phonon coupling strength  $\lambda = -\epsilon_\perp E_0^{43,44}$ . As a result, the cyclic  $\Delta$ -type transition among all three spin levels of the NV center has been formed, as illustrated in Fig. 1b. The whole system can be written as  $H_2 = \omega_+ \sigma_{+1,+1} + \omega_- \sigma_{-1,-1} + \omega_m a^\dagger a + [\lambda \sigma_{+1,-1} a + G_1 \sigma_{0,+1} \exp(i\omega_1 t) + G_2 \sigma_{0,-1} \exp(i\omega_2 t) + \text{H.c.}]$ , where H.c. represents the conjugate Hermitian, and  $\omega_\pm = D_{gs} + \epsilon_\parallel E_z \pm g_s \mu_B B_z$ . In the interacting picture, the Hamiltonian  $H_2$  changes into

$$H_3 = \Delta_1 \sigma_{+1,+1} + \Delta_2 \sigma_{-1,-1} + (\lambda \sigma_{+1,-1} a + G_1 \sigma_{0,+1} + G_2 \sigma_{0,-1} + H.c.), \tag{3}$$

where  $\Delta_{1(2)} = \omega_{+(-)} - \omega_{1(2)}$  represent the corresponding detunings between the transition frequencies  $\omega_{+(-)}$  of the NV center and the frequencies of the classical microwave fields, respectively. Here the relation  $\omega_1 - \omega_2 = \omega_m$  has been set.

**Generation of the SCS.** In this section, we focus on how to generate the SCS via the dynamical strain-mediated coupling mechanism, based on the cyclic  $\Delta$ -type transition structure.

To intuitively describe the main mechanism of how to create the SCS, we transform the cyclic  $\Delta$ -type transition configuration in Fig. 1b into an effective  $\Lambda$ -type transition structure in Fig. 1c. Here the transitions  $|+1\rangle \leftrightarrow |\pm\rangle$  are induced by two displaced quantized phonon fields in Fig. 1c, which results from the quantized phonon field coupling to the transition  $|+1\rangle \leftrightarrow |-1\rangle$  in Fig. 1b, through an unitary transformation. Concretely speaking, the dressed states  $|\pm\rangle$  have the following relation between the  $|-1\rangle$  and  $|0\rangle$  states as  $|+\rangle = \cos(\theta/2) |-1\rangle + \sin(\theta/2) |0\rangle$  and  $|-\rangle = -\sin(\theta/2) |-1\rangle + \cos(\theta/2) |0\rangle$  with the mixing angle  $\theta = \arctan[2G_2/\Delta_2]$ .

Based on this  $\Lambda$ -type transition structure in Fig. 1c, the effective Hamiltonian  $H_{\it eff}$  has the following form  $H_{\it eff} = H_{11} + H_{+-}$  with

$$H_{11} = (\Delta_1 + \omega_p p p^{\dagger} + \omega_q q q^{\dagger}) \sigma_{+1,+1}, \tag{4}$$

and

$$H_{+-} = (\varepsilon_+ - \omega_p p^{\dagger} p) \sigma_{+,+} + (\varepsilon_- - \omega_q q^{\dagger} q) \sigma_{-,-} + \eta(p^{\dagger} q \sigma_{+,-} + p q^{\dagger} \sigma_{-,+}), \tag{5}$$

where  $p=a+k_1$  and  $q=a-k_2$  with  $k_1=G_1\tan(\theta/2)/\lambda$ ,  $k_2=G_1\cot(\theta/2)/\lambda$ .  $\sigma_{j,k}=|j\rangle\langle k|$   $(j,k=+1,\pm)$ . The other controllable parameters are  $\lambda_1=\lambda\cos(\theta/2)$ ,  $\lambda_2=\lambda\sin(\theta/2)$ ,  $\eta=\frac{2\Delta_1-\Delta_2}{2\Delta_1\Delta_-}\lambda_1\lambda_2$ ,  $\omega_{p(q)}=\lambda_{1(2)}^2/\Delta_{+(-)}$ ,

 $\varepsilon_{\pm} = \Delta_2/2 \pm \omega$ , and  $\Delta_{\pm} = \Delta_1 - \varepsilon_{\pm}$  with the dressed frequency  $\omega = \sqrt{G_2^2 + \Delta_2^2/4}$ . The details of our deductions are presented in Section of Method.

In the large detuning case  $\Delta_{\pm} \gg \lambda_{1(2)}$ , the upper state  $|+1\rangle$  can be adiabatically eliminated by using the rotating wave approximation and the method of Fröhlich-Nakajima transformation (FNT)<sup>48</sup>. As a result, we obtain the Hamiltonian  $H_{+-} = H_{+}\sigma_{+,+} + H_{-}\sigma_{-,-}$  with

$$H_{+} = -\omega_{p} a^{\dagger} a - \omega_{p} k_{1} (a + a^{\dagger}) + \varepsilon_{+} - \omega_{p}^{2} k_{1}^{2}, \tag{6}$$

and

$$H_{-} = -\omega_q a^{\dagger} a + \omega_q k_2 (a + a^{\dagger}) + \varepsilon_{-} - \omega_q^2 k_2^2. \tag{7}$$

Note that the Hamiltonian  $H_{+-}$  can be employed to generate the SCS because both the Hamiltonian  $H_+$  and  $H_-$  also describe the forced harmonic oscillator.

Assuming that the whole system is initially in the state  $|\psi(0)\rangle = \sin(\theta/2)| + \rangle |0\rangle_m + \cos(\theta/2)| - \rangle |0\rangle_m$ , where the DMR is initially prepared in the vacuum state  $|0\rangle_m$ , while the NV center is in the ground state  $|0\rangle = \sin(\theta/2)| + \rangle + \cos(\theta/2)| - \rangle^{14}$ . According to the unitary operator  $U = \exp(-iH_{+-}t)$ , the whole system evolves into the time-dependent state

$$|\psi(t)\rangle = \sin(\theta/2)A|\alpha, +\rangle + \cos(\theta/2)B|\beta, -\rangle,$$
 (8)

where  $|\alpha\rangle = |-k_1 + k_1 \exp(i\omega_p t)\rangle$  and  $|\beta\rangle = |k_2 - k_2 \exp(i\omega_q t)\rangle$  represent the coherent states of DMR.  $A = \exp[ik_1^2 \sin(\omega_p t) - i\varepsilon_+ t]$  and  $B = \exp[ik_2^2 \sin(\omega_q t) - i\varepsilon_- t]$  are the time-dependent global phases, and we

denote the phase difference between  $|\alpha\rangle$  and  $|\beta\rangle$  by  $\delta_{AB}$ . Note that the state  $|\psi(t)\rangle$  is an entangled cat state with different coherent states  $|\alpha\rangle$  and  $|\beta\rangle$  ( $\alpha \neq \beta$ ), and it is an entanglement between the mechanical phonon state and the NV center. If the readout result of the NV center is  $|0\rangle$ , the mechanical phonon state becomes  $|\psi_p(t)\rangle = A \sin^2(\theta/2) |\alpha\rangle + B \cos^2(\theta/2) |\beta\rangle$ , which is a standard mechanical SCS. In experiments, the relation between the population of  $|0\rangle$  and the phase difference  $\delta_{AB}$  has been used to directly detect the SCS  $|\psi_{a}(t)\rangle$ . In our work, we will verify the existence of the mechanical SCS by calculating the overlap between these two coherent states  $|\alpha\rangle$  and  $|\beta\rangle$  as

$$|\langle \beta | \alpha \rangle|^2 = \exp[-R(t)], \tag{9}$$

$$R(t) = 2(k_1 + k_2)^2 - 4k_1k_2\sin^2(\omega_d t/2) - 2(k_1 + k_2)[k_1\cos(\omega_p t) + k_2\cos(\omega_q t)]$$
(10)

with  $\omega_d = \omega_p - \omega_a$ .

In the following, we use the quantum master equation to investigate the decoherence effect as

$$\dot{\rho} = -i[H_{+-}, \rho(t)] + \gamma \mathcal{D}[\sigma_{-+}]\rho + \kappa (\overline{n}_{th} + 1) \mathcal{D}[a]\rho + \kappa \overline{n}_{th} \mathcal{D}[a^{\dagger}]\rho, \tag{11}$$

where  $\mathcal{D}[o]\rho = o\rho o^{\dagger} - (o^{\dagger}o\rho + \rho o^{\dagger}o)/2$  is the standard Lindblad superoperator that describes the dissipation effects.  $\gamma$  and  $\kappa$  represent the decay rate of the NV center and the DMR, respectively.  $\bar{n}_{th} = 1/[\exp(\hbar\omega_m/k_BT) - 1]$ denotes the equilibrium phonon occupation number at temperature T with  $k_B$  the Boltzmann constant. In the basis  $\{|+\rangle, |-\rangle\}$ , using the superoperator method<sup>41</sup>, the analytical results can be obtained as

$$\dot{\rho}_{++}(t) = -i \Big[ H_+, \, \rho_{++}(t) \Big] + \mathcal{L} \rho_{++}(t), \tag{12}$$

$$\dot{\rho}_{-}(t) = -i[H_{-}, \rho_{-}(t)] + \mathcal{L}\rho_{-}(t), \tag{13}$$

$$\dot{\rho}_{+-}(t) = -iH_{+}\rho_{+-}(t) + i\rho_{+-}(t)H_{-} + \mathcal{L}\rho_{+-}(t), \tag{14}$$

$$\dot{\rho}_{-+}(t) = i\rho_{-+}(t)H_{+} - iH_{-}\rho_{-+}(t) + \mathcal{L}\rho_{-+}(t), \tag{15}$$

where  $\dot{p}_{ii}(t) = \langle i|\dot{p}(t)|j\rangle$  (i, j=+,-) represent the matrix elements of the  $\dot{p}(t)$ , and the superoperator is  $\mathcal{L} = \kappa (2\ddot{a} \cdot a^{\dagger} - a^{\dagger}a \cdot - a^{\dagger}a)/2$ . In our calculation, the operator disenable equation  $\exp(N + P)$  $\exp[N(1-e^{-\xi})/\xi] \exp(P) = \exp(P) \exp[N(e^{\xi}-1)/\xi]$  has been used, when  $[N, P] = NP - PN = \xi N$ . If the readout result of the NV center is  $|0\rangle$ , the density matrix with respect to the mechanical phonon state is

$$\rho(t) = \sin^{4}(\theta/2) |\exp(T_{p}^{k_{1}})|^{2} |\alpha_{d}\rangle\langle\alpha_{d}| + \cos^{4}(\theta/2) |\exp(T_{q}^{k_{2}})|^{2} |\beta_{d}\rangle\langle\beta_{d}|$$

$$+ \{\sin^{2}(\theta) \exp(-i\varepsilon_{d}t) \exp(i\omega_{l}^{k}t) \times \exp(T_{p}^{k_{1}}) \exp(T_{q}^{k_{2}})^{*} |\alpha_{d}\rangle\langle\beta_{d}|/4$$

$$+ H.c.\},$$
(16)

where,  $|\alpha_d\rangle$  and  $|\beta_d\rangle$  represent the coherent states with  $\alpha_d=\omega_p k_1 [\exp(i\omega_p t-\kappa t/2)-1]/(\omega_p+i\kappa/2)$  and  $\beta_d=-\omega_q k_2 [\exp(i\omega_q t-\kappa t/2)-1]/(\omega_q+i\kappa/2)$ . The other parameters are  $\varepsilon_d=\varepsilon_+-\varepsilon_-$ ,  $\omega_l^{\ k}=\omega_p k_1^2-\omega_q k_2^2$ , and


$$T_{p(q)}^{k_{1(2)}} = \frac{\omega_{p(q)}^{2} k_{1(2)}^{2}}{2(\omega_{p(q)} + i\kappa/2)^{2}} \left\{ \frac{\kappa^{2}/2 - i\kappa\omega_{p(q)}}{(\omega_{p(q)}^{2} + \kappa^{2}/4)} \times [2\Gamma\cos(\omega_{p(q)}t) - \Gamma^{2} - 1] + [2i\Gamma\sin(\omega_{p(q)}t) - (2i\omega_{p(q)} - \kappa)t + \Gamma^{2} - 1] \right\}$$

$$(17)$$

with dissipative factor  $\Gamma = \exp(-\kappa t/2)$ . Using the similar method, we also calculate the overlap between theses two coherent states  $|\alpha_d\rangle$  and  $|\beta_d\rangle$  in the dissipative case as  $|\langle \beta_d | \alpha_d \rangle|^2 = \exp[-R_d(t)]$  with

$$\begin{split} R_{d}(t) &= (\Gamma^{2} + 1)[f(p)k_{1}^{2} + f(q)k_{2}^{2}] - 2\Gamma[f(p)k_{1}^{2}\cos(\omega_{p}t) \\ &+ f(q)k_{2}^{2}\cos(\omega_{q}t)] + 2M(\omega_{p}\omega_{q} + \kappa^{2}/4)\{\Gamma^{2}\cos(\omega_{d}t) \\ &- \Gamma[\cos(\omega_{p}t) + \cos(\omega_{q}t)] + 1\} - \kappa\omega_{d}M\{\Gamma^{2}\sin(\omega_{d}t) \\ &- \Gamma[\sin(\omega_{p}t) - \sin(\omega_{q}t)]\}, \end{split}$$
 (18)

where  $M = \omega_p \omega_q k_1 k_2 / [(\omega_p \omega_q + \kappa^2/4)^2 + \omega_d^2 \kappa^2/4]$  and  $f(x) = \omega_x^2 / (\omega_x^2 + \kappa^2/4)(x = p \text{ and } q)$ . In Fig. 2, we plot the time evolution of R(t) (non-dissipative case) in Eq. (10) and  $R_d(t)$  (dissipative case) in Eq. (18) under the different Rabi frequencies  $G_2$ . In non-dissipative case, as shown in Fig. 2(a), the exponent R(t)of the overlap between coherent states changed periodically and symmetrically, and it has the identical maximum value in each period, which means that the overlap  $|\langle \beta_d | \alpha_d \rangle|$  becomes minimum at these times. Besides,



**Figure 2.** The relationship between the exponent R(t) (non-dissipative case,  $\kappa=0$ ) and  $R_d(t)$  (dissipative case,  $\kappa=0.02$ ) and the time t is plotted in (a) and (b), respectively. The parameters are  $\lambda=1$ ,  $G_1=G_2$ ,  $\Delta_1=2G_2$ , and  $\theta=\pi/2$ . The red-solid line, blue-dashed line and black-dash-dotted line denote the case of  $G_2=1$ , 1.5 and 2, respectively.

we find that the larger the amount of Rabi frequency  $G_2$  is, the higher the maximal value of R(t) can reach, at the cost of the longer oscillation period. In dissipative case in Fig. 2(b), the symmetry and the periodicity on  $R_d(t)$  is destroyed, and the maximum values of the exponent  $R_d(t)$  decreases due to the dissipation effect. Combining this dissipative case with non-dissipative case, we come to the conclusion that the common feature that the small modulation overimposed on the larger modulation exists in both the non-dissipative and dissipative case, and the period of R(t) is partly determined by the Rabi frequency  $G_2$ . Therefore the generation of SCS can be optimized by adjusting the Rabi frequency  $G_3$ .

To visualize the time evolution of the mechanical phonon state, we firstly calculate the phase-space quasiprobability distribution using the Husimi Q function  $^{49,50}$  spanned by the dimensionless field quadratures  $\mathrm{Re}(\alpha_Q)$  and  $\mathrm{Im}(\alpha_Q)$ . The Husimi Q function is defined as  $Q(\alpha_Q)=|\langle\alpha_Q|\psi_p(t)\rangle|^2/\pi$  with  $\alpha_Q=x+iy$  the arbitrary complex number and  $\psi_p(t)$  the mechanical phonon state. In Fig. 3, we plot the Husimi Q function of DMR phonon state at different times as t=0,9,17,28,39,47,56, which correspond to the extreme time point of the exponent R(t) ( $G_2=1.5$ ) in the first period in Fig. 2(a). In the upper panel of Fig. 3 (the non-dissipative case with  $\kappa=0$ ), one can find that the DMR is initially prepared in the vacuum state, then gradually changed into the SCS, and finally evolve back into the vacuum state. In the presence of dissipation effect, i.e.,  $\kappa=0.02$ , as shown in the lower panel of the Fig. 3, the SCS can also be obtained with high-fidelity because the dissipation effect brings slight influence.

To show the interference fringes and nonclassicality of the mechanical SCS, the phase-space quasiprobability distribution using the Wigner function was also calculated in the phase space<sup>51</sup>, as shown in Fig. 4. In general, the quantum state could be judged to be nonclassical by checking that the Wigner function is negative in phase space. The Wigner function has the relation with the density operator  $\rho$  of the quantum state in the Fock state representation as  $W(\alpha_W) = 2 \operatorname{tr}[D^{-1}(\alpha_W)\rho D(\alpha_W)P]/\pi$  with the displacement operator  $D(\alpha_W) = \exp(\alpha_W a^\dagger - \alpha_W^* a)$  ( $\alpha_W = x + iy$  is the arbitrary complex number) and the parity operator  $P = \exp(i\pi a^\dagger a)^{52}$ . In Fock space, we have  $P|n\rangle = (-1)^n|n\rangle$ . Similar to the Husimi Q function, we plot the Wigner function on the SCS at the different times under the non-dissipative case (upper panel of the Fig. 4) and dissipative case (lower panel of the Fig. 4). One can find that the interference fringes appear under these two different cases, and it verifies the quantum features of the mechanical SCS of DMR due to the negative values of the Wigner function. Note that several experimental schemes<sup>53,54</sup> have been proposed to measure the Winger function with respect to the vibrational cat state.





Figure 3. (a–g) Represent the Husimi Q function  $(Q(\alpha_Q) = |\langle \alpha_Q | \psi_p(t) \rangle|^2/\pi$  with  $\alpha_Q = x + iy$  the arbitrary complex number and  $\psi_p(t)$  the mechanical phonon state) of SCS at the different times t=0,9,17,28,39,47,56. The upper and lower panel represent the non-dissipative case  $(\kappa=0)$  and dissipative case  $(\kappa=0.02)$ , respectively. The other parameters are the same as the Fig. 2(a).



**Figure 4.** (a–g) Represent the Wigner function  $(W(\alpha_W) = 2 \text{ tr} [D^{-1}(\alpha_W) \rho D(\alpha_W) P]/\pi$  with the displacement operator  $D(\alpha_W) = \exp(\alpha_W a^\dagger - \alpha_W^* a)$  ( $\alpha_W = x + iy$  is the arbitrary complex number), the density operator  $\rho$  of the mechanical phonon state and the parity operator  $P = \exp(i\pi a^\dagger a)$ ) of SCS at the different times t = 0, 9, 17, 28, 39, 47, 56. The upper and lower panel represent the non-dissipative case ( $\kappa = 0$ ) and dissipative case ( $\kappa = 0.02$ ), respectively. The other parameters are the same as the Fig. 2(a).

To more describe quantitatively the quantum features of the mechanical SCS of DMR, we employ the negative volume  $\delta$  of the Wigner function defined by Kenfack  $et~al.^{55}$ , as a quantitative measure of nonclassicality, where a non-zero value of  $\delta$  indicates nonclassicality. The negative volume associated with a quantum state  $|\psi\rangle$  can be written as

$$\delta = \iint [|W_{\psi}(x, p)| - W_{\psi}(x, p)] dxdp$$

$$= \iint |W_{\psi}(x, p)| dxdp - 1,$$
(19)

where

$$W_{\psi}(x, p) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} \langle x - y/2 | \psi \rangle \langle \psi | x + y/2 \rangle \exp(ipy) \, dy, \tag{20}$$

is the Wigner function of state  $|\psi\rangle$  in the coherent state representation, which satisfies the normalization condition  $\iint W_{\psi}(x,p)\,dxdp=1$ . It means that the larger the value of the negative volume  $\delta$  is, the stronger the classicality of the mechanical phonon state is. In Fig. 5, the negative volume  $\delta$  of Wigner function associated with the mechanical SCS is shown as a function of time t, where the non-zero value of  $\delta$  verifies the nonclassicality of the SCS, and the dissipation effect decreases the nonclassicality of the SCS during most of the evolution process.




Figure 5. The negative volume  $\delta$  of the Wigner function of SCS as a function of the time t. The red-dashed, blue-solid, and black-dashed lines represent the  $\kappa = 0$ ,  $\kappa = 0.02$ , and  $\kappa = 0.05$ , respectively. The other parameters are the same as the Fig. 4.

#### Discussion

We briefly address the experimental feasibility of our scheme. In general, the strain field could be produced by means of the thin piezoelectric  $^{11-17}$  or piezomagnetic  $^{56,57}$  film grown on the surface of the diamond layer, where, the piezoelectric or piezomagnetic film behaves as a transducer that transforms the signal between the external electric or magnetic field signal and the lattice vibration. Applying a voltage (magnetic field) across the piezoelectric film (piezomagnetic film), the strain field formed into the DMR. It implies that the strain field can be well controlled by the external electric or magnetic field. For a doubly clamped diamond beam with dimensions  $L\gg w$ , h, the coupling strength  $\lambda$  between the NV center and the DMR can be estimated by using the Euler-Bernoulli thin beam elasticity theory  $^{58}$ . If the NV center is located near the surface of the beam, we have  $\lambda/2\pi\approx 180\sqrt{\hbar/(L^3w\sqrt{\rho E})}$  GHz with  $\rho$  and E the mass density and the Young's modulus of diamond, respectively. Assuming that the dimension of the DMR is on the scale of  $\sim \mu m^{17}$  and the DMR's vibrational frequency  $\omega_m$  is about  $0.1\sim 10~{\rm GHz}^{18}$ , the coupling strength  $\lambda$  can be estimated as several kHz in our case. Besides, high-quality factor DMRs with  $Q=10^5-10^6$  have also been demonstrated in experiments  $^{21,22}$ , therefore the mechanical damping rate  $\kappa=\omega_m/Q$  can be less than 1 kHz, which could be further decreased by reducing the DMR's dimensions or improving the quality factor of DMR.

In addition, the preparation of the initial state in our scheme requires that the DMR is cool to the quantized ground state and the NV center is in the ground state, which can be realized in experiments. On one hand, several groups have demonstrated the mechanical resonator can be cooled to the quantum ground state  $^{39,61,62}$ . For example,  $^{61,62}$  is  $^{61,62}$ . ple, the experiment using conventional cryogenic refrigeration has demonstrated that the mechanical resonator can be cooled to its quantum ground state with the maximum phonon number  $\langle n_{\rm max} \rangle < 0.07$  and the probability >93%, through coupling the microwave-frequency mechanical oscillator to a superconducting phase qubit<sup>39</sup>. Additionally, recent work<sup>62</sup> has realized an approximately 10 MHz micromechanical oscillator cooling to the quantum ground state with the phonon occupation  $0.34 \pm 0.05$  using the sideband cooling technique, where the mechanical resonator is parametrically coupled to a superconducting microwave resonant circuit. Very recently, the topical review<sup>63</sup> summarized three theoretical phonon cooling protocols on the present DMR system in our model, established for both strain and magnetically mediated interactions between the orbital and spin degrees of freedom of NV center and DMR phonons. On the other hand, the ground state of the NV center is also derived experimentally<sup>14</sup>. In addition, the excellent coherence properties of NV center has been demonstrated in experiment with the spin relaxation time  $T_1 \sim 100 \,\text{seconds}^{64}$  and the spin dephasing time  $T_2 = 10 \,\text{milliseconds}^4$  at low temperature. Comparing with the decay of the DMR, the decay of the NV center is negligible. Furthermore, we emphasized that our work has the essential difference with the previous work<sup>26</sup>, where the coupling was achieved by means of a strong magnetic field gradient from a nearby magnetic tip, and the spatial quantum superposition state for the diamond's center-of-mass oscillation was generated via the magnetic field gradient induced the NV center electron spin and the mechanical oscillation coupling. Here we use intrinsic strain coupling between mechanical oscillator and the NV center to generate the macroscopic quantum superposition states. Larger magnetic field gradient is not needed.

For more technical aspects, we notice that a hybrid device that integrating a microscale diamond beam with a single embedded NV center spin to a superconducting coplanar waveguide cavity was proposed<sup>65</sup>, where the diamond beam phonon can strongly couple to the cavity photon via a dielectric interaction through an ac electric field. Based on this method, quantum state transfer between the mechanical mode and the cavity can be realized. For instance, if we map the mechanical phonon state into the cavity field completely, the Wigner function of SCS can be constructed by quantum homodyne tomography via detecting the cavity photons<sup>50,66,67</sup>. In addition, several theoretical and experimental schemes have been proposed to detect the cat state in different physical

systems, perhaps the most possible achievable measurement to our mechanical SCS is analogy of the experimental techniques applied in the system of trapped ion<sup>68,69</sup>, where the quantum mechanical interference between the localized wave packets of the motional states of ions can be directly measured by detecting the probability of the ion's internal state.

In summary, we utilize the cyclic  $\Delta$ -type transition structure in the DMR system to generate the nonclassical mechanical SCS through the dynamical strain-mediated coupling mechanism. Using this cyclic population transfer, the populations of the different energy levels of the NV center can be selectively transferred by controlling classical fields, and the SCS can be created by adjusting the controllable parameters of the system. Our scheme also provides another route for building a distributed QIP architecture, where each DMR unit acts as a quantum node, and quantum information is transferred and processed in and between the NV centers at spatially different DMR. It may be a significant step toward the future full-scale quantum-information processor based on increasingly developed nanoscale solid-state technology.

#### Method

The deduction of the Hamiltonian  $H_{+-}$  (Eqs (6 and 7)) using the FNT method. Firstly, we consider the sub-Hamiltonian  $H_u=\Delta_2\sigma_{-1,-1}+G_2(\sigma_{0,-1}+\sigma_{-1,0})$  containing the two lower states  $|0\rangle$  and  $|-1\rangle$  of the NV center. It is easy to derive its eigenstates as  $|+\rangle = \cos(\theta/2) |-1\rangle + \sin(\theta/2) |0\rangle$  and  $|-\rangle = -\sin(\theta/2) |-1\rangle$  $+\cos(\theta/2)$   $|0\rangle$  and the corresponding eigenvalues  $\varepsilon_{\pm}=\Delta_2/2\pm\omega$ . The mixing angle is  $\theta=\arctan[2G_2/\Delta_2]$  and the dressed frequency is  $\omega = \sqrt{G_2^2 + \Delta_2^2/4}$ .

Thus, we can rewritten the total Hamiltonian of the system H as  $H = H_f + H_i$  with

$$H_f = \Delta_1 \sigma_{+1,+1} + \varepsilon_+ \sigma_{+,+} + \varepsilon_- \sigma_{-,-} \tag{21}$$

and

$$H_i = \lambda_1 p \sigma_{+1,+} - \lambda_2 q \sigma_{+1,-} + H.c.,$$
 (22)

where  $p = a + k_1$  and  $q = a - k_2$  can be treated as the displaced boson operators. The other four controllable parameters are  $\lambda_1 = \lambda \cos(\theta/2)$ ,  $\lambda_2 = \lambda \sin(\theta/2)$ ,  $k_1 = G_1 \tan(\theta/2)/\lambda$ , and  $k_2 = G_1 \cot(\theta/2)/\lambda$ , respectively. Therefore, the effective  $\Lambda$ -like three-level transition of NV center has been established, as shown in Fig. 1c. The transition  $|+1\rangle \leftrightarrow |\pm\rangle$  are coupled to the displaced quantized phonon field with the operators p and q, respectively, and the corresponding coupling strengths are  $\lambda_1$  and  $\lambda_2$ . As a result, the quantized phonon field coupling to the transition  $|+1\rangle \leftrightarrow |-1\rangle$  has been transformed into two displaced quantized phonon field coupling two transition  $|+1\rangle \leftrightarrow |\pm\rangle$ by this unitary transformation.

In order to better understand the mechanism to prepare the nonclassical mechanical SCS, in the large detuning limit ( $\Delta_{\pm} \gg \lambda_{1(2)}$ ), we can separate the upper state  $|\pm 1\rangle$  from the two lower states  $|\pm \rangle$  and adiabatically eliminate the transitions  $|+1\rangle \leftrightarrow |\pm\rangle$  by using the FNT method<sup>48</sup>. We consider the total Hamiltonian H consisting of the free Hamiltonian  $H_f$  and the interaction Hamiltonian  $H_b$  and use the general perturbation theory to treat the interaction part  $H_i$  as a perturbation term comparing with the free part  $H_i$ . Then, an unitary transformation is used to the Hamiltonian H, and derived an effective Hamiltonian  $H_{eff}$  as  $H_{eff} = \exp(-O)H \exp(O)$  with O an anti-Hermitian operator. Further, by using the Baker-Campbell-Hausdorff formula<sup>70</sup>, the effective Hamiltonian  $H_{eff}$  could be simplified to

$$H_{eff} \cong H_f + [H_i, O]/2, \tag{23}$$

where the condition  $H_i + [H_{\beta} O] = 0$  has been used, which determines the form of anti-Hermitian operator O.

In general, the eigenvectors  $\{|i\rangle = |+1\rangle$ ,  $|\pm\rangle$  (corresponding to the eigenvalues  $E_i = \Delta_1$ ,  $\varepsilon_{\pm}$ ) for the free Hamiltonian  $H_i$  are known. Taking the matrix elements of the equation  $H_i + [H_i, O] = 0$  on the basis  $|i\rangle$ , we can obtain  $O = \eta_1 p \sigma_{+1,+} + \eta_2 q \sigma_{+1,-} - \eta_1 p^{\dagger} \sigma_{+,+1} - \eta_2 q^{\dagger} \sigma_{-,+1}$  with  $\eta_1 = -\lambda_1 / \Delta_+$ ,  $\eta_2 = \lambda_2 / \Delta_-$ , and  $\Delta_{\pm} = \Delta_1 - \varepsilon_{\pm}$ . Then, substituting the expression of  $H_{\mathfrak{S}}$   $H_{\mathfrak{b}}$  and O into the Eq. (23), the effective Hamiltonian  $H_{eff}$  can be

rewritten as  $H_{eff} = H_{11} + H_{+-}$  with

$$H_{11} = (\Delta_1 + \omega_p p p^{\dagger} + \omega_q q q^{\dagger}) \sigma_{+1,+1}, \tag{24}$$

and

$$H_{+-} = (\varepsilon_{+} - \omega_{p} p^{\dagger} p) \sigma_{+,+} + (\varepsilon_{-} - \omega_{q} q^{\dagger} q) \sigma_{-,-} + \eta(p^{\dagger} q \sigma_{+,-} + p q^{\dagger} \sigma_{-,+})$$
(25)

with the parameters  $\eta=\frac{2\Delta_1-\Delta_2}{2\Delta_+\Delta_-}\lambda_1\lambda_2$  and  $\omega_{p\,(q)}=\lambda_{\mathrm{I}(2)}^2/\Delta_{+(-)}$ . By substituting the expressions  $p=a+k_1$  and  $q=a-k_2$  into the Hamiltonian  $H_{11}$ , we can obtain  $H_{11}=H_d\sigma_{+1,+1}$  with  $H_d\cong(\omega_p+\omega_q)aa^\dagger+(k_1\omega_p-k_2\omega_q)(a+a^\dagger)$  describing a typical forced harmonic actilities

In addition, using the rotating wave approximation, we write the Hamiltonian  $H_{+-}$  as  $H_{+-} = H_{+}\sigma_{+,+} + H_{+,+}$  $H_{-}\sigma_{-,-}$  with

$$H_{+} = -\omega_{p}a^{\dagger}a - \omega_{p}k_{1}(a+a^{\dagger}) + \varepsilon_{+} - \omega_{p}^{2}k_{1}^{2}, \tag{26}$$

and

$$H_{-} = -\omega_{a}a^{\dagger}a + \omega_{a}k_{2}(a+a^{\dagger}) + \varepsilon_{-} - \omega_{a}^{2}k_{2}^{2}. \tag{27}$$

# References

- 1. Childress, L. et al. Coherent Dynamics of Coupled Electron and Nuclear Spin Qubits in Diamond. Science 314, 281-285 (2006).
- 2. Rondin, L. et al. Magnetometry with nitrogen-vacancy defects in diamond. Rep. Prog. Phys. 77, 056503 (2014).
- 3. Jelezko, F. et al. Observation of Coherent Oscillations in a Single Electron Spin. Phys. Rev. Lett. 92, 076401 (2004).
- 4. Balasubramanian, G. et al. Ultralong spin coherence time in isotopically engineered diamond. Nat. Mater. 8, 383–387 (2009).
- 5. Maurer, P. C. et al. Room-Temperature Quantum Bit Memory Exceeding One Second. Science 336, 1283-1286 (2012).
- 6. van der Sar, T. et al. Decoherence-protected quantum gates for a hybrid solid-state spin register. Nature 484, 82-86 (2012).
- 7. Yao, N. Y. et al. Scalable architecture for a room temperature solid-state quantum information processor. Nat. Commun. 3, 800 (2012).
- 8. Liang, Z.-T. et al. Proposal for implementing universal superadiabatic geometric quantum gates in nitrogen-vacancy centers. Phys. Rev. A 93, 040305(R) (2016).
- 9. Imamoğlu, A. Cavity QED Based on Collective Magnetic Dipole Coupling: Spin Ensembles as Hybrid Two-Level Systems. *Phys. Rev. Lett.* **102**, 083602 (2009).
- Yang, W. L. et al. Quantum simulation of an artificial Abelian gauge field using nitrogen-vacancy-center ensembles coupled to superconducting resonators. Phys. Rev. A 86, 012307 (2012).
- superconducting resonators. *Phys. Rev. A* **86**, 012307 (2012).

  11. Ovartchaiyapong, P. *et al.* Dynamic strain-mediated coupling of a single diamond spin to a mechanical resonator. *Nat. Commun.* **5**, 4429 (2014).
- Teissier, J. et al. Strain coupling of a nitrogen-vacancy center spin to a diamond mechanical oscillator. Phys. Rev. Lett. 113, 020503 (2014).
- 13. Barfuss, A. et al. Strong mechanical driving of a single electron spin. Nat. Phys. 11, 820-824 (2015).
- 14. MacQuarrie, E. R. et al. Mechanical spin control of nitrogen-vacancy centers in diamond. Phys. Rev. Lett. 111, 227602 (2013).
- 15. MacQuarrie, E. R. *et al.* Coherent control of a nitrogen-vacancy center spin ensemble with a diamond mechanical resonator. *Optica*. **2**, 233–238 (2015).
- 16. MacQuarrie, E. R. et al. Continuous dynamical decoupling of a single diamond nitrogen-vacancy center spin with a mechanical resonator. Phys. Rev. B 92, 224419 (2015).
- 17. Bennett, S. D. *et al.* Phonon-induced spin-spin interactions in diamond nanostructures: application to spin squeezing. *Phys. Rev. Lett.* **110**, 156402 (2013).
- 18. Kepesidis, K. V. et al. Phonon cooling and lasing with nitrogen-vacancy centers in diamond. Phys. Rev. B 88, 064105 (2013).
- Meesala, S. et al. Enhanced Strain Coupling of Nitrogen-Vacancy Spins to Nanoscale Diamond Cantilevers. Phys. Rev. Applied 5, 034010 (2016).
- 20. Xu, Q., Yang, W. L. & Yin, Z.-q. Phonon induced two-mode squeezing of nitrogen-vacancy center ensembles. arXiv:1512.08605 (2015).
- 21. Tao, Y., Boss, J. M., Moores, B. A. & Degen, C. L. Single-crystal diamond nanomechanical resonators with quality factors exceeding one million. *Nat. Commun.* 5, 3638 (2014).
- 22. Khanaliloo, B., Mitchell, M., Hryciw, A. C. & Barclay, P. E. High-Q/V Monolithic Diamond Microdisks Fabricated with Quasi-isotropic Etching. *Nano Letters* 15, 5131–5136 (2015).
- 23. Bose, S., Jacobs, K. & Knight, P. L. Preparation of nonclassical states in cavities with a moving mirror. *Phys. Rev. A* 56, 4175–4186 (1997).
- 24. Armour, A. D., Blencowe, M. P. & Schwab, K. C. Entanglement and Decoherence of a Micromechanical Resonator via Coupling to a Cooper-Pair Box. *Phys. Rev. Lett.* **88**, 148301 (2002).
- Romero-Isart, O. et al. Large Quantum Superpositions and Interference of Massive Nanometer-Sized Objects. Phys. Rev. Lett. 107, 020405 (2011).
   Yin, Z. Q., Li, T. C., Zhang, X. & Duan, L. M. Large quantum superpositions of a levitated nanodiamond through spin-
- optomechanical coupling. Phys. Rev. A 88, 033614 (2013).

  27. Marshall W. Simon C. Penrose R. & Bouwmeester D. Towards Quantum Superpositions of a Mirror Phys. Rev. Lett. 91, 130401
- 27. Marshall, W., Simon, C., Penrose, R. & Bouwmeester, D. Towards Quantum Superpositions of a Mirror. *Phys. Rev. Lett.* **91**, 130401 (2003).
- 28. Abdi, M. et al. Quantum State Engineering with Circuit Electromechanical Three-Body Interactions. Phys. Rev. Lett. 114, 173602 (2015).
- 29. Nielsen, M. A. & Chuang, I. L. Quantum Computation and Quantum Information (Cambridge, 2000).
- Zhu, S.-L., Wang, Z. D. & Zanardi, P. Geometric Quantum Computation and Multiqubit Entanglement with Superconducting Qubits inside a Cavity. Phys. Rev. Lett. 94, 100502 (2005).
- 31. Yu, L. B. *et al.* Implementing multi-qubit entanglement of two-level systems inside a superconducting phase qubit. *Euro. Phys. J. D.* **61**, 499–505 (2011).
- 32. Gangat, A. A. Observing controlled state collapse in a single mechanical oscillator via a direct probe of energy variance. *Phys. Rev.* A 88, 063846 (2013).
- 33. Kippenberg, T. J. & Vahala, K. J. Cavity Optomechanics: Back-Action at the Mesoscale. Science 321, 1172–1176 (2008).
- Vlastakis, B. et al. Deterministically Encoding Quantum Information Using 100-Photon Schröinger Cat States. Science 342, 607-610 (2013).
- 35. Leibfried, D. et al. Creation of a six-atom 'Schröinger cat' state. Nature 438, 639-642 (2005).
- 36. Raimond, J. M., Brune, M. & Haroche, S. Reversible Decoherence of a Mesoscopic Superposition of Field States. *Phys. Rev. Lett.* **79**, 1964–1967 (1997).
- 37. Ourjoumtsev, A. et al. Generation of optical 'Schröinger cats' from photon number states. Nature 448, 784-786 (2007).
- 38. Huang, X. M. H. et al. Nanoelectromechanical systems: Nanodevice motion at microwave frequencies. Nature 421, 496 (2003).
- 39. O'Connell, A. D. et al. Quantum ground state and single-phonon control of a mechanical resonator. Nature 464, 697-703 (2010).
- Liao, J.-Q., Huang, J.-F. & Tian, L. Generation of macroscopic Schrödinger's cat states in qubit-oscillator systems. Phys. Rev. A 93, 033853 (2016).
- 41. Peixoto de Faria, J. G. & Nemes, M. C. Dissipative dynamics of the Jaynes-Cummings model in the dispersive approximation: analytical results. *Phys. Rev. A* **59**, 3918–3925 (1999).
- 42. Song, W.-l. *et al.* One-step generation of multipartite entanglement among nitrogen-vacancy center ensembles. *Sci. Rep.* **5**, 7755 (2015).
- 43. Maze, J. R. Gali *et al.* Properties of nitrogen-vacancy centers in diamond: the group theoretic approach. *New J. Phys.* **13**, 025025 (2011).
- 44. Doherty, M. W. et al. Theory of the ground-state spin of the NV<sup>-</sup> center in diamond. Phys. Rev. B 85, 205203 (2012).
- 45. Fuchs, G. D. et al. Gigahertz Dynamics of a Strongly Driven Single Quantum Spin. Science 326, 1520-1522 (2009).
- 46. Yale, C. G. et al. All-optical control of a solid-state spin using coherent dark states. Proc. Natl. Acad. Sci. USA 110, 7595–7600 (2013).
- 47. Dolde, F. et al. Electric-field sensing using single diamond spins. Nat. Phys. 7, 459-463 (2011).

- 48. Fröhlich, H. Theory of the superconducting state. I. the ground state at the absolute zero of temperature. *Phys. Rev.* **79**, 845–856 (1950).
- 49. Kôdi, H. Some formal properties of the density matrix. Proc. Phys. Math. Soc. Jpn. 22, 264-314 (1940).
- 50. Kirchmair, G. et al. Observation of quantum state collapse and revival due to the single-photon Kerr effect. Nature 495, 205-209 (2013)
- 51. Wigner, E. P. On the Quantum Correction For Thermodynamic Equilibrium. Phy. Rev. 40, 749-759 (1932).
- 52. Cahill, K. E. & Glauber, R. J. Density Operators and Quasiprobability Distributions. Phys. Rev. 177, 1882-1902 (1969).
- 53. Leibfried, D. et al. Experimental Determination of the Motional Quantum State of a Trapped Atom. Phys. Rev. Lett. 77, 4281–4285 (1996).
- 54. Lutterbach, L. G. & Davidovich, L. Method for Direct Measurement of the Wigner Function in Cavity QED and Ion Traps. *Phys. Rev. Lett.* **78**, 2547–2550 (1997).
- 55. Kenfack, A. & Zyczkowski, K. Negativity of the Wigner function as an indicator of non-classicality. *J. Opt. B*, *Quantum Semiclass. Opt.* 6, 396–404 (2004).
- 56. Cai, J., Jelezko, F. & Plénio, M. B. Hybrid sensors based on colour centres in diamond and piezoactive layers. *Nat. Commun.* 5, 4065 (2014).
- 57. Schuetz, M. J. A. et al. Universal Quantum Transducers Based on Surface Acoustic Waves. Phys. Rev. X 5, 031031 (2015).
- 58. Landau, L. D. & Lifshitz, E. M. Theory of Elasticity (Butterworth-Heinemann, Oxford, 1986).
- Soykal, Ö. O., Ruskov, R. & Tahan, C. Sound-Based Analogue of Cavity Quantum Electrodynamics in Silicon. Phys. Rev. Lett. 107, 235502 (2011).
- 60. Albrecht, A., Retzker, A., Jelezko, F. & Plenio, M. B. Coupling of nitrogen vacancy centres in nanodiamonds by means of phonons. *New J. Phys.* 15, 083014 (2013).
- 61. Wilson-Rae, I., Zoller, P. & Imamoğlu, A. Laser Cooling of a Nanomechanical Resonator Mode to its Quantum Ground State. *Phys. Rev. Lett.* **92**, 075507 (2004).
- 62. Teufel, J. D. et al. Sideband cooling of micromechanical motion to the quantum ground state. Nature 475, 359-363 (2011).
- 63. Lee, D. et al. Topical Review: Spins and mechanics in diamond. arXiv:1609.00418 (2016).
- 64. Jarmola, A. et al. Temperature- and Magnetic-Field-Dependent Longitudinal Spin Relaxation in Nitrogen-Vacancy Ensembles in Diamond. Phys. Rev. Lett. 108, 197601 (2012).
- 65. Li, P.-B. et al. Hybrid quantum device based on NV centers in diamond nanomechanical resonators plus superconducting waveguide cavities. Phys. Rev. Applied 4, 044003 (2015).
- 66. Hofheinz, M. et al. Synthesizing arbitrary quantum states in a superconducting resonator. Nature 459, 546-549 (2009).
- 67. Bertet, P. et al. Direct measurement of the Wigner function of a one-photon Fock state in a cavity. Phys. Rev. Lett. 89, 200402 (2002).
- 68. Monroe, C., Meekhof, D. M., King, B. E. & Wineland, D. J. A. "Schrödinger cat" superposition state of an atom. Science 272, 1131–1136 (1996).
- 69. Kienzler, D. et al. Observation of quantum interference between separated mechanical oscillator wave packets. Phys. Rev. Lett. 116, 140402 (2016).
- 70. Witschel, W. Ordered products of exponential operators by similarity transformations. Int. J. Quantum Chem. 20, 1233-1241 (1981).

# **Acknowledgements**

We thank Shi-Liang Zhu for enlightening discussions. This work was supported by the NKRDP of China (Grant No. 2016YFA0301803), the NSFC (Grant No. 11474153), and the PCSIRT (Grant No. IRT1243). W. L. Yang was supported by the NSFC (Grants No. 11574353 and No. 11274351). Z. Q. Yin was supported by the NSFC (Grants No. 61435007 and No. 11474177).

#### **Author Contributions**

H.Q.Z. and Y.W.L. conceive the idea and carry out the research. H.Q.Z., Y.W.L., C.C.Y. and Y.Z.Q. discuss the results and write the manuscript.

# **Additional Information**

**Competing financial interests:** The authors declare no competing financial interests.

How to cite this article: Hou, Q. et al. Generation of macroscopic Schrödinger cat state in diamond mechanical resonator. Sci. Rep. 6, 37542; doi: 10.1038/srep37542 (2016).

**Publisher's note:** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit http://creativecommons.org/licenses/by/4.0/

© The Author(s) 2016