

# **OPEN** Generalised monogamy relation of convex-roof extended negativity in multi-level systems

Received: 29 June 2016 Accepted: 19 October 2016 Published: 18 November 2016

Tian Tian, Yu Luo & Yongming Li

In this paper, we investigate the generalised monogamy inequalities of convex-roof extended negativity (CREN) in multi-level systems. The generalised monogamy inequalities provide the upper and lower bounds of bipartite entanglement, which are obtained by using CREN and the CREN of assistance (CRENOA). Furthermore, we show that the CREN of multi-qubit pure states satisfies some monogamy relations. Additionally, we test the generalised monogamy inequalities for qudits by considering the partially coherent superposition of a generalised W-class state in a vacuum and show that the generalised monogamy inequalities are satisfied in this case as well.

Quantum entanglement is one of the most important physical resources in quantum information processing 1-4. As distinguished from classical correlations, quantum entanglement cannot be freely shared among many objects. We call this important phenomenon of quantum entanglement monogamy<sup>5,6</sup>. The property of monogamy may be as fundamental as the no-cloning theorem<sup>7</sup>, which gives rise to structures of entanglement in multipartite settings<sup>8,9</sup>. Some monogamy inequalities have been studied to apply entanglement to more useful quantum information processing. The property of monogamy property has been considered in many areas of physics: it can be used to extract an estimate of the quantity of information about a secret key captured by an eavesdropper in quantum cryptography<sup>10,11</sup>, as well as the frustration effects observed in condensed matter physics<sup>12,13</sup> and even black-hole physics<sup>14,15</sup>

The monogamy relation of entanglement is a way to characterise different types of entanglement distribution. The first monogamy relation was named the Coffman-Kundu-Wootters (CKW) inequality<sup>8</sup>. The monogamy property can be interpreted as the following statement: the amount of entanglement between A and B plus the amount of entanglement between A and C cannot be greater than the amount of entanglement between A and the BC pair. Osborne and Verstraete later proved that the CKW inequality also holds in an n-qubit system<sup>9</sup>. Other types of monogamy relations for entanglement were also proposed. Studies have found that the monogamy inequality holds in terms of some entanglement measures, negativity<sup>16</sup>, squared CREN<sup>17</sup>, entanglement of formation<sup>18–20</sup>, Rényi entropy<sup>21</sup> and Tsallis entropy<sup>22,23</sup>. The monogamy property of other physical resources, such as discord and steering<sup>24</sup>, has also been discussed. There can be several inequivalent types of entanglement among the subsystems in multipartite quantum systems, and the amount of different types of entanglement might not be directly comparable to one another. Regula et al. studied multi-party quantum entanglement and found that there was strong monogamy<sup>25</sup>. Additionally, generalised monogamy relations of concurrence for N-qubit systems were also proposed by Zhu et al.26.

In this paper, we study the generalised monogamy inequalities of CREN in multi-qubit systems. We first recall some basic concepts of entanglement measures. Then, monogamy inequalities are given by the concurrence and negativity of the *n*-qubit entanglement. Furthermore, we consider some states in a higher-dimensional quantum system and find that the generalised monogamy inequalities also hold for these states. We specifically test the generalised monogamy inequalities for qudits by considering the partially coherent superposition of a generalised W-class state in a vacuum, and we show that the generalised monogamy inequalities are satisfied in this case as well. These relations also give rise to a type of trade-off in inequalities that is related to the upper and lower bounds of CRENOA. It shows the bipartite entanglement between AB and the other qubits: especially under partition AB, a two-qubit system is different from the previous monogamy inequality that is typically used.

College of Computer Science, Shaanxi Normal University, Xi'an, 710062, China. Correspondence and requests for materials should be addressed to Y.Li. (email: liyonqm@snnu.edu.cn)

#### Results

This paper is organised as follows: in the first subsection, we recall some basic concepts of concurrence and negativity. We present the monogamy relations of concurrence and negativity in the second subsection. In the third subsection, the generalised monogamy inequalities of CREN are given. The fourth subsection includes some examples that verify these results.

**Preliminaries: concurrence and negativity.** For any bipartite pure state  $|\psi\rangle_{AB}$  in a  $d\otimes d'$   $(d\leq d')$  quantum system with its Schmidt decomposition,

$$|\psi\rangle_{AB} = \sum_{i=0}^{d-1} \sqrt{\lambda_i} |ii\rangle, \quad \lambda_i \ge 0, \sum_{i=0}^{d-1} \lambda_i = 1,$$
 (1)

the concurrence  $\mathcal{C}ig(\ket{\psi}_{AB}ig)$  is defined as  $^{27}$ 

$$C(|\psi\rangle_{AB}) = \sqrt{2[1 - \text{Tr}(\rho_A^2)]}, \qquad (2)$$

where  $\rho_A = tr_B(|\psi\rangle_{AB}\langle\psi|)$ . For any mixed state  $\rho_{AB}$ , its concurrence is defined as

$$C(\rho_{AB}) = \min \sum_{i} p_{i} C(|\psi_{i}\rangle_{AB}), \tag{3}$$

where the minimum is taken over all possible pure state decompositions  $\{p_i, |\psi_i\rangle_{AB}\}$  of  $\rho_{AB}$ . Similarly, the concurrence of assistance (COA) of  $\rho_{AB}$  is defined as<sup>28</sup>

$$C_a(\rho_{AB}) = \max \sum_i p_i C(|\psi_i\rangle_{AB}), \tag{4}$$

where the maximum is taken over all possible pure state decompositions  $\{p_i, |\psi_i\rangle_{AB}\}$  of  $\rho_{AB}$ .

Another well-known quantification of bipartite entanglement is negativity. For any bipartite pure state  $|\psi\rangle_{AB}$ , the negativity  $\mathcal{N}(|\psi\rangle_{AB})$  is

$$\mathcal{N}(|\psi\rangle_{AB}) = 2\sum_{i < j} \sqrt{\lambda_i \lambda_j} = (\text{Tr}_{\sqrt{\rho_A}})^2 - 1, \tag{5}$$

where  $\rho_A = tr_B(|\psi\rangle_{AB}\langle\psi|)$ .

For any bipartite state  $ho_{AB}$  in the Hilbert space  $\mathcal{H}_{\mathcal{A}}\otimes\mathcal{H}_{\mathcal{B}}$  negativity is defined as  $^{29}$ 

$$\mathcal{N}(\rho_{AB}) = \frac{\|\rho_{AB}^{T_A}\| - 1}{2},\tag{6}$$

where  $\rho_{AB}^{T_A}$  is a partial transposition with respect to the subsystem A, ||X|| denotes the trace norm of X; i.e.,  $||X|| \equiv \text{Tr}\sqrt{XX^{\dagger}}$ . Negativity is a computable measure of entanglement, which is a convex function of  $\rho_{AB}$ . It disappears if, and only if,  $\rho_{AB}$  is separable for the  $2 \otimes 2$  and  $2 \otimes 3$  systems<sup>30</sup>. For the purposes of this discussion, we use the following definition of negativity:

$$\mathcal{N}(\rho_{AB}) = \|\rho_{AB}^{T_A}\| - 1. \tag{7}$$

For any maximally entangled state in a two-qubit system, this negativity is equal to 1. CREN gives a perfect discrimination of positive partial transposition-bound entangled states and separable states in any bipartite quantum system  $^{31,32}$ . For any mixed state  $\rho_{AB}$ , CREN is defined as

$$\mathcal{N}_{c}(\rho_{AB}) = \min \sum_{i} p_{i} \mathcal{N}(|\psi_{i}\rangle_{AB}), \tag{8}$$

where the minimum is taken over all possible pure state decompositions  $\{p_i, |\psi_i\rangle_{AB}\}$  of  $\rho_{AB}$ . For any mixed state  $\rho_{AB}$ , CRENOA is defined as  $^{17}$ 

$$\mathcal{N}_a(\rho_{AB}) = \max \sum_i p_i \mathcal{N}(|\psi_i\rangle_{AB}),\tag{9}$$

where the maximum is taken over all possible pure state decompositions  $\{p_i, |\psi_i\rangle_{AB}\}$  of  $\rho_{AB}$ .

CREN is equivalent to concurrence for any pure state with Schmidt rank- $2^{17}$ , and consequently, it follows that for any two-qubit mixed state  $\rho_{AB} = \sum_i p_i |\psi_i\rangle\langle\psi_i|$ :

$$\mathcal{N}_{c}(\rho_{AB}) = \min \sum_{i} p_{i} \mathcal{N}(|\psi_{i}\rangle_{AB}) = \min \sum_{i} p_{i} \mathcal{C}(|\psi_{i}\rangle_{AB}) = \mathcal{C}(\rho_{AB})$$
(10)

and

$$\mathcal{N}_{a}(\rho_{AB}) = \max \sum_{i} p_{i} \mathcal{N}(|\psi_{i}\rangle_{AB}) = \max \sum_{i} p_{i} \mathcal{C}(|\psi_{i}\rangle_{AB}) = \mathcal{C}_{a}(\rho_{AB}), \tag{11}$$

where the minimum and the maximum are taken over all pure state decompositions  $\{p_i, |\psi_i\rangle_{AB}\}$  of  $\rho_{AB}$ .

Monogamy relations of concurrence and negativity. The CKW inequality<sup>8</sup> was first defined as

$$\mathcal{C}^2(\rho_{A|BC}) \ge \mathcal{C}^2(\rho_{AB}) + \mathcal{C}^2(\rho_{AC}),\tag{12}$$

where  $C(\rho_{A|BC})$  is the concurrence of a three-qubit state  $\rho_{A|BC}$  for any bipartite cut of subsystems between A and BC. Similarly, the dual inequality in terms of COA is as follows<sup>33</sup>:

$$\mathcal{C}^2(\rho_{A|BC}) \le \mathcal{C}_a^2(\rho_{AB}) + \mathcal{C}_a^2(\rho_{AC}). \tag{13}$$

For any pure state  $|\psi\rangle_{A_1...An}$  in an n-qubit system  $A_1\otimes...\otimes A_n$ , where  $A_i\cong C^2$  for i=1,...,n, a generalisation of the CKW inequality is

$$C^{2}(|\psi\rangle_{A_{1}|A_{2}...A_{n}}) \ge C^{2}(\rho_{A_{1}A_{2}}) + ... + C^{2}(\rho_{A_{1}A_{n}}).$$
(14)

The dual inequality in terms of the COA for n-qubit states has the form<sup>17</sup>

$$C^{2}(|\psi\rangle_{A_{1}|A_{2}...A_{n}}) \leq C_{a}^{2}(\rho_{A_{1}A_{2}}) + ... + C_{a}^{2}(\rho_{A_{1}A_{n}}).$$
(15)

when the rank of the matrix is 2, we have

$$C(|\psi\rangle_{A_1|A_2...A_n}) = \mathcal{N}(|\psi\rangle_{A_1|A_2...A_n}). \tag{16}$$

Combining Eq. (10) with Eq. (11), we have

$$\mathcal{C}(\rho_{A_i A_j}) = \mathcal{N}_c(\rho_{A_i A_j}), \, \mathcal{C}_a(\rho_{A_i A_j}) = \mathcal{N}_a(\rho_{A_i A_j}), \tag{17}$$

where  $i, j \in \{1, ..., n\}, i \neq j$ .

For any *n*-qubit pure state  $|\psi\rangle_{A_{n-A}n}$ , we have

$$\mathcal{N}^{2}(|\psi\rangle_{A_{1}|A_{2}...A_{n}}) \ge \mathcal{N}_{c}^{2}(\rho_{A_{1}A_{2}}) + ... + \mathcal{N}_{c}^{2}(\rho_{A_{1}A_{n}}). \tag{18}$$

The dual inequality<sup>17</sup> in terms of CRENOA is as follows:

$$\mathcal{N}^{2}(|\psi\rangle_{A_{1}|A_{2}...A_{n}}) \leq \mathcal{N}^{2}_{a}(\rho_{A_{1}A_{2}}) + ... + \mathcal{N}^{2}_{a}(\rho_{A_{1}A_{n}}). \tag{19}$$

**Monogamy inequalities of CREN.** For a  $2\otimes 2\otimes m$  quantum pure state  $|\psi\rangle_{ABC}$ , it has been shown that  $\mathcal{C}_a^2(\rho_{AB})=\mathcal{C}^2(\rho_{AB})+\tau_2^C(|\psi\rangle_{ABC})^{33}$ , where  $\tau_2^C(|\psi\rangle_{ABC})=\mathcal{C}^2(|\psi\rangle_{A|BC})-\mathcal{C}^2(\rho_{AB})-\mathcal{C}^2(\rho_{AC})$  is the three-tangle of concurrence.  $\mathcal{C}(|\psi\rangle_{A|BC})$  is the concurrence under bipartition A|BC for pure state  $|\psi\rangle_{ABC}$ . Namely,

$$C^{2}(|\psi\rangle_{A|BC_{1}...C_{n-2}}) = C_{a}^{2}(\rho_{AB}) + C^{2}(\rho_{A|C_{1}...C_{n-2}}).$$
(20)

Similarly, considering that CREN is equivalent to concurrence by Eq. (17), we have

$$\mathcal{N}^{2}(|\psi\rangle_{A|BC_{1}...C_{n-2}}) = \mathcal{N}^{2}_{a}(\rho_{AB}) + \mathcal{N}^{2}_{c}(\rho_{A|C_{1}...C_{n-2}}). \tag{21}$$

The concurrence is related to the linear entropy of a state<sup>34</sup>

$$T(\rho) = 1 - \text{Tr}(\rho^2). \tag{22}$$

Given a bipartite state  $\rho$ ,  $T(\rho)$  has the property<sup>35</sup>,

$$T(\rho_A) + T(\rho_B) \ge T(\rho_{AB}) \ge |T(\rho_A) - T(\rho_B)|. \tag{23}$$

From the definition of pure state concurrence in Eq. (2) together with Eq. (22), we have

$$C^{2}(|\psi\rangle_{AB|C_{1}...C_{n-2}}) = 2[1 - \text{Tr}(\rho_{AB}^{2})] = 2T(\rho_{AB}).$$
(24)

Now, we provide the following theorems:

Theorem 1. For any  $2 \otimes 2 \otimes 2$  tripartite mixed state  $\rho_{ABC}$  we have

$$\mathcal{N}_{a}^{2}(\rho_{A|BC}) \le \mathcal{N}_{a}^{2}(\rho_{B|AC}) + \mathcal{N}_{a}^{2}(\rho_{C|AB}).$$
 (25)

Proof. Let  $\rho_{ABC} = \sum_i p_i |\psi_i\rangle_{ABC} \langle \psi_i|$  be an optimal decomposition realising  $\mathcal{N}_a(\rho_{A|BC})$ ; that is,

Figure 1. The example shows the reciprocal relation of squared CRENOA in a four-qubit system.

$$\mathcal{N}_a(\rho_{A|BC}) = \max_i p_i \mathcal{N}(|\psi_i\rangle_{A|BC}), \tag{26}$$

where  $\rho_{BC} = \text{Tr}_A |\psi_i\rangle_{ABC} \langle\psi_i|$ ,  $\rho_B = \text{Tr}_{AC} |\psi_i\rangle_{ABC} \langle\psi_i|$  and  $\rho_C = \text{Tr}_{AB} |\psi_i\rangle_{ABC} \langle\psi_i|$ , and we have

$$\mathcal{N}^{2}(|\psi_{i}\rangle_{A|BC}) = \mathcal{C}^{2}(|\psi_{i}\rangle_{A|BC}) = 2T(\rho_{A}) = 2T(\rho_{BC}). \tag{27}$$

Combining Eq. (23) with Eq. (24), we have

$$2T(\rho_{BC}) \leq 2T(\rho_{B}) + 2T(\rho_{C})$$

$$= \mathcal{C}^{2}(|\psi_{i}\rangle_{B|AC}) + \mathcal{C}^{2}(|\psi_{i}\rangle_{C|AB})$$

$$= \mathcal{N}^{2}(|\psi_{i}\rangle_{B|AC}) + \mathcal{N}^{2}(|\psi_{i}\rangle_{C|AB}). \tag{28}$$

The third equality holds because CREN and concurrence are equal for any rank-2 pure state. Therefore, we obtain

$$\mathcal{N}^{2}(|\psi_{i}\rangle_{A|BC}) \leq \mathcal{N}^{2}(|\psi_{i}\rangle_{B|AC}) + \mathcal{N}^{2}(|\psi_{i}\rangle_{C|AB}). \tag{29}$$

Combining Eq. (26) with Eq. (29), we finally get

$$\mathcal{N}_{a}^{2}(\rho_{A|BC}) \le \mathcal{N}_{a}^{2}(\rho_{B|AC}) + \mathcal{N}_{a}^{2}(\rho_{C|AB}).$$
 (30)

Thus, the proof is completed.

Theorem 1 shows a simple relationship of CRENOA in a tripartite quantum system. The monogamy inequality shows that the entanglement A|BC cannot be greater than the sum of the entanglement B|AC and the entanglement C|AB. Taking an easy example, when considering a three-qubit state, the following equation exists:  $|\psi\rangle_{ABC}=a|010\rangle+b|100\rangle$  where  $|a|^2+|b|^2=1$ . Using a simple calculation, the following equation can be obtained:  $\mathcal{N}_a^2(\rho_{A|BC})=\mathcal{N}_a^2(\rho_{B|AC})+\mathcal{N}_a^2(\rho_{C|AB})$  where the state  $|\psi\rangle_{ABC}$  saturates the monogamy inequality in Eq. (25). Moreover, the iteration of Eq. (25) leads us to the generalized monogamy inequality in multi-qubit quantum systems.

Corollary 1. For any multi-party mixed state  $\rho_{A_1|A_2\dots A_n}$  in an n-qubit system<sup>36</sup>, the following monogamy inequality exists:

$$\mathcal{N}_{a}^{2}(\rho_{A_{1}|A_{2}...A_{n}}) \leq \sum_{i=2}^{n} \mathcal{N}_{a}^{2}(\rho_{A_{i}|A_{1}...A_{i-1}A_{i+1}...A_{n}}) \leq \sum_{i=2}^{n} \sum_{j=1, j \neq i}^{n} \mathcal{N}_{a}^{2}(\rho_{A_{i}A_{j}}). \tag{31}$$

The meaning of the first inequality is clear the bipartite entanglement between  $\rho_{A_1}$  and the other qubits, when taken as a group cannot be greater than the sum of the n-1 individual bipartite entanglements between  $\rho_{A_i}$  ( $i\neq 1$ ) and the other remaining qubits. We now start to consider a four-qubit system. As shown in Fig. 1, the squared CRENOA with respect to the bipartition (A|BCD) is not greater than the sum of the three squared CRENOAs (the three possible bipartitions are B|ACD, C|ABD and D|ABC).

The meaning of the second inequality is clear the sum of the bipartite entanglements between  $\rho_{A_i}$   $(i \neq 1)$  and the other remaining qubits cannot be greater than the sum of the bipartite entanglements  $\rho_{A_iA_i}$   $(i \neq 1, j \neq i)$ .

Theorem 2. For any *n*-qubit pure state  $|\psi\rangle_{ABC_1...C_{n-2}}$ , we have

$$2\mathcal{N}_{a}^{2}(\rho_{AB}) + \sum_{i=1}^{n-2} \mathcal{N}_{a}^{2}(\rho_{AC_{i}}) + \sum_{i=1}^{n-2} \mathcal{N}_{a}^{2}(\rho_{BC_{i}}) \ge \mathcal{N}^{2}(|\psi\rangle_{AB|C_{1}...C_{n-2}}), \tag{32}$$

where  $\rho_{AB}=\operatorname{Tr}_{C_1\dots C_{n-2}}(|\psi\rangle\langle\psi|)$ ,  $\rho_{AC_i}=\operatorname{Tr}_{BC_1\dots C_{i-1}C_{i+1}\dots C_{n-2}}(|\psi\rangle\langle\psi|)$  and  $\rho_{BC_i}=\operatorname{Tr}_{AC_1\dots C_{i-1}C_{i+1}\dots C_{n-2}}(|\psi\rangle\langle\psi|)$ . Proof. From the result of Theorem 1, we find that the generalised monogamy inequality can be easily obtained by using the superposition of states. We now consider  $\mathcal{N}^2(|\psi\rangle_{AB|C_1\dots C_{n-2}})$ . When the rank of the matrix is 2, we have

$$\mathcal{N}^{2}(|\psi\rangle_{AB|C_{1}...C_{n-2}}) = \mathcal{C}^{2}(|\psi\rangle_{AB|C_{1}...C_{n-2}}) = 2T(\rho_{AB}). \tag{33}$$

Combining Eq. (23) with Eq. (24), we get the relationship

$$2T(\rho_{AB}) \leq 2T(\rho_{A}) + 2T(\rho_{B})$$

$$= C^{2}(|\psi_{i}\rangle_{A|BC_{1}...C_{n-2}}) + C^{2}(|\psi_{i}\rangle_{B|AC_{1}...C_{n-2}})$$

$$= \mathcal{N}^{2}(|\psi_{i}\rangle_{A|BC_{1}...C_{n-2}}) + \mathcal{N}^{2}(|\psi_{i}\rangle_{B|AC_{1}...C_{n-2}}).$$
(34)

The third equality follows from the fact that CREN and concurrence are equal for any rank-2 pure state.

$$\mathcal{N}^{2}(|\psi_{i}\rangle_{AB|C_{1}...C_{n-2}}) \leq \mathcal{N}^{2}(|\psi_{i}\rangle_{A|BC_{1}...C_{n-2}}) + \mathcal{N}^{2}(|\psi_{i}\rangle_{B|AC_{1}...C_{n-2}}).$$
(35)

For a mixed state, CRENOA is expressed as  $\mathcal{N}(|\psi_i\rangle_{A|BC_1...C_{n-2}})$  , and we have

$$\mathcal{N}_{a}(\rho_{A|BC_{1}...C_{n-2}}) = \max_{i} \sum_{i} p_{i} \mathcal{N}(|\psi_{i}\rangle_{A|BC_{1}...C_{n-2}}).$$
 (36)

Furthermore, when combining this with Eq. (35), we finally get

$$\mathcal{N}^{2}(|\psi\rangle_{AB|C_{1}...C_{n-2}}) \leq \mathcal{N}^{2}_{a}(\rho_{A|BC_{1}...C_{n-2}}) + \mathcal{N}^{2}_{a}(\rho_{B|AC_{1}...C_{n-2}})$$
(37)

and

$$\mathcal{N}_{a}^{2}(\rho_{A|BC_{1}...C_{n-2}}) \leq \mathcal{N}_{a}^{2}(\rho_{AB}) + \sum_{i=1}^{n-2} \mathcal{N}_{a}^{2}(\rho_{AC_{i}}),$$

$$\mathcal{N}_{a}^{2}(\rho_{B|AC_{1}...C_{n-2}}) \leq \mathcal{N}_{a}^{2}(\rho_{BA}) + \sum_{i=1}^{n-2} \mathcal{N}_{a}^{2}(\rho_{BC_{i}}).$$
(38)

Combining Eq. (37) with Eq. (38), we have Eq. (32). In other words, we give an upper bound about  $\mathcal{N}^2(|\psi\rangle_{AB|C_1...C_{n-2}})$ , i.e.,

$$2\mathcal{N}_{a}^{2}(\rho_{AB}) + \sum_{i=1}^{n-2} \mathcal{N}_{a}^{2}(\rho_{AC_{i}}) + \sum_{i=1}^{n-2} \mathcal{N}_{a}^{2}(\rho_{BC_{i}}) \ge \mathcal{N}^{2}(|\psi\rangle_{AB|C_{1}...C_{n-2}}).$$
(39)

This completes the proof.

Theorem 2 shows that the entanglement between AB and the other qubits cannot be greater than the sum of the individual entanglements between A and each of the n-1 remaining qubits and the individual entanglements between B and each of the n-1 remaining qubits. Theorem 2 provides a polygamy-type upper bound of multi-qubit entanglement between the two-qubit system AB and the other (n-2)-qubit system  $C_1C_2...C_{n-2}$  in terms of the squared CRENOA. Especially under partition AB, a two-qubit system is different from the previous monogamy inequality. When  $|\psi\rangle_{AB|C_1...C_{n-2}} = |\psi\rangle_A \otimes |\psi\rangle_{B|C_1...C_{n-2}}$ , the calculation results in  $\mathcal{N}_a^2(\rho_{AB}) = 0$ ,  $\mathcal{N}_a^2(\rho_{AC_i}) = 0$ . Consequently, the polygamy-type relation is obtained as shown in Eq. (19).

Finally, consider the following four-qubit state:  $|\psi\rangle_{ABCD} = a|0100\rangle + b|0010\rangle + c|0001\rangle$  where  $|a|^2 + |b|^2 + |c|^2 = 1$ . We can easily get the following equations:  $\mathcal{N}_a^2(\rho_{AB}) = \mathcal{N}_a^2(\rho_{AC}) = \mathcal{N}_a^2(\rho_{AD}) = 0$  and  $\mathcal{N}_a^2(\rho_{BC}) + \mathcal{N}_a^2(\rho_{BD}) = \mathcal{N}^2(|\psi\rangle_{AB|CD}) = \frac{16}{9}$ . Therefore, the state  $|\psi\rangle_{ABCD}$  saturates the monogamy inequality in Eq. (32).

Theorem 3. For any *n*-qubit pure state  $|\psi\rangle_{ABC_1...C_{n-2}}$ ,

$$\mathcal{N}^{2}(|\psi\rangle_{AB|C_{1}...C_{n-2}}) \ge |\sum_{i=1}^{n-2} \mathcal{N}_{a}^{2}(\rho_{AC_{i}}) - \sum_{i=1}^{n-2} \mathcal{N}_{a}^{2}(\rho_{BC_{i}})|, \tag{40}$$

where  $\rho_{AB}=\mathrm{Tr}_{C_1\dots C_{n-2}}(|\psi\rangle\langle\psi|)$ ,  $\rho_{AC_i}=\mathrm{Tr}_{BC_1\dots C_{i-1}C_{i+1}\dots C_{n-2}}(|\psi\rangle\langle\psi|)$  and  $\rho_{BC_i}=\mathrm{Tr}_{AC_1\dots C_{i-1}C_{i+1}\dots C_{n-2}}(|\psi\rangle\langle\psi|)$ . Proof. We have the following property for linear entropy<sup>35</sup>:

$$T(\rho_{AB}) \ge |T(\rho_A) - T(\rho_B)|. \tag{41}$$

Combining Eq. (24) with Eq. (41), we have

$$2[1 - \text{Tr}(\rho_{AB}^2)] \ge |2[1 - \text{Tr}(\rho_A^2)] - 2[1 - \text{Tr}(\rho_B^2)]|$$
(42)

and

$$C^{2}(|\psi\rangle_{AB|C_{1}...C_{n-2}}) \ge |C^{2}(|\psi\rangle_{A|BC_{1}...C_{n-2}}) - C^{2}(|\psi\rangle_{B|AC_{1}...C_{n-2}})|.$$
(43)

By using the equivalent relation between concurrence and CREN (see Eq. (17)), we have

$$\mathcal{N}^{2}(|\psi\rangle_{AB|C_{1}\cdots C_{n-2}}) \geq |\mathcal{N}^{2}(|\psi\rangle_{A|BC_{1}\cdots C_{n-2}}) - \mathcal{N}^{2}(|\psi\rangle_{B|AC_{1}\cdots C_{n-2}})|. \tag{44}$$

There is a relationship between CREN and CRENOA (see Eq. (21)):

$$\mathcal{N}^{2}(|\psi\rangle_{A|BC_{1}...C_{n-2}}) = \mathcal{N}^{2}_{a}(\rho_{AB}) + \mathcal{N}^{2}_{c}(\rho_{A|C_{1}...C_{n-2}})$$
(45)

$$\mathcal{N}^{2}(|\psi\rangle_{B|AC_{1}...C_{n-2}}) = \mathcal{N}^{2}_{a}(\rho_{BA}) + \mathcal{N}^{2}_{c}(\rho_{B|C_{1}...C_{n-2}}). \tag{46}$$

Putting the above two equalities into Eq. (44), we get

$$|\mathcal{N}_{c}^{2}(\rho_{A|C_{1}...C_{n-2}}) - \mathcal{N}_{c}^{2}(\rho_{B|C_{1}...C_{n-2}})| \geq |\sum_{i=1}^{n-2} \mathcal{N}_{c}^{2}(\rho_{AC_{i}}) - \sum_{i=1}^{n-2} \mathcal{N}_{a}^{2}(\rho_{BC_{i}})|$$

$$\geq |\sum_{i=1}^{n-2} \mathcal{N}_{a}^{2}(\rho_{AC_{i}}) - \sum_{i=1}^{n-2} \mathcal{N}_{a}^{2}(\rho_{BC_{i}})|.$$
(47)

Similar to the above derivation, we give a lower bound about  $\mathcal{N}^2(\ket{\psi}_{AB|C_1...C_{n-2}})$ , i.e.,

$$\mathcal{N}^{2}(|\psi\rangle_{AB|C_{1}...C_{n-2}}) \ge |\sum_{i=1}^{n-2} \mathcal{N}_{a}^{2}(\rho_{AC_{i}}) - \sum_{i=1}^{n-2} \mathcal{N}_{a}^{2}(\rho_{BC_{i}})|.$$
(48)

This lower bound is a direct consequence of CREN.

Theorem 3 shows that the entanglement between AB and the other qubits cannot be less than the absolute value of the difference between both the individual entanglements between A and each of the n-1 remaining qubits and the individual entanglements between B and each of the n-1 remaining qubits. Theorem 3 provides a monogamy-type lower bound of multi-qubit entanglement between the two-qubit system AB and the other (n-2)-qubit system  $C_1C_2...C_{n-2}$  in terms of the squared CRENOA. When  $|\psi\rangle_{AB|C_1...C_{n-2}} = |\psi\rangle_B \otimes |\psi\rangle_{A|C_1...C_{n-2}}$ ,  $\mathcal{N}_a^2(\rho_{BC}) = 0$ , and so we obtain the CWK-type relation in Eq. (18).

Finally, we consider the following four-qubit state  $|\psi\rangle_{ABCD} = a|1000\rangle + b|0010\rangle + c|0001\rangle$  where  $|a|^2 + |b|^2 + |c|^2 = 1$ , from which we can easily obtain the following equations:  $\mathcal{N}_a^2(\rho_{BC}) = \mathcal{N}_a^2(\rho_{BD}) = 0$  and  $\mathcal{N}_a^2(\rho_{AC}) + \mathcal{N}_a^2(\rho_{AD}) = \mathcal{N}^2(|\psi\rangle_{AB|CD}) = \frac{16}{9}$ . Therefore, the state  $|\psi\rangle_{ABCD}$  saturates the monogamy inequality in Eq. (40). Therefore, a generalised monogamy inequality using negativity and CRENOA in an n-qubit is proposed. These relations also give rise to a type of trade-off in inequalities that is related to the upper and lower bounds of CRENOA.

Remark. It is interesting to note that the properties of CREN are based on the subadditivity of linear entropy. However, negativity violates this subadditivity in general conditions<sup>37–39</sup>.

**Examples.** In this section, we use some special states to study generalised monogamy inequalities. First, we consider the (Greenberger-Horne-Zeilinger) GHZ state and W state in Examples 1 and 2. Second, we consider two states in the higher-dimensional system in Examples 3 and 4. Example 1. For an arbitrary pure GHZ state in an *n*-qubit system:

$$|GHZ\rangle = a|0\rangle^{\otimes n} + b|1\rangle^{\otimes n},\tag{49}$$

where  $|a|^2+|b|^2=1$ . The generalized GHZ state is satisfied with the previous CKW inequality. We will now show that the generalised GHZ state satisfies the generalised monogamy inequalities. We have  $\rho_1=\rho_2=\ldots=\rho_n=a^2|0\rangle$   $\langle 0|+b^2|1\rangle\langle 1|$ . It is straightforward to check:  $\mathcal{N}_a^2(\rho_{A_1|A_2\ldots A_n})=\mathcal{N}_a^2(\rho_{A_2|A_1\ldots A_n})=\ldots=\mathcal{N}_a^2(\rho_{A_1|A_1\ldots A_{i-1}A_{i+1}\ldots A_n})=4|(ab)^2|$  and  $\mathcal{N}_a^2(\rho_{A_1A_2})=\mathcal{N}_a^2(\rho_{A_2A_3})=\ldots=\mathcal{N}_a^2(\rho_{A_iA_{i+1}})=4|ab|, \mathcal{N}_a^2(\rho_{A_1A_2|A_3\ldots A_n})=4|(ab)^2|$ . Therefore:

$$\mathcal{N}_{a}^{2}(\rho_{A_{1}|A_{2}\dots A_{n}}) = 4|(ab)^{2}| \le 4(n-1)|(ab)^{2}| = \sum_{i=2}^{n} \mathcal{N}_{a}^{2}(\rho_{A_{i}|A_{1}\dots A_{i-1}A_{i+1}\dots A_{n}}), \tag{50}$$

$$2\mathcal{N}_{a}^{2}(\rho_{AB}) + \sum_{i=1}^{n-2} \mathcal{N}_{a}^{2}(\rho_{AC_{i}}) + \sum_{i=1}^{n-2} \mathcal{N}_{a}^{2}(\rho_{BC_{i}}) = 8(n-1)|(ab)^{2}| \ge 4|(ab)^{2}|$$

$$= \mathcal{N}^{2}(|\psi\rangle_{AB|C_{1}...C_{n-2}}), \tag{51}$$

$$\mathcal{N}^{2}(|\psi\rangle_{AB|C_{1}...C_{n-2}}) = 4|(ab)^{2}| \ge 0 = |\sum_{i=1}^{n-2} \mathcal{N}_{a}^{2}(\rho_{AC_{i}}) - \sum_{i=1}^{n-2} \mathcal{N}_{a}^{2}(\rho_{BC_{i}})|.$$
(52)

Example 2. For a pure state  $|W\rangle$  in an *n*-qubit system:

$$|W\rangle = \frac{1}{\sqrt{n}}(|10\dots 0\rangle + |01\dots 0\rangle\dots + |00\dots 1\rangle),\tag{53}$$

with  $\sum_{i=1}^{n}\left|\frac{1}{\sqrt{n}}\right|^2=1$ . It is very important to understand the saturation of the previous CKW inequality. Using a simple calculation, we have  $\rho_1=\rho_2=\ldots=\rho_n=\frac{1}{n}(|1\rangle\langle 1|)+\frac{n-1}{n}(|0\rangle\langle 0|)$ . It is straightforward to check:

 $\mathcal{N}_{a}^{2}(\rho_{A_{1}|A_{2}\dots A_{n}}) = \mathcal{N}_{a}^{2}(\rho_{A_{2}|A_{1}\dots A_{n}}) = \dots = \mathcal{N}_{a}^{2}(\rho_{A_{1}|A_{1}\dots A_{i-1}A_{i+1}\dots A_{n}}) = \frac{4(n-1)}{n^{2}}.$   $\mathcal{N}_{a}^{2}(\rho_{A_{1}A_{2}}) = \mathcal{N}_{a}^{2}(\rho_{A_{2}A_{3}}) = \dots = \mathcal{N}_{a}^{2}(\rho_{A_{i}A_{i+1}}) = \frac{4}{n^{2}}, \\ \mathcal{N}^{2}(\rho_{A_{1}A_{2}|A_{3}\dots A_{n}}) = \frac{8(n-2)}{n^{2}}.$  In the same way, we get the following inequalities:

$$\mathcal{N}_{a}^{2}(\rho_{A_{1}|A_{2}...A_{n}}) = \frac{4(n-1)}{n^{2}} \le \frac{4(n-1)^{2}}{n^{2}} = \sum_{i=2}^{n} \mathcal{N}_{a}^{2}(\rho_{A_{i}|A_{1}...A_{i-1}A_{i+1}...A_{n}}), \tag{54}$$

$$2\mathcal{N}_{a}^{2}(\rho_{AB}) + \sum_{i=1}^{n-2} \mathcal{N}_{a}^{2}(\rho_{AC_{i}}) + \sum_{i=1}^{n-2} \mathcal{N}_{a}^{2}(\rho_{BC_{i}}) = \frac{8(n-1)}{n^{2}} \ge \frac{8(n-2)}{n^{2}} = \mathcal{N}^{2}(|\psi\rangle_{AB|C_{1}...C_{n-2}}), \tag{55}$$

$$\mathcal{N}^{2}(|\psi\rangle_{AB|C_{1}\dots C_{n-2}}) = \frac{8(n-2)}{n^{2}} \ge 0 = |\sum_{i=1}^{n-2} \mathcal{N}_{a}^{2}(\rho_{AC_{i}}) - \sum_{i=1}^{n-2} \mathcal{N}_{a}^{2}(\rho_{BC_{i}})|.$$
(56)

From the above results, we discover that the generalised GHZ state and W state satisfy our inequalities. We further explore the condition of the generalised inequalities in higher-dimensional systems. We consider the following examples:

Example 3. For a pure, totally antisymmetric state  $|\psi_{ABC}\rangle$  in a  $3 \otimes 3 \otimes 3$  system<sup>40</sup>:

$$|\psi_{ABC}\rangle = \frac{1}{\sqrt{6}}(|123\rangle - |132\rangle + |231\rangle - |213\rangle + |312\rangle - |321\rangle).$$
 (57)

This special quantum state is not satisfied with the previous CKW inequality<sup>41</sup> but it is established in generalised monogamy inequalities. We can easily obtain  $\mathcal{N}_a^2(\rho_{A|BC}) = \mathcal{N}_a^2(\rho_{B|AC}) = \mathcal{N}_a^2(\rho_{C|AB}) = 4$  and further obtain the inequalities  $\mathcal{N}_a^2(\rho_{A|BC}) \leq \mathcal{N}_a^2(\rho_{B|AC}) + \mathcal{N}_a^2(\rho_{C|AB})$ . We now explore theorems 2 and 3. First, we have  $\mathcal{N}_a^2(\rho_{AB}) = 1, \mathcal{N}_a^2(\rho_{AC}) = 1, \mathcal{N}_a^2(\rho_{BC}) = 1$  and  $\mathcal{N}^2(|\psi\rangle_{AB|C}) = 4$ . Therefore, we obtain the following inequalities:

$$2\mathcal{N}_a^2(\rho_{AB}) + \mathcal{N}_a^2(\rho_{AC}) + \mathcal{N}_a^2(\rho_{BC}) \ge \mathcal{N}^2(|\psi\rangle_{AB|C}) \ge |\mathcal{N}_a^2(\rho_{AC}) - \mathcal{N}_a^2(\rho_{BC})|. \tag{58}$$

Example 4. The *n*-qudit generalised W-class state in higher-dimensional quantum systems is very useful in quantum information theory<sup>42</sup>. We verify whether the generalised monogamy inequalities hold in higher-dimensional systems using a special example. First, we recall the definition of n-qudit generalised W-class state<sup>43</sup>,

$$|W_n^d\rangle_{A_1...A_n} = \sum_{i=1}^{d-1} (a_{1i}|i0...0\rangle + a_{2i}|0i...0\rangle + ... + a_{ni}|00...i\rangle),$$
(59)

where  $\sum_{s=1}^{n}\sum_{i=1}^{d-1}|a_{si}|^{2}=1$ . Let  $|\psi\rangle_{A_{1}...A_{n}}$  be an n-qudit pure state in a superposition of an n-qudit generalised W-class state and vacuum;

$$|\psi\rangle_{A_1\dots A_n} = \sqrt{p} |W_n^d\rangle_{A_1\dots A_n} + \sqrt{1-p} |0\dots 0\rangle_{A_1\dots A_n},\tag{60}$$

for some  $0 \le p \le 1$ .

For the squared negativity  $\mathcal{N}^2$  of  $|\psi\rangle_{A_1...A_n}$  with respect to the bipartition between  $A_1$  and the other qudits, the reduced density matrix  $\rho_{A_1}$  of  $|\psi\rangle_{A_1...A_n}$  onto subsystem  $A_1$  is obtained as

$$\rho_{A_{1}} = \operatorname{Tr}_{A_{2}...A_{n}} |\psi\rangle_{A_{1}A_{2}...A_{n}} \langle \psi | 
= p \sum_{i,j=1}^{d-1} a_{1i} a_{ij}^{*} |i\rangle_{A_{1}} \langle j| + [p\Omega + (1-p)] |0\rangle_{A_{1}} \langle 0| + \sqrt{p(1-p)} 
\times \left[ \sum_{i=1}^{d-1} a_{1i} |i\rangle_{A_{1}} \langle 0| + \sum_{j=1}^{d-1} a_{1j}^{*} |0\rangle_{A_{1}} \langle j| \right],$$
(61)

where  $\Omega = \sum_{s=2}^n \sum_{i=1}^{d-1} |a_{si}|^2 = 1 - \sum_{j=1}^{d-1} |a_{1j}|^2$ . When considering the  $|\psi\rangle_{A_1|A_2...A_n}$  state, we need to obtain the eigenvalue of the matrix by applying the definition of pure state negativity in Eq. (5). Using a simple calculation, we find that the matrix has rank-2 and we have

$$\mathcal{N}^2\Big(|\psi\rangle_{A_1|A_2...A_n}\Big) = \left[\left(\operatorname{Tr}\sqrt{\rho_{A_1}}\right)^2 - 1\right]^2 = 4\lambda_i\lambda_j = 4p^2(1-\Omega)\Omega.$$
(62)

We now consider the case in which n = 2. The remaining cases follow analogously. The two-qudit reduced density matrix  $\rho_{A_1A_2}$  of  $|\psi\rangle_{A_1A_2...A_n}$  is obtained as

$$\rho_{A_{1}A_{2}} = \operatorname{Tr}_{A_{3}...A_{n}} |\psi\rangle_{A_{1}A_{2}...A_{n}} \langle \psi | 
= p \sum_{i,j=1}^{d-1} \left[ a_{1i} a_{1j}^{*} |i0\rangle_{A_{1}A_{2}} \langle j0|a_{1i} a_{2j}^{*} |i0\rangle_{A_{1}A_{2}} \langle 0j| + a_{2i} a_{1j}^{*} |0i\rangle_{A_{1}A_{2}} \langle j0| 
+ a_{2i} a_{2j}^{*} |0i\rangle_{A_{1}A_{2}} \langle 0j| \right] + (p\Omega_{2} + 1 - p) |00\rangle_{A_{1}A_{2}} \langle 00| + \sqrt{p(1-p)} 
\times \sum_{k=1}^{d-1} \left[ (a_{1k}|k0\rangle + a_{2k}|0k\rangle)_{A_{1}A_{2}} \langle 00| + |00\rangle_{A_{1}A_{2}} (a_{1k}^{*} \langle k0| + a_{2k}^{*} \langle 0k|) \right],$$
(63)

where  $\Omega_2 = 1 - \sum_{i=1}^{d-1} (a_{1i}^2 + a_{2i}^2)$ . For convenient calculation, we consider two unnormalised states:

$$|\widetilde{x}\rangle = \sqrt{p} \sum_{i=1}^{d-1} \left( a_{1i} |i0\rangle_{A_1 A_2} + a_{2i} |0i\rangle_{A_1 A_2} \right) + \sqrt{1-p} |00\rangle_{A_1 A_2}, |\widetilde{y}\rangle = \sqrt{\Omega_2} |00\rangle_{A_1 A_2}.$$
(64)

Consequently,  $\rho_{A_1A_2}$  can be represented as  $\rho_{A_1A_2}=|\widetilde{x}\rangle_{A_1A_2}\langle\widetilde{x}|+|\widetilde{y}\rangle_{A_1A_2}\langle\widetilde{y}|$ , where  $|\widetilde{x}\rangle$  and  $|\widetilde{y}\rangle$  are unnormalised states of the subsystems  $A_1A_2$ . By the HJW theorem<sup>44</sup>, any pure-state decomposition  $\rho_{A_1A_2}=\sum_h^r|\widetilde{\psi}_h\rangle_{A_1A_2}\langle\widetilde{\psi}_h|$ with size r > 2 can be obtained by an  $r \times r$  unitary matrix  $u_{hl}$  such that

$$\left|\widetilde{\psi}_{h}\right\rangle_{A_{1}A_{2}} = u_{hl}\left|\widetilde{x}\right\rangle_{A_{1}A_{2}} + u_{h2}\left|\widetilde{y}\right\rangle_{A_{1}A_{2}} \tag{65}$$

for each h, for the normalized state  $|\psi_h\rangle_{A_1A_2}=|\widetilde{\psi}_h\rangle_{A_1A_2}/\sqrt{p_h}$  with  $p_h=|\langle\widetilde{\psi}_h||\widetilde{\psi}_h\rangle|$ . We apply the definition of mixed state negativity in Eqs (8 and 63), and then we have the two-tangle based on

the CREN of  $\rho_{A_1A_2}$  as

$$\mathcal{N}_{c}^{2}(\rho_{A_{1}A_{2}}) = \min \sum_{i} p_{i} \mathcal{N}^{2}(|\psi_{i}\rangle_{A_{1}A_{2}}) = 4p^{2}(1-\Omega) \sum_{i=1}^{d-1} |a_{2i}|^{2} = 4p^{2}(1-\Omega)\Omega',$$
(66)

where  $\Omega' = \sum_{i=1}^{d-1} |a_{2i}|^2$ . From the definition of pure state negativity in Eqs (9 and 63), we have

$$\mathcal{N}_{c}^{2}(\rho_{A_{1}A_{2}|A_{3}...A_{n}}) = \min \sum_{i} p_{i} \mathcal{N}^{2}(|\psi_{i}\rangle_{A_{1}A_{2}|A_{3}...A_{n}}) = 4p\Omega_{2}(1 - \Omega_{2}).$$
(67)

We now try to verify the generalised monogamy inequalities of CREN in an n-qudit system. For convenient calculation, we assume that  $\sum_{i=1}^{d-1}a_{1i}^2=a$ ,  $\sum_{i=1}^{d-1}a_{2i}^2=b$ ,  $\sum_{i=1}^{d-1}a_{1i}^4=A$ ,  $\sum_{i=1}^{d-1}a_{2i}^4=B$ . We first consider the generalisation of Theorem 1.

$$\mathcal{N}_{a}^{2}(\rho_{A_{1}|A_{2}...A_{n}}) = \mathcal{N}_{a}^{2}(\rho_{A_{2}|A_{1}...A_{n}}) = ... = \mathcal{N}_{a}^{2}(\rho_{A_{i}|A_{1}...A_{i-1}A_{i+1}...A_{n}})$$

$$= 4p^{2}(1 - \Omega)\Omega = 4p^{2}(1 - a)a. \tag{68}$$

This special quantum state is satisfied with the generalised monogamy inequality in Eq. (25) i.e.,

$$\mathcal{N}_{a}^{2}(\rho_{A_{1}|A_{2}...A_{n}}) \leq \sum_{i=2}^{n} \mathcal{N}_{a}^{2}(\rho_{A_{i}|A_{1}...A_{i-1}A_{i+1}...A_{n}}). \tag{69}$$

For the generalisation of Theorem 2, the left of Eq. (32) is

$$2\mathcal{N}_a^2(\rho_{AB}) + \sum_{i=1}^{n-2} \mathcal{N}_a^2(\rho_{AC_i}) + \sum_{i=1}^{n-2} \mathcal{N}_a^2(\rho_{BC_i}). \tag{70}$$

Using Eqs (8 and 62) we can simplify the calculation to

$$\mathcal{N}_{a}^{2}(\rho_{AB}) + \sum_{i=1}^{n-2} \mathcal{N}_{a}^{2}(\rho_{AC_{i}}) = \mathcal{N}_{c}^{2}(\rho_{A|BC_{1}...C_{n}}) = 4p^{2}(1-\Omega)\Omega = 4p^{2}(1-a)a$$
(71)

and

$$\sum_{i=1}^{n-2} \mathcal{N}_a^2(\rho_{BC_i}) = \mathcal{N}_c^2(\rho_{B|C_1\dots C_n}) = 4p^2(1-\Omega')\Omega' = 4p^2(1-b)b.$$
(72)

After some calculations, we have

$$2\mathcal{N}_{a}^{2}(\rho_{AB}) + \sum_{i=1}^{n-2} \mathcal{N}_{a}^{2}(\rho_{AC_{i}}) + \sum_{i=1}^{n-2} \mathcal{N}_{a}^{2}(\rho_{BC_{i}}) = \mathcal{N}_{a}^{2}(\rho_{AB}) + \sum_{i=1}^{n-2} \mathcal{N}_{a}^{2}(\rho_{AC_{i}})$$

$$+ \sum_{i=1}^{n-2} \mathcal{N}_{a}^{2}(\rho_{BC_{i}}) + \mathcal{N}_{a}^{2}(\rho_{AB}) = \mathcal{N}_{c}^{2}(\rho_{A|BC_{1}...C_{n}}) + \mathcal{N}_{c}^{2}(\rho_{B|C_{1}...C_{n}}) + \mathcal{N}_{a}^{2}(\rho_{AB})$$

$$= 4p^{2}(1-a)a + 4p^{2}(1-b)b + 4p^{2}ab.$$

$$(73)$$

Second, taking Eq. (67) to the right side of Eq. (32), we then have

$$\mathcal{N}^{2}(|\psi\rangle_{AB|C_{1}...C_{n-2}}) = 4p\Omega_{2}(1 - \Omega_{2}) = 4p^{2}[1 - (a+b)](a+b). \tag{74}$$

After a straightforward calculation, we obtain

$$2\mathcal{N}_{a}^{2}(\rho_{AB}) + \sum_{i=1}^{n-2} \mathcal{N}_{a}^{2}(\rho_{AC_{i}}) + \sum_{i=1}^{n-2} \mathcal{N}_{a}^{2}(\rho_{BC_{i}}) - \mathcal{N}^{2}(|\psi\rangle_{AB|C_{1}...C_{n-2}}) = 12p^{2}ab \ge 0.$$
(75)

Therefore, this *n*-qudit pure state is satisfied with the generalised monogamy inequality in Eq. (32). In other words, the test of the Theorem 2 has been accomplished. Next, we verify Theorem 3. First, we consider the term CREN from Eq. (40):

$$\sum_{i=1}^{n-2} \mathcal{N}_a^2(\rho_{AC_i}) = \mathcal{N}_c^2(\rho_{A|BC_1...C_n}) - \mathcal{N}_a^2(\rho_{AB}) = 4p^2(1-\Omega)\Omega - 4p^2(1-\Omega)\Omega'$$

$$= 4p^2a(1-a-b). \tag{76}$$

Calculating the absolute value of the difference between Eqs (72 and 76), we obtain

$$\left| \sum_{i=1}^{n-2} \mathcal{N}_a^2 \left( \rho_{AC_i} \right) - \sum_{i=1}^{n-2} \mathcal{N}_a^2 \left( \rho_{BC_i} \right) \right| = |4p^2 (a - a^2 - ab + b^2 - b)| = 4p^2 (a - a^2 - ab + b^2 - b). \tag{77}$$

It is easy to check  $4p^2(a - a^2 - ab + b^2 - b) > 0$ , as

$$0 \le a + b \le 1 \implies a(a + b) \le a \implies a^2 + ab - b + b^2 < a(a + b) \le a$$
  

$$\implies a^2 + ab - b + b^2 - a < 0 \implies a - a^2 - ab + b^2 - b > 0$$
  

$$\implies 4p^2(a - a^2 - ab + b^2 - b) > 0.$$
(78)

After a straightforward calculation, we have

$$\mathcal{N}^{2}\Big(|\psi\rangle_{AB|C_{1}...C_{n-2}}\Big) - \left|\sum_{i=1}^{n-2} \mathcal{N}_{a}^{2}\Big(\rho_{AC_{i}}\Big) - \sum_{i=1}^{n-2} \mathcal{N}_{a}^{2}\Big(\rho_{BC_{i}}\Big)\right| = 4p^{2}b(2 - 2b - a) \ge 0.$$
(79)

Therefore, this n-qudit pure state satisfies the generalised monogamy inequality in Eq. (40). We have now verified the generalised monogamy inequalities. In other words, the generalised monogamy inequality are satisfied with the n-qudit pure state for all three of our theorems.

# **Conclusions**

In this paper, we have used CREN to study different types of monogamy relations. In particular, we have shown that CREN satisfies the generalised monogamy inequalities. We have investigated the CKW-like inequalities and generalised monogamy inequalities. Furthermore, the generalised monogamy inequalities related to CREN and CRENOA were obtained by n-qubit states. These relations also give rise to a type of trade-off in inequalities that is related to the upper and lower bounds of CRENOA. Finally, we have shown that the partially coherent superposition of the generalised W-class state and vacuum extensions of CREN satisfies the generalised monogamy inequalities. We believe that the generalised monogamy inequalities can be useful in quantum information theory. This paper was based on the linear entropy. To continue this work, we will study the nature of other entropy further in the future work. We hope that our work will be useful to the quantum physics.

# References

- 1. Horodecki, R., Horodecki, P., Horodecki, M. & Horodecki, K. Quantum entanglement. Rev. Mod. Phys. 81, 865 (2009).
- 2. Bennett, C. H. et al. Teleporting an unknown quantum state via dual classical and Einstein-Podolsky-Rosen channels. Phys. Rev. Lett. 70, 1895 (1993).
- 3. Popescu, S., Nonlocality beyond quantum mechanics. Nature Phys. 10, 264 (2014).
- 4. Eisert, J., Cramer, M. & Plenio, M. B. Area laws for the entanglement entropy—a review. Rev. Mod. Phys. 82, 277 (2010).
- 5. Koashi, M. & Winter, A. Monogamy of quantum entanglement and other correlations. Phys. Rev. A 69, 022309 (2004).
- 6. Terhal, B. Is entanglement monogamous? IBM J. Res. Dev. 48, 71 (2004).
- 7. Kay, A., Kaszlikowski, D. & Ramanathan, R. Optimal cloning and singlet monogamy. Phys. Rev. Lett. 103, 050501 (2009).
- 8. Coffman, V., Kundu, J. & Wootters, W. K. Distributed entanglement. Phys. Rev. A 61, 052306 (2000).
- 9. Osborne, T. J. & Verstraete, F. General monogamy inequality for bipartite qubit entanglement. Phys. Rev. Lett. 96, 220503 (2006).
- 10. Bennett, C. H. Quantum cryptography using any two nonorthogonal states. *Phys. Rev. Lett.* **68**, 3121 (1992).
- 11. Barrett, J., Hardy, L. & Kent, A. No signaling and quantum key distribution. Phys. Rev. Lett. 95, 010503 (2005).

- 12. Dowling, M. R., Doherty, A. C. & Wiseman, H. M. Entanglement of indistinguishable particles in condensed-matter physics. *Phys. Rev. A* 73, 052323 (2006).
- 13. Ma, X. S. et al. Quantum simulation of the wavefunction to probe frustrated heisenberg spin systems. Nat. Phys. 7, 399 (2009).
- 14. Kabat, D. Black hole entropy and entropy of entanglement. Nuclear Physics B 453.1 (1995).
- 15. Lloyd, S. & Preskill, J. Unitarity of black hole evaporation in final-state projection models. J. High Energy Phys. 08126 (2014).
- 16. Ou, Y. C. & Fan, H. Monogamy inequality in terms of negativity for three-qubit states. Phys. Rev. A 75, 062308 (2007).
- 17. Kim, J. S., Das, A. & Sanders, B. S. Entanglement monogamy of multipartite higher-dimensional quantum systems using convex-roof extended negativity. *Phys. Rev. A* **79**, 012329 (2009).
- 18. de Oliveira, T. R., Cornelio, M. F. & Fanchini, F. F. Monogamy of entanglement of formation. Phys. Rev. A 89, 034304 (2014).
- 19. Bai, Y.-K., Xu, Y.-F. & Wang, Z. D. General monogamy relation for the entanglement of formation in multiqubit systems. *Phys. Rev.* Lett.113, 100503 (2014).
- 20. Lancien, C. et al. Should entanglement measures be monogamous or faithful? arXiv:1604.02189 (2016).
- 21. Song, W. et al. General monogamy relation of multiqubit systems in terms of squared Rényi- $\alpha$  entanglement. Phys. Rev. A 93, 022306 (2016).
- 22. Luo, Y., Tian, T., Shao, L.-H. & Li, Y.-M. General monogamy of Tsallis *q*-entropy entanglement in multiqubit systems. *Phys. Rev. A* 93, 062340 (2016).
- 23. Yuan, G. M. et al. Monogamy relation of multi-qubit systems for squared Tsallis-q entanglement. Sci. Rep. 6, 28719 (2016).
- 24. Bai, Y.-K., Zhang, N., Ye, M.-Y. & Wang, Z. D. Exploring multipartite quantum correlations with the square of quantum discord. *Phys. Rev. A* 88, 012123 (2013).
- 25. Regula, B., Martino, S. D., Lee, S. & Adesso, G. Strong monogamy conjecture for multiqubit entanglement: The four-qubit case. *Phys, Rev. Lett.* 113, 110501 (2014).
- 26. Zhu, X.-N. & Fei, S.-M. Generalized monogamy relations of concurrence for N-qubit systems. Phys. Rev. A 92, 062345 (2015).
- 27. Wootters, W. K. Entanglement of formation of an arbitrary state of two qubits. Phys. Rev. Lett. 80, 2245 (1998).
- 28. Laustsen, T., Verstraete, F. & Van enk, S. J. Local versus joint measurements for the entanglement of assistance. *Quantum Inf. Comput.* 3, 64 (2003).
- 29. Vidal, G. & Werner, R. F. Computable measure of entanglement. Phys. Rev. A 65, 032314 (2002).
- 30. Horodecki, M., Horodecki, P. & Horodecki, R. Mixed-State entanglement and distillation: Is there a "Bound" entanglement in nature? *Phys. Rev. Lett.* **80**, 5239 (1998).
- 31. Horodeki, P. Separability criterion and inseparable mixed states with positive partial transposition. Phys. Lett. A. 232, 333 (1997).
- 32. Dur, W., Cirac, J. I., Lewenstein, M. & Bru ß, D. Distillability and partial transposition in bipartite systems. Phys. Rev. A 61, 062313 (2000).
- 33. Yu, C.-S. & Song, H.-S. Measurable entanglement for tripartite quantum pure states of qubits. Phys. Rev. A 76, 022324 (2007).
- 34. Santos, E. & Ferrero, M. Linear entropy and Bell inequalities. Phys. Rev. A 62, 024101 (2000).
- 35. Zhang, C.-J., Gong, Y.-X., Zhang, Y.-S. & Guo, G.-C. Observable estimation of entanglement for arbitrary finite-dimensional mixed states. *Phys. Rev. A* **78**, 042308 (2008).
- 36. Luo, Y. & Li, Y.-M. Monogamy of  $\alpha$ th power entanglement measurement in qubit systems. *Ann. Phys.* **362**, 511 (2015).
- 37. Rastegin, A. E. Some general properties of unified entropies. J. Stat. Phys. 143, 1120 (2011).
- 38. Rossignoli, R., Canose, N. & Ciliberti, L. Generalized entropic measures of quantum correlations. Phys. Rev. A 82, 052342 (2010).
- 39. Hu, X.-H. & Ye, Z.-X. Generalized quantum entropy. J. Math. Phys. 47, 023502 (2006).
- 40. Ou, Y. C. Violation of monogamy inequality for higher-dimensional objects. Phys. Rev. A 75, 034305 (2007).
- 41. Choi, J. H. & San Kim, J. Negativity and strong monogamy of multiparty quantum entanglement beyond qubits. *Phys. Rev. A* **92**, 042307 (2015).
- 42. Li, L.-Z. & Qiu, D.-W. The states of W -class as shared resources for perfect teleportation and superdense coding. *J. Phys. A.* 40, 10871 (2007).
- 43. Kim, J. S. & Sanders, B. C. Generalized W -class state and monogamy relation. J. Phys. A. 41, 495301 (2008).
- 44. Hughston, L. P., Jozsa, R. & Wootters, W. K. A complete classification of quantum ensembles having a given density matrix. *Phys. Lett. A.* **183**, 14 (1993).

### **Acknowledgements**

It is a pleasure to thank F. G. Zhang for inspiring discussions. We thank the anonymous referees for their valuable comments. This work was supported by the National Nature Science Foundation of China (Grant No. 1127123), the Higher School Doctoral Subject Foundation of Ministry of Education of China (Grant No. 20130202110001) and the Fundamental Research Funds for the Central Universitie (Grant No. 2016CBY003).

## **Author Contributions**

T.T. and Y. Luo contributed the idea. T.T. performed the calculations and wrote the main manuscript. Y. Luo checked the calculations. Y. Li improved the manuscript. All authors contributed to the discussion and reviewed the manuscript.

# **Additional Information**

**Competing financial interests:** The authors declare no competing financial interests.

**How to cite this article**: Tian, T. *et al.* Generalised monogamy relation of convex-roof extended negativity in multi-level systems. *Sci. Rep.* **6**, 36700; doi: 10.1038/srep36700 (2016).

**Publisher's note:** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit http://creativecommons.org/licenses/by/4.0/

© The Author(s) 2016