

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/342958208>

geemap: A Python package for interactive mapping with Google Earth Engine

Article in The Journal of Open Source Software · July 2020

DOI: 10.21105/joss.02305

CITATIONS

9

READS

1,913

1 author:



[Qiusheng Wu](#)

University of Tennessee

65 PUBLICATIONS 1,210 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Localization of geographical analyses [View project](#)



Mapping Vernal Pools in the Northeastern United States [View project](#)

geemap: A Python package for interactive mapping with Google Earth Engine

Qiusheng Wu¹

¹ Department of Geography, University of Tennessee, Knoxville, TN 37996, United States

DOI: [10.21105/joss.02305](https://doi.org/10.21105/joss.02305)

Software

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

Editor: [Hugo Ledoux](#) ↗

Reviewers:

- [@fbiljecki](#)
- [@steflhermitte](#)

Submitted: 22 May 2020

Published: 15 July 2020

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Summary

geemap is a Python package for interactive mapping with [Google Earth Engine](#) (GEE), which is a cloud computing platform with a [multi-petabyte catalog](#) of satellite imagery and geospatial datasets (e.g., Landsat, Sentinel, MODIS, NAIP) (Gorelick et al., 2017). During the past few years, GEE has become very popular in the geospatial community and it has empowered numerous environmental applications at local, regional, and global scales. Some of the notable environmental applications include mapping global forest change (Hansen et al., 2013), global urban change (Liu et al., 2020), global surface water change (Pekel, Cottam, Gorelick, & Belward, 2016), wetland inundation dynamics (Wu et al., 2019), vegetation phenology (Li et al., 2019), and time series analysis (Kennedy et al., 2018).

GEE provides both JavaScript and Python APIs for making computational requests to the Earth Engine servers. Compared with the comprehensive [documentation](#) and interactive IDE (i.e., [GEE JavaScript Code Editor](#)) of the GEE JavaScript API, the GEE Python API lacks good documentation and lacks functionality for visualizing results interactively. The **geemap** Python package is created to fill this gap. It is built upon [ipyleaflet](#) and [ipywidgets](#), enabling GEE users to analyze and visualize Earth Engine datasets interactively with Jupyter notebooks.

geemap Audience

geemap is intended for students and researchers who would like to utilize the Python ecosystem of diverse libraries and tools to explore Google Earth Engine. It is also designed for existing GEE users who would like to transition from the GEE JavaScript API to a Python API. The automated JavaScript-to-Python [conversion module](#) of the **geemap** package can greatly reduce the time needed to convert existing GEE JavaScripts to Python scripts and Jupyter notebooks.

geemap Functionality

The interactive mapping functionality of the **geemap** package is built upon [ipyleaflet](#) and [folium](#), both of which rely on Jupyter notebooks for creating interactive maps. A key difference between [ipyleaflet](#) and [folium](#) is that [ipyleaflet](#) is built upon [ipywidgets](#) and allows bidirectional communication between the frontend and the backend, enabling the use of the map to capture user input, while [folium](#) is meant for displaying static data only (QuantStack, 2019). It should be noted that [Google Colab](#) currently does not support [ipyleaflet](#). Therefore, if one wants to use **geemap** on Google Colab, one should import `geemap.eefolium` as `geemap`, which provides limited interactive mapping functionality. To utilize the full interactive mapping

functionality of **geemap**, one should import `geemap` on a local computer or secured server with Jupyter notebook installed.

The key functionality of **geemap** is organized into several modules:

- [geemap](#): the main module for interactive mapping with Google Earth Engine, `ipyleaflet`, and `ipywidgets`.
- [eefolium](#): a module for interactive mapping with Earth Engine and `folium`. It is designed for users to run `geemap` with Google Colab.
- [conversion](#): utilities for automatically converting Earth Engine JavaScripts to Python scripts and Jupyter notebooks.
- [basemaps](#): a module for adding various XYZ and WMS tiled basemaps.
- [legends](#): a module for adding customized legends to interactive maps.

geemap Tutorials

Various tutorials and documentation are available for using **geemap**, including:

- [20+ video tutorials with corresponding notebook examples](#)
- [360+ Jupyter notebook examples for using Google Earth Engine](#)
- [Complete documentation on geemap modules and methods](#)

Acknowledgements

The author would like to thank the developers of `ipyleaflet` and `ipywidgets`, which empower the interactive mapping functionality of **geemap**, including [Martin Renou](#), [Sylvain Corlay](#), and [David Brochart](#). The author would also like to acknowledge source code contributions from [Justin Braaten](#), [Cesar Aybar](#), [Oliver Burdekin](#), [Diego Garcia Diaz](#), and [Stephan Büttig](#).

References

- Gorelick, N., Hancher, M., Dixon, M., Ilyushchenko, S., Thau, D., & Moore, R. (2017). Google Earth Engine: Planetary-scale geospatial analysis for everyone. *Remote Sensing of Environment*, 202, 18–27. doi:[10.1016/j.rse.2017.06.031](#)
- Hansen, M. C., Potapov, P. V., Moore, R., Hancher, M., Turubanova, S. A., Tyukavina, A., Thau, D., et al. (2013). High-resolution global maps of 21st-century forest cover change. *Science*, 342(6160), 850–853. doi:[10.1126/science.1244693](#)
- Kennedy, R. E., Yang, Z., Gorelick, N., Braaten, J., Cavalcante, L., Cohen, W. B., & Healey, S. (2018). Implementation of the LandTrendr Algorithm on Google Earth Engine. *Remote Sensing*, 10(5), 691. doi:[10.3390/rs10050691](#)
- Li, X., Zhou, Y., Meng, L., Asrar, G. R., Lu, C., & Wu, Q. (2019). A dataset of 30 m annual vegetation phenology indicators (1985–2015) in urban areas of the conterminous United States. *Earth System Science Data*. doi:[10.5194/essd-11-881-2019](#)
- Liu, X., Huang, Y., Xu, X., Li, X., Li, X., Ciais, P., Lin, P., et al. (2020). High-spatiotemporal-resolution mapping of global urban change from 1985 to 2015. *Nature Sustainability*, 1–7. doi:[10.1038/s41893-020-0521-x](#)

- Pekel, J.-F., Cottam, A., Gorelick, N., & Belward, A. S. (2016). High-resolution mapping of global surface water and its long-term changes. *Nature*, 540(7633), 418–422. doi:[10.1038/nature20584](https://doi.org/10.1038/nature20584)
- QuantStack. (2019). Interactive GIS in Jupyter with ipyleaflet - Jupyter Blog. Retrieved from <https://blog.jupyter.org/interactive-gis-in-jupyter-with-ipyleaflet-52f9657fa7a>
- Wu, Q., Lane, C. R., Li, X., Zhao, K., Zhou, Y., Clinton, N., DeVries, B., et al. (2019). Integrating LiDAR data and multi-temporal aerial imagery to map wetland inundation dynamics using Google Earth Engine. *Remote Sensing of Environment*, 228, 1–13. doi:[10.1016/j.rse.2019.04.015](https://doi.org/10.1016/j.rse.2019.04.015)