

## State Categories and Response Functors

---

*1st draft*

F. William Lawvere\*

*Dedicated to Walter Noll*

The concept of state, which was already included among the Aristotelian categories, is logically presupposed by all science, yet becomes clear only after extended development of several particular sciences.

In recent decades, important advances in the understanding of the concept of state have been made by Walter Noll [1] and others [2], yet further simplification will allow still wider use of these advances. With the help of the modern theory of categories, due to Eilenberg and Mac Lane [3], many parts of geometry and analysis have been simplified through unification. The following considerations are offered in the conviction that category theory can also assist in unifying the diverse concepts of the foundation of continuum physics with each other and with other parts of mathematics.

Plato's observation (incorrectly attributed to Heracleitus) that "all is flux" and "nothing is", does not lead to exact science, for exact science requires sober consideration of precise, distinguishable states. On the

\* supported by NSF


other hand, states are just the way-stations in definite processes of change, so that neither states nor processes can do without the other. Further, among processes there are special ones that are not processes at all. These observations are descriptive of the relationship of objects, morphisms, and identity morphisms in a mathematical category. Thus the theory of mathematical categories provides precise terminology and constructions for a significant part of the theory of states. For example, the notion of a cyclic process, beginning and ending at a given state, may be identified with that of endomorphism of a given object. But what particular structure and properties will categories arising in this manner from continuum physics enjoy? Can we hope to characterize such categories mathematically? It appears that they have very strong properties, some of which I shall try to explain in this paper.

Perhaps one day a (useful and not merely speculative) analysis of the concept of time will arise from a deeper study of state categories themselves, but (at least in the continuous case) we must here consider duration -as an additional structure on a category.


Definition

If  $x$  is a morphism in a category  $X$  then the interval  $I(x)$  is the category constructed as follows:

An object of  $\underline{I}(x)$  is any ordered pair  $(\alpha_0, \alpha_1)$  of morphisms of  $\underline{x}$  for which  $x = \alpha_1 \alpha_0$  and a morphism  $(\alpha_0, \alpha_1) \xrightarrow{a} (\beta_0, \beta_1)$  in  $\underline{I}(x)$  is any morphism of  $\underline{x}$  such that  $\beta_0 = a\alpha_0$  and  $\beta_1 a = \alpha_1$



(of course  $\beta_1 a \alpha_0 = x$  for any morphism  $a \xrightarrow{a} \beta$  in  $\underline{I}(x)$ . If  $\beta \xrightarrow{b} \gamma$  is another morphism in  $\underline{I}(x)$ , then  $a \xrightarrow{ba} \gamma$  is easily verified to be another morphism in  $\underline{I}(x)$ )



thus defining the composition operation needed to make  $\underline{I}(x)$  a category.

#### Proposition

If  $x_0 \xrightarrow{x} x_1$  is a given morphism in a category  $\underline{x}$ , then there is a canonical "forgetful" functor  $\underline{I}(x) \longrightarrow \underline{x}$  which to any  $x_0 \xrightarrow{\alpha_0} A \xrightarrow{\alpha_1} x_1$  with  $\alpha_0 \alpha_1 = x$  assigns the object  $A$  in  $\underline{x}$ . The category  $\underline{I}(x)$  always has an initial object  $0_x = (1_{x_0}, x)$  and a terminal object  $1_x = (x, 1_{x_1})$

which map to  $x_0, x_1$  respectively via the canonical functor  $\underline{I}(x) \longrightarrow \underline{X}$ , while the unique morphism  $o_x \longrightarrow 1_x$  in  $\underline{I}(x)$  maps to  $x$ .

We may sometimes write  $A \in \underline{X}$  to indicate that  $A$  is the image of the canonical functor  $\underline{I}(x) \longrightarrow \underline{X}$ .

#### Definition

A functor  $\underline{X} \xrightarrow{d} \underline{T}$  will be called a duration if for every morphism  $x$  in  $\underline{X}$  the induced functor  $\underline{I}(x) \xrightarrow{d} \underline{I}(dx)$  is an isomorphism of categories. This definition will be used mainly when  $\underline{T}$  is a commutative monoid considered as a category with one object.

#### Remark

Our definition  $\underline{I}(x) \xrightarrow{\sim} \underline{I}(dx)$  of duration functor  $d$  is equivalent to the requirement that  $d$  enjoy "unique lifting of factorization", i.e.  $d(x) = t_1 + t_0 \Rightarrow$  implies there is a unique pair  $x_1, x_0$  for which  $x_1 x_0 = x$ ,  $d(x_1) = t_1, d(x_0) = t_0$ .

#### Proposition

If  $\underline{X} \xrightarrow{d} \underline{T}$  is a duration then each morphism  $x \xrightarrow{x} x'$  in  $\underline{X}$  determines  $t = d(x)$  in  $\underline{T}$  and a function (the succession of states determined by the process  $x$ )

$$|\underline{I}(t)| \xrightarrow{|x|} |\underline{X}|$$

from the interval of length  $t$  in  $\underline{T}$  into the set of objects of  $\underline{X}$ , whose value at the initial object of the interval is  $x$  and whose value at the terminal object of the interval is  $x'$ .

Proof

Compose the inverse of the functor induced by  $d$  with the canonical forgetful functor

$$\underline{I}(t) \xleftarrow{\sim} \underline{I}(x) \longrightarrow \underline{x}.$$

Proposition

If  $\underline{x} \xrightarrow{d} \underline{T}$  is a duration and if  $\underline{T}$  is a category in which every morphism is an epimorphism, then  $\underline{x}$  also has the property that every morphism is an epimorphism.

Proof

Suppose  $x_0 \xrightarrow{\alpha} A \xrightarrow[x'']{x'} x_1$  are such that  $x'\alpha = x''\alpha$ ; we must show  $x' = x''$ . Since  $d(x')d(\alpha) = d(x'')d(\alpha)$  and since  $d(\alpha)$  is an epimorphism, we have  $d(x') = d(x'')$ .

Denoting by  $x = x'\alpha = x''\alpha$  the common value, we see that  $\underline{I}(x) \xrightarrow[d]{\approx} \underline{I}(dx)$  takes the two objects  $(\alpha, x')$ ,  $(\alpha, x'')$  to the same object of  $\underline{I}(dx)$ ; hence these objects are the same, so that in particular  $x' = x''$ .

Proposition

Let  $\underline{T}$  be a monoid (category with single object) whose composition is commutative and written additively.

Suppose  $\underline{T}$  satisfies the two cancellation properties

$$t' + s = t'' + s \implies t' = t''$$

$$t + s = 0 \implies t = 0 \text{ and } s = 0$$

Then for any category  $\underline{X}$  which admits a duration functor  $\underline{X} \xrightarrow{d} \underline{T}$  valued in  $\underline{T}$  (i.e.  $d(\alpha_1 \alpha_0) = d(\alpha_1) + d(\alpha_0)$ ,  $d(l_A) = 0$ ) one has that every morphism  $\alpha$  in  $\underline{X}$  is both an epimorphism and a monomorphism but not an isomorphism nor an idempotent unless  $\alpha = l_A$  is an identity. That is, any  $\alpha$  is both right cancellable and left cancellable, but

$$\begin{aligned}\alpha_1 \alpha_0 &= l_X \implies \alpha_0 = \alpha_1 = l_X \\ \alpha \alpha &= \alpha \implies \alpha = l_A\end{aligned}$$

Proof

The first cancellation law for  $\underline{T}$  says that all morphisms in  $\underline{T}$  are epimorphisms, so by the previous proposition the same holds for  $\underline{X}$ . Since the previous proposition can be dualized and since  $\underline{T}$  is commutative, it also follows that all morphisms in  $\underline{X}$  are monomorphisms. On the other hand if we have even a one-sided inverse (retraction or section) then

$$\alpha_1 \alpha_0 = l_X$$

$$d(\alpha_1) + d(\alpha_0) = 0$$

so that, by the second cancellation property,  $d\alpha_i = 0$  and hence  $\alpha_i = l_X$  since  $d$  is "locally" injective  $\underline{I}(l_X) \xrightarrow{\sim} \underline{I}(dl_X) = \underline{1}$ . Of course, the non-existence of non-identity cyclic processes  $\alpha$  which are idempotent  $\alpha\alpha = \alpha$  already follows from the fact that each  $\alpha$  is an epimorphism.

Remark

The primary example of monoid  $\underline{T}$  for continuum mechanics is the set of non-negative real numbers under the operation of addition, so that the category  $\underline{I}(t)$  becomes isomorphic to the linearly ordered set  $[0, t]$ . For comparison we may also consider the discrete case where  $\underline{T}$  is the additive monoid of non-negative natural numbers; a striking difference between the two is that  $t_1 > 0, t_2 > 0 \implies \underline{I}(t_1) \approx \underline{I}(t_2)$  are isomorphic categories in the continuous case, whereas in the discrete case the size of an interval is intrinsic in the sense that a category can have at most one duration functor to the additive monoid of natural numbers. The properties stated in the above proposition are in the discrete case related to those which led Leroux [4] to a powerful generalization of the Möbius inversion formula.

Example

If  $\underline{T}$  is any category and if  $T_o$  is any object, consider the category  $\underline{T}/T_o$  whose objects are morphisms  $T \longrightarrow T_o$  in  $\underline{T}$  and whose morphisms are commutative triangles of the form

$$\begin{array}{ccc} T' & \xrightarrow{\quad} & T \\ & \searrow & \swarrow \\ & T_o & \end{array}$$

in  $\underline{T}$ . The obvious forgetful functor  $\underline{T}/T_o \longrightarrow \underline{T}$  is always a duration. If  $\underline{T}$  has left cancellation (more

generally, if every morphism to  $T_0$  in  $\underline{T}$  is a monomorphism) then  $\underline{T}/T_0$  has the stronger property of being a poset (that is, a category in which there is at most one morphism  $x \rightarrow x'$  for each given ordered pair of objects). In case  $\underline{T}$  is a monoid (so  $T_0$  is the unique object) then  $\underline{T}/T_0$  is usually referred to as the divisibility of  $\underline{T}$ ; for example if  $\underline{T}$  is additive,  $t' \rightarrow t$  in  $\underline{T}/T_0$  iff  $t' \geq t$ , and the forgetful duration functor

$$\underline{T}/T_0 \longrightarrow T$$


is usually denoted as "difference", which is a functor because  $t'' \geq t' \geq t$  implies

$$(t'' - t') + (t' - t) = t' - t.$$

Incidentally, the category of categories seems the only reasonable context in which fundamental relationships between objects as disparate as posets and monoids become structure-preserving morphisms.

Example

Given a set  $|X|$  and a suitable monoid  $T$ , a path-category can be constructed by considering the elements of the set as objects and functions  $|\underline{I}(t)| \xrightarrow{x} |X|$  as morphisms. Composition is defined by using a push-out property of



$$\text{where } j_0(s_0, s_1) = (s_0, s_1 + t_1)$$

$$j_1(u_0, u_1) = (t_0 + u_0, u_1)$$

The push-out property (i.e. the unique existence of the indicated  $x$ ) follows if  $\underline{T}$  has the property that all its intervals are linearly ordered; for if  $w \notin I(t_0 + t_1)$ , then the alternative  $w \leq t$  or  $t \leq w$  allows the determination of  $x(w)$  using either  $\alpha_0$  or  $\alpha_1$  as appropriate. Note that a subcategory (of a category with duration) which is moreover closed with respect to factorization, is again a category with duration; in practice, examples are such sub-categories, consisting of processes physically possible in certain contexts, of path-categories.


Smoothness properties such as differentiability are not preserved under the above push-out, yet in spite of this severe handicap the convenience of having a category (i.e. closure under composition) has traditionally been considered more compelling, leading precisely to the notion of

"piecewise smooth" paths. The linear ordering appears again below.

Definition

A freeze of an object  $X$  in a category  $\underline{X}$  is an endomorphism  $f$  of  $X$  with the property that for any factorization

$$f = \alpha_1 \alpha_0$$



in  $\underline{X}$  one has  $X' = X$  (i.e. that  $\alpha_0, \alpha_1$  must also be endomorphisms).

Thus intuitively, a freeze is a process (for example a deformation) throughout which the state (or configuration) remains unchanged.

Proposition

If  $\underline{X}$  is a category in which

$$\underline{I}(l_X) \xrightarrow{\sim} \underline{1} \text{ is a one-element category for any object } X$$

and


$\underline{I}(x)$  is a linear order for any endomorphism  $x$ , then the freezes of  $X$  form a submonoid of the endomorphism monoid  $\underline{X}(X, X)$  of all cyclic processes beginning and ending at  $X$ .

Proof

The first assumption is precisely the condition that  $l_X$  has no non-trivial factorizations, so is certainly a freeze.

?  
What, then, is  
the change that takes  
place in the process?


We must then show that the composition of two freezes  $f, g$  is a freeze. Consider any factorization  $gf = \alpha_1 \alpha_0$ .



By the linear ordering, either there is a  $u$  such that  $\alpha_0 = uf$  and  $\alpha_1 u = g$ , or there is a  $v$  such that  $gv = \alpha_1$  and  $f = \alpha_1 \alpha_0$ . In the latter case  $X' = X$  because  $f$  is a freeze and in the former case  $X' = X$  because  $g$  is a freeze. Hence  $gf$  is a freeze.

obvious!

Note that in a monoid every morphism is a freeze, and since in fact the condition that  $f$  be a freeze of  $A$  in any category  $\underline{X}$ , is equivalent to the requirement that the canonical functor from the interval of  $f$  maps entirely into the submonoid of  $\underline{X}$  consisting of cyclic processes.



#### Proposition

If  $\underline{X} \xrightarrow{d} T$  is a duration with values in a monoid, then any section of  $d$  consists entirely of freezes of some object  $A$  of  $\underline{X}$ .

Proof

By a section of a functor  $d$  is meant a functor  $\underline{T} \xrightarrow{\varphi} \underline{X}$  such that  $d\varphi = 1_{\underline{T}}$ . If  $\underline{T}$  is a monoid then  $\varphi(0) = A$  is an object and  $\varphi$  parameterizes a certain submonoid of  $\text{end}_{\underline{X}}(A)$  which is isomorphic to  $\underline{T}$ . Now we use the special property that  $d$  is a duration to show that each  $\varphi(t)$  is actually a freeze. The diagram

$$\begin{array}{ccccc} \underline{I}(\varphi(t)) & \xrightarrow{\sim} & \underline{I}(d\varphi(t)) & = & \underline{I}(t) \longrightarrow \underline{I}(\varphi(t)) \\ \downarrow & & \downarrow & & \downarrow \\ \underline{X} & \xrightarrow{d} & \underline{T} & \xrightarrow{\varphi} & \underline{X} \end{array}$$

Since  $d$  is  
a duration

shows that  $\underline{I}(\varphi(t))$  maps into a monoid in  $\underline{X}$  since the functor goes via a monoid  $\underline{T}$ .

A section  $\varphi$  of a monoid-valued duration need not meet all freezes at the object  $A$  which  $\varphi$  meets, although it must if all freezes at  $A$  commute. Indeed for any duration  $d$  and any two endomorphisms of  $A$ , if  $d(f) = d(g)$  and  $fg = gf$  then  $f = g$ .

For studies as diverse as control theory and the theory of material constitutive relations, it is necessary to consider explicitly the relation between processes of "deformation" which can be in principle freely carried out on "configuration" states and the processes on

internal states which must necessarily accompany the former. This relation we will take to be that of "fibration", to be defined presently. From a certain point of view the processes that it should be possible to "freely" carry out in any situation must include the freezes, although in fact the cost of the supplies necessary to keep a configuration frozen may be substantial.

Definition

A category  $\underline{C}$  equipped with a duration functor  $\underline{C} \xrightarrow{d} \underline{T}$  to non-negative reals will be called a configuration category (and its morphisms referred to as deformation processes)<sup>\*</sup> in case for every object  $C$  of  $\underline{C}$  there is a given section  $\varphi_C$  of  $d$  for which  $\varphi_C(0) = C$ .

Remark

Of course in applications the configurations may include thermodynamical and other components as well as the purely mechanical aspects (such as infinitesimal metric) usually considered as configurations.

The duration may be considered as the supply of time required to carry out a process, in the sense of the general definition which follows:

\* how should one think of the endomorphisms in  $\underline{C}$ ? (in particular, the freezes). Surely they don't deform the configuration  $C$ , because that is constant during the freeze.

### Definition

A functor  $\underline{x} \xrightarrow{Q} \underline{y}$  whose codomain is a commutative monoid will often be called a **supply**, and  $Q(x)$  referred to as the supply of  $Q$  needed for the process  $x$ .

### Proposition

Let  $p: \underline{v} \longrightarrow \underline{w}$  be a given functor between commutative monoids and let  $\underline{x} \xrightarrow{\Delta v} \underline{v}$ ,  $\underline{x} \xrightarrow{w} \underline{w}$  be supplies.

Suppose  $\underline{x}$  has a duration to a commutative monoid  $\underline{t}$  whose intervals are linearly ordered and has  $I(0) = \mathbf{1}$  and that  $\underline{w}$  is equipped with a submonoid  $\underline{w}_+$  of "non-negative" elements. Define

$$\underline{x}_p = \left\{ x \in \underline{x} \mid \text{for all } y, u, v, \text{ if } x = u y v \text{ then } w(y) \leq p(\Delta v)(y) \right\}$$

where  $\leq$  is defined using  $\underline{w}_+$  in the usual manner. Then  $\underline{x}_p$  is a subcategory of  $\underline{x}$  with the same objects and the restriction of  $d$  to  $\underline{x}_p$  is still a duration.


### Proof

If  $x$  is any identity process then any subprocess  $y$


(i.e.  $x = u y v$ ) is actually  $y = x$  since  $I(x) \geq I(0) = \mathbf{1}$ ;

$(w)(id)$  and  $(\Delta v)(id)$  are both zero, so  $x \in \underline{x}_p$ .


If  $x \xrightarrow{x_0} x' \xrightarrow{x_1} x''$  are each in  $\underline{x}_p$  and if  $y$  is any subprocess of  $x_1 x_0$ ,



then by the linear ordering of  $\underline{I}(x)$  (where  $x = x_1 x_0$ ) we can reduce the question whether  $x \in \underline{x}_p$  to three cases: namely (A)  $(v, uy) \leq (x_0, x_1)$  or (B)  $(x_0, x_1) \leq (v, uy)$ ; in the second case (B),  $y$  is a subprocess of  $x_1$ , (hence  $y$  satisfies the inequality) while in the first case (A) there is  $z$  for which



is commutative. Then  $z$  is a subprocess of  $x_0$ , hence in  $\underline{x}_p$ . Thus there is a further alternative (Al):  
( $z, x_1 \leq (y, u)$  or (A2)  $(y, u) \leq (z, x_1)$ ), in the second case (A2) of which  $y$  is a subprocess of  $z$ , hence in  $\underline{x}_p$ , while in the first case (Al) there exists  $w$  for which



is commutative. Then  $w$  is a subprocess of  $x_1$ , hence in  $\underline{x}_p$ , so that  $y$  is a composite of  $w$  and  $z$ , both of which satisfy the inequality concerning the bound  $p$  on the average pressure.

But the calculation following shows that the processes satisfying the inequality form a subcategory:

$$\begin{aligned} W(y_1 y_o) &= Wy_1 + Wy_o \leq p(\Delta V)y_1 + p(\Delta V)y_o \\ &= p((\Delta V)y_1 + (\Delta V)y_o) \\ &= p(\Delta V)(y_1 y_o). \end{aligned}$$

Thus  $y$  satisfies the inequality as was to be shown.

Remark

If we assume  $(\Delta V)(\varphi) = 0$  and  $W(\varphi) \leq 0$  for freezes  $\varphi$ , (i.e. that work must be done on the system to keep it frozen), then  $\underline{X}_p$  will contain all freezes of  $\underline{X}$ . We have described this (oversimplified) example partly to suggest the relativity of connectedness. Here we use the term "connected" in the naïve sense of category theory, so that the set of components of a category  $\underline{X}$  is obtained by applying the left adjoint of the full inclusion of the category of sets (as discrete categories) into the category of categories; thus  $X, X'$  are in the same component of  $\underline{X}$  if there exists a morphism  $X \dashrightarrow X'$  in  $\underline{X}$  or in other words,  $X'$  is "accessible" from  $X$ . Alternatively, instead of taking the symmetric-transitive closure of the accessibility relation, one could instead require  $X \dashrightarrow X'$  leading to smaller "strong" components. The following general remark applies not only to naïve components, but also to various refinements or limiting versions of the notion of accessibility. Although a state category  $\underline{X}_p$  may have several components and thus

= materials?

appear to be a conglomeration of state categories of several distinct materials, increasing the technology to  $p' \geq p$  may lead to qualitative change in the sense that in  $\underline{X}_p$ , it is possible to process one of these materials into another one. We would like to consider the inclusion  $\underline{X}_p \hookrightarrow \underline{X}_{p'}$ , as a morphism in the category of state categories and hence we do not make connectedness or accessibility part of our general axioms. In the situation described the inclusion would induce a surjective map from the set of components of  $\underline{X}_p$  to the set of components of  $\underline{X}_{p'}$ .

Now we temporarily ignore the duration structure to define the central requirement of determinism. Consider a functor  $\underline{X} \xrightarrow{\Pi} \underline{C}$  in which we think of  $\Pi(X) = C$  as signifying that  $C$  is the configuration in which the internal state  $X$  manifests itself and similarly  $\Pi(x) = \mathcal{J}$  signifies that  $\mathcal{J}$  is the deformation process which accompanies the internal-state-process  $x$ . We require that  $\Pi$  is a (discrete-op) fibration in the sense of the following definition:

Definition

A functor  $\underline{X} \xrightarrow{\Pi} \underline{C}$  is a (discrete-op) fibration if whenever  $C \xrightarrow{\mathcal{J}} C'$  in  $\underline{C}$ , then for any  $X$  for which  $\Pi(X) = C$ , there is a unique morphism  $x$  in  $\underline{X}$  for which both

$$\Pi(x) = \mathcal{J}$$

$$\text{domain} = X.$$

Denote by  $\mathcal{J} \cdot X$  the codomain in  $\underline{X}$  of this unique  $x$ .

Proposition

If  $\widehat{\Pi}$  is a (discrete op-) fibration, then  $x_1 \xrightarrow{x} x_2$ ,  
 $\widehat{\Pi} x_1 = \widehat{\Pi} x_2 = C$  and  $\widehat{\Pi}(x) = 1_C$  implies  $x_1 = x_2$  and  
 $x = 1_x$  (i.e. the fibers of  $\widehat{\Pi}$  are discrete as categories).

One has

$$1_C \cdot x = x$$

and whenever  $C \xrightarrow{f_0} C' \xrightarrow{f_1} C''$  and  $\widehat{\Pi}(x) = C$ , one has

$$f_1 \cdot (f_0 \cdot x) = (f_1 f_0) \cdot x$$

in the fiber over  $C''$ . In this way the category of all  
(discrete op-) fibrations with given base category  $\underline{C}$  is  
equivalent to the category

$$\mathcal{S}_{\underline{C}}$$

of all set-valued covariant functors on  $\underline{C}$ .

Proposition

If  $\underline{X} \xrightarrow{\widehat{\Pi}} \underline{C}$  is a (discrete op-) fibration then  $\widehat{\Pi}$  is  
in particular a "duration" ( $\underline{I}(x) \xrightarrow{\sim} \underline{I}(\widehat{\Pi}(x))$  for all  $x$ )  
and hence the composite  $d \widehat{\Pi}$  is a duration for  $\underline{X}$  whenever  
 $\underline{C} \xrightarrow{d} \underline{T}$  is a duration for  $\underline{C}$ .

Remark

We may refer in a general way to functors  $\underline{X} \longrightarrow \underline{Y}$   
as "outputs", (which may include measurements of "inputs")  
reserving the term "supply" to the case where  $\underline{Y}$  is an  
additive monoid and the term "response functor" for a

situation in which both  $\underline{X}$ ,  $\underline{Y}$  are equipped with duration supplies to the same  $\underline{T}$  and the triangle

$$\begin{array}{ccc} \underline{X} & \xrightarrow{\quad} & \underline{Y} \\ & \searrow & \swarrow \\ & \underline{T} & \end{array}$$

commutes.

A supply  $Q$  is said to be independent of path in case there is codiscrete category  $\underline{U}$  (that is, one in which there is a unique morphism  $\underline{U} \longrightarrow \underline{U}'$  for any ordered pair  $\underline{U}, \underline{U}'$  of objects) and some pair  $E, \Delta$  of functors for which  $Q = \Delta E$  in

$$\begin{array}{ccc} \underline{X} & \xrightarrow{Q} & \underline{Q} \\ & \searrow^E & \swarrow^\Delta \\ & \underline{U} & \end{array}$$

It follows that the image of  $Q$  is a subgroup of the monoid  $\underline{Q}$ .

A frequently arising situation is that in which a supply  $Q: \underline{X} \longrightarrow \underline{Q}$  is absolutely continuous relative to another supply  $S: \underline{C} \longrightarrow \underline{A}$  (which is considered also as a supply on  $\underline{X}$  by composing with a given fibration  $\underline{X} \xrightarrow{\uparrow} \underline{C}$ ). This relationship involves the existence of a response

$$\underline{X} \xrightarrow{\theta} \underline{\text{Hom}}(\underline{A}, \underline{Q})$$


valued in a suitable category whose objects are additive maps (i.e. functors) from the additive monoid A to the additive monoid Q, and is usually written

$$Q = \int \theta ds$$

meaning that for each morphism x in X, Q(x) is "approximated" by

$$\sum \theta(x_i) \cdot s(x_i)$$

for finite partitions



( $x = x_n x_{n-1} \dots x_2 x_1$ ,  $x_i \in x_i$ ) which are sufficiently fine in the sense that

$$d(x_i) \leq \delta(x_i) \quad i = 1, \dots, n$$

for a suitable positive function  $\delta$  on  $|I(x)|$ ,  $d$  being the duration functor of X.

A constitutive relation over a configuration category C involves a pair

$$\begin{array}{ccc} \underline{x} & \xrightarrow{\theta} & \underline{y} \\ \widetilde{\pi} \downarrow & & \\ \underline{c} & & \end{array}$$

where  $\hat{\Pi}$  is a (discrete op-) fibration and where  $\theta$  is an output (response or supply). Some aspects  $\eta \theta$  of  $\theta$  may depend only on  $C$  in the sense that there exists  $S$  with  $s \hat{\Pi} = \eta \theta$ . If all of the relevant response  $\theta$  depends only on  $C$ , the state category  $X \xrightarrow{\hat{\Pi}} C$  may as well be reduced to  $C$  and we are dealing with one of the possible "elastic" responses  $C \xrightarrow{\theta} Y$  whose configuration category is  $C$ . In general a survey of all possible constitutive relations over a given  $C$  involves a range of possibilities for both the state category  $X \xrightarrow{\hat{\Pi}} C$  fibered over  $C$  and for the output. Let us consider separately some of the reasonable restrictions on each of these two aspects. The restrictions on constitutive relations which will be treated here, are of a very general logical nature and so provide at best a background for the more profound restrictions on the second aspect in the study of which Walter Noll was one of the pioneers. (I refer here to the derivation of equations of state from dissipation inequalities.)

For any given configuration category  $C$  whose duration is valued in either the natural numbers or the non-negative real numbers, there is a natural restriction of a general nature that can be imposed on state categories over  $C$ .

#### Definition

Let  $sh(C) \hookrightarrow S^C$  be the full subcategory consisting of all objects corresponding to (discrete op-) fibrations  $X \xrightarrow{\hat{\Pi}} C$  which satisfy the following conditions:

Let  $C$  be a given configuration, and suppose that for every deformation  $C \xrightarrow{\alpha} A$  satisfying

$$d(\alpha) > 0$$

we are given a state  $x_\alpha$  of configuration  $A$  in such a way that for any deformations  $A \xrightarrow{\beta} B$ ,  $C \xrightarrow{\alpha} A$ ,  $C \xrightarrow{\beta} B$  for which  $\beta = \beta \alpha$  and  $d(\alpha) > 0$  (hence  $d(\beta) > 0$ ) we have

$$\beta \cdot x_\alpha = x_\beta;$$

then there is a unique state  $x$  of configuration  $C$  such that for all  $C \xrightarrow{\alpha} A$  with  $d(\alpha) > 0$

$$x_\alpha = \alpha \cdot x$$

The above condition may be considered as a kind of "causal completeness": If all the possible "effects"  $\alpha \cdot x$  (and hence  $\theta(\alpha \cdot x)$  for all outputs  $\theta$ ) for all  $\alpha$  with  $d(\alpha) > 0$  are coherently present in the state category  $\underline{X}$ , then the "cause"  $x$  itself should also be present and uniquely determined. It can be shown that the category  $sh(\underline{C})$  is a topos [5], since the causal completeness condition is actually a "sheaf" condition with respect to a particularly simple example of Grothendieck covering which is induced via the duration from the positivity structure on the time translation monoid  $\underline{T}$ . In the autonomous case, where  $\underline{C} \xrightarrow{d} \underline{T}$  is an isomorphism,  $sh(\underline{C}) \cong sh(\underline{T})$  consists, if  $\underline{T}$  is discrete, of invertible "discrete dynamical systems", but if  $\underline{T}$  is continuous, of possibly dissipative but semi-continuous actions of the additive monoid of non-negative reals. In the

general non-autonomous case the duration functor  
 $\underline{C} \longrightarrow \underline{T}$  induces an exact pair of adjoint functors  
(morphism of toposes)  $sh(\underline{C}) \longrightarrow sh(\underline{T})$ . Moreover,  
the categories  $sh(\underline{C})$  of state categories are quite  
special among toposes even for arbitrary configuration  
categories  $\underline{C}$ , as can be deduced from work of Barel,  
Diaconescu, Freyd, Johnstone and Rosenthal (see  
Johnstone [5a]).

Theorem

For any configuration category  $\underline{C}$  over a commutative  
monoid  $\underline{T}$  with cancellation, there is a topological  
space  $\mathfrak{X}(\underline{C})$  and a topological group  $G$  such that there  
are topos morphisms

$$\mathfrak{X}(\underline{C}) \longrightarrow sh(\underline{C}) \longrightarrow \mathcal{B}_G$$

where the first is a surjective local homeomorphism  
and the second (to the Boolean classifying topos of  $G$ )  
is "localic" (meaning roughly that  $sh(\underline{C})$  can be  
reconstructed from an "internal topological space" with  
 $G$ -action). For discrete time  $\underline{T} = \mathbb{N}$ , the space  $\mathfrak{X}(\underline{C})$   
depends only on the graph  $\underline{C}_1 = \{j \in \underline{C} \mid d(j) = 1\}$  and is a  
generalization of Cantor space and Baire space (which  
themselves are in fact examples of  $\mathfrak{X}(\underline{C})$  for particular  
 $\underline{C}$ ). For continuous time, there is a second "skeletal"  
inclusion  $sh(\underline{C}) \xrightarrow{i_\nabla} \mathcal{S}^{\underline{C}}$  left adjoint  $i^*$  to the original  
 $\uparrow$   
to the left adjoint

inclusion  $i_*$ ; hence  $sh(\underline{C})$  is locally connected, with a "components" functor  $sh(\underline{C}) \longrightarrow S$  assigning to each  $\underline{X}$  its set of components.

In fact, the conclusions in the above theorem require only the fact that all deformations are epimorphisms in  $\underline{C}$  (to obtain the local homeomorphic surjection from the space  $\mathfrak{X}(\underline{C})$ ) and that every deformation is a monomorphism (to obtain the localic representation over  $B_G$ ); the existence of the further left adjoint in the continuous case is at bottom due just to the fact that every strictly positive duration can be expressed as the sum of two strictly positive durations. This particular conjunction of properties is only beginning to be studied in topos theory, yet in fact even our very general definition of duration functor and configuration category implies still more stringent requirements on the topos  $sh(\underline{C})$ , for example, those arising from the non-existence of non-trivial isomorphisms among the deformations and from the interaction between the causal completeness (sheaf) condition and the plentitude of freezes. Thus not only the already existing general topos theory, but also an emerging special branch thereof, provides a powerful background for the construction and comparison within  $sh(\underline{C})$  of the various possible internal state categories which present the same configurational face.

P. 43 - 1

A further condition on a state category  $\underline{X} \xrightarrow{\hat{\Pi}} \underline{C}$ , considered in effect by Walter Noll [1] as part of semi-elasticity, is that among all states above a given configuration  $C$ , there is a unique one  $X$  which is relaxed in the sense that

$$f \cdot X = X$$

for every freeze  $f$  at  $C$ . Of course, such a state is in particular one which admits freezes in the sense of the intrinsic structure of the category  $\underline{X}$  with duration. Relative to a given state category over  $\underline{C}$ , one can define an elastic deformation in  $\underline{C}$  to be any deformation  $C \xrightarrow{\alpha} C'$  such that if  $X$  is a relaxed state with  $\hat{\Pi} X = C$ , then  $\alpha \cdot X$  is relaxed. For a fibration  $\hat{\Pi}$  satisfying Noll's condition, these form a subcategory  $\underline{C}_{\text{elast}} \hookrightarrow \underline{C}$  containing all configurations, and there is a functorial section  $\lambda$  of  $\hat{\Pi}$  over it

$$\begin{array}{ccc} & X & \\ \lambda \nearrow & \downarrow \hat{\Pi} & \\ \underline{C}_{\text{elast}} & \xrightarrow{\quad} & \underline{C} \end{array}$$

such that if  $\varPhi$  is any section of  $d$ ,  $\varPhi$  actually has values in  $\underline{C}_{\text{elast}}$ , and  $\lambda \varPhi$  is a section of the duration  $d \hat{\Pi}$  of  $\underline{X}$ .

Now a standard example of a fibration over  $\underline{C}$  is  $\underline{C}(C_0, -)$  for a particular configuration  $C_0$ ; in considering this as a state category, those states which are of configuration  $C$  are the "finite histories"  $C_0 \longrightarrow C$  which began at  $C_0$  and which end at  $C$ ; the fibration property is just the action by composition in  $\underline{C}$ .

(Caution: In the case of discrete time this object is not in the subcategory  $\text{sh}_{\underline{C}} \hookrightarrow \mathcal{S}^{\underline{C}}$  and its sheaf completion must be considered instead.) The object

$$\mathcal{H}_{\text{fin}}(\underline{C}) = \sum_{c_o \in \underline{C}} \underline{C}(c_o, -)$$

has as its states all the finite histories. The famous Yoneda Lemma implies the following:

Proposition

If  $\underline{x}$  is any object of  $\text{sh}(\underline{C})$ , there is a one-to-one correspondence between functors of state categories over  $\underline{C}$

$$\underline{C}(c_o, -) \longrightarrow \underline{x}$$

and choices of a single state of  $\underline{x}$  of configuration  $c_o$ . Similarly, each functor of state categories over  $\underline{C}$

$$\mathcal{H}_{\text{fin}}(\underline{C}) \longrightarrow \underline{x}$$

is uniquely determined by a function  $\lambda$  which assigns to each configuration  $C$  a state  $\lambda(C)$  of underlying configuration  $C$ .

In case  $\underline{x}$  has the (fragment of) semi-elastic structure given by a function  $\lambda$  assigning to every configuration  $C$  the relaxed state  $\lambda(C)$  of configuration  $C$ , we thus obtain a standard map

$$\begin{array}{ccc} \mathcal{H}_{\text{fin}}(\underline{C}) & \xrightarrow{\tilde{\lambda}} & \underline{x} \\ & \searrow & \swarrow \pi \\ & \underline{C} & \end{array}$$


expressing one conception of how what is (German 'Wesen') results from what was (German 'gewesen'). Using topology, limits, and further conditions on X, it may be possible to extend  $\hat{\lambda}$  to some larger subcategory

$$\mathcal{H}_{\text{fin}}(C) \hookrightarrow ? \hookrightarrow \mathcal{H}(C)$$

of the category of infinite histories (which are defined [6] to be sections  $\underline{T}/O \xrightarrow{h} \underline{C}$  of  $d$  over the difference functor  $\underline{T}/O \xrightarrow{\Delta} T$ ); the inclusion of  $\mathcal{H}_{\text{fin}}$  into  $\mathcal{H}$  is defined using continuation into the past by constants. Special interest attaches to those semi-elastic state categories X for which the map  $\hat{\lambda}$  (or such an extension of it) maps surjectively to X, or at least densely in the sense that within a given family of preferred "continuous" outputs, a pair of continuous outputs agree on X provided they agree when composed with  $\hat{\lambda}$ . Such a condition on X seems to imply a drastic reduction of the range of possibilities for state categories compared to the proper class  $\text{sh}(C)$ . However, one must also consider the possibility of enlarging  $C' \longrightarrow C$  (the notion of configuration (and deformation)), for example a purely mechanical C may be enlarged to a thermomechanical C'. Thus in the extreme we could imagine being given only a fibration X  $\longrightarrow T$  over the time monoid alone, an autonomous necessity without any given notion C of deformation process. Deeper knowledge of X may however

be accompanied by the possibility of constructing a category  $\underline{C}$  of freely possible deformations, a fibration  $\underline{x} \xrightarrow{\pi} \underline{C}$  and a functor  $\mathcal{H}_?(\underline{C}) \longrightarrow \underline{x}$  which is surjective or dense.

Let us now consider the second aspect of a constitutive relation, namely the outputs, responses, and supplies (all taken as one here for simplicity). If the recipient category  $\underline{Y}$  for this has been specified, and also the configuration category  $\underline{C}$ , then from a pragmatic point of view the "only" role of a state category



is to account for observed correlations between processes in  $\underline{C}$  and in  $\underline{Y}$ \*). The above diagram suggests that given only a pair of categories  $\underline{C}, \underline{Y}$  there is a universal state category  $\mathcal{F}_{\underline{C}}(\underline{Y})$  to which all others can be reduced insofar as output into  $\underline{Y}$  is concerned. Indeed this follows from the construction, recently utilized by Grothendieck in homotopy theory [7], of an adjoint pair of functors

$$\text{Cat} \xleftarrow{\mathcal{G}_{\underline{C}}} \mathcal{S}_{\underline{C}}^{\underline{C}} \xrightarrow{\mathcal{D}_{\underline{C}}}$$

$$\begin{array}{c} \text{set}^{\underline{C}} \xleftarrow{\quad} \underline{C}^{\text{op}} \\ \mathcal{G}_{\underline{C}} = R \downarrow \text{Set}^{\underline{C}} \xrightarrow{\quad} \text{Cat} \end{array}$$

\*) Of course the pragmatic point of view contrasts with the materialist view which considers that a state category reflects the objective essence of the thing being studied, so will in particular remain the basis of responses valued in undiscovered  $\underline{Y}'$ . etc.

in which the left adjoint  $\mathcal{G}_{\underline{C}}$  is the Grothendieck construction assigning to any set-valued functor on  $\underline{C}$  the total category of the associated (discrete op-) fibration;  $\mathcal{F}_{\underline{C}} = \mathcal{G}_{\underline{C}} \circ \mathcal{D}_{\underline{C}}$  is the endofunctor of Cat which we seek.

Theorem

There is a fibration  $\mathcal{F}_{\underline{C}}(\underline{Y}) \longrightarrow \underline{C}$  over  $\underline{C}$  equipped with a canonical output  $\mathcal{F}_{\underline{C}}(\underline{Y}) \xrightarrow{\epsilon} \underline{Y}$  to  $\underline{Y}$  such that for any fibration  $\underline{X} \xrightarrow{\pi} \underline{C}$  and any functor  $\underline{X} \xrightarrow{S} \underline{Y}$ , there is a unique morphism of fibrations

$$\begin{array}{ccc} \underline{X} & \xrightarrow{S} & \mathcal{F}_{\underline{C}}(\underline{Y}) \\ \pi \searrow & \swarrow \epsilon & \downarrow \\ \underline{C} & & \end{array}$$

for which  $S = \epsilon \circ \hat{S}$ . In fact the set of states of configuration A in  $\mathcal{F}_{\underline{C}}(\underline{Y})$  is the set

$$\text{Cat } (A/\underline{C}, \underline{Y})$$

of all functors  $\eta$  to  $\underline{Y}$  defined on the category of future deformations which start at A, and the action of any deformation  $A \xrightarrow{\alpha} A'$  on these states is defined by composing functors

$$A'/\underline{C} \xrightarrow{\sum \alpha} A/\underline{C} \xrightarrow{\eta} \underline{Y}$$

with the functor  $\sum_{\alpha}$  of composing in  $\underline{C}$  with  $\alpha$ . The canonical output  $\epsilon$  evaluates a state  $\eta$  at the object  $l_A$  of  $A/\underline{C}$ . If  $\underline{Y}$  and  $\underline{C}$  are equipped with duration functors and we consider responses  $S$ , a universal state category is again obtained by restricting the functors taken as states to be responses themselves. If instead  $\underline{Y}$  is a group, so that a universal state category with respect to supplies of  $\underline{Y}$  is sought, and if we exploit the fact that each category  $A/\underline{C}$  is actually a poset when all morphisms in  $\underline{C}$  are epimorphisms, then a universal state  $(A, \eta)$  can be considered as just a  $\underline{Y}$ -valued mapping defined on the set of all deformations starting from  $A$ .

The above theorem thus clarifies the status of the "definition of state" given by Banfi and Fabrizio [8].

Combining the two extreme examples of state categories we have constructed, we can construct a third.

Proposition

Given two categories  $\underline{C}$ ,  $\underline{Y}$ , a map of fibrations over  $\underline{C}$

$$\mathcal{H}_{fin}(\underline{C}) \longrightarrow \mathcal{F}_{\underline{C}}(\underline{Y})$$

is determined by specifying for each  $C$  an arbitrary functor  $C/C \xrightarrow{\eta_C} \underline{Y}$ . The image of such a map in  $sh(\underline{C})$  is a state category over  $\underline{C}$  with an output  $\epsilon$  to  $\underline{Y}$  satisfying both of the requirements that the states are determined by histories and distinguished by  $\underline{Y}$ -outputs

in the sense that if  $\eta, \eta'$  are two distinct states of the same configuration A, then there is some deformation  $A \xrightarrow{\alpha} B$  such that  $\epsilon(\alpha \cdot \eta_1) \neq \epsilon(\alpha \cdot \eta_2)$ .

The choice of  $\eta_C$  for each C must thus be justified in order to render the above image construction of a state category for C, Y, somewhat canonical. Further investigation is needed to determine whether a choice of the type made by Noll, based on further axioms which relate a fading memory concept to relaxed states, can be characterized algebraically.

Buffalo,  
May 1986

## R e f e r e n c e s

---

- 1 Walter Noll "A new Mathematical Theory of Simple Materials"  
The Foundations of Mechanics and Thermodynamics  
(Selected Papers) Springer 1974, pp 243-292.
- 2 Bernard D. Coleman and David R. Owen  
"Recent Research on the Foundations of Thermo-  
dynamics" in Categories in Continuum Physics  
Springer Lecture Notes # 1174, (1986)  
pp 83 - 99
- 3 S. Eilenberg and S. Mac Lane  
"A General Theory of Natural Equivalences"  
Trans. Amer. Math. Soc. 58, (1945) pp 231-264.
- 4 Pierre Leroux "Les Catégories de Möbius" Cahiers de Topologie  
et géométrie différentielle 16 (1975)  
pp 280 - 282.
- 5 Peter Johnstone Topos Theory 1977 Academic Press
- 5' " " " "How General is a Generalized Space?"  
Dowker Memorial Volume, London Math. Soc.  
Lecture Notes # 93 (1985) pp 77 - 112.
- 6 F. William Lawvere Introduction to Categories in Continuum Physics  
Springer Lecture Notes # 1174, 1986
- 7 A. Grothendieck "In Pursuit of Stacks" 1983
- 8 C. Banfi and M. Fabrizio "Sulla nozione di stato per un elemento  
materiale" Bollettino UMI 15-A (1978)  
pp 405 - 415