# University of Nebraska - Lincoln DigitalCommons@University of Nebraska - Lincoln

Papers in Herpetology

Papers in the Biological Sciences

9-2015

## Spreading Holiday Spirit and Northwestern Salamanders, *Ambystoma gracile* (Baird 1859) (Caudata: Ambystomatidae), Across the USA

Michael R. Rochford University of Florida, miker@ufl.edu

Jeffrey M. Lemm

San Diego Zoo Institute for Conservation Research, cjlemm@cox.net

Kenneth Krysko

Florida Museum of Nahral History, Division of Herpetology, University of Florida, Gainesville, Florida, kenneyk@flmnh.ufl.edu

Louis A. Somma

Florida State Collection of Arthropods, somma@ufl.edu

Robert W. Hansen

Clovis, California, hansenranch2@gmail.com

#### See next page for additional authors

Follow this and additional works at: http://digitalcommons.unl.edu/biosciherpetology

Part of the <u>Aquaculture and Fisheries Commons</u>, <u>Population Biology Commons</u>, <u>Terrestrial and Aquatic Ecology Commons</u>, and the <u>Zoology Commons</u>

Rochford, Michael R.; Lemm, Jeffrey M.; Krysko, Kenneth; Somma, Louis A.; Hansen, Robert W.; and Mazzotti, Frank J., "Spreading Holiday Spirit and Northwestern Salamanders, *Ambystoma gracile* (Baird 1859) (Caudata: Ambystomatidae), Across the USA" (2015). *Papers in Herpetology*. 15.

http://digitalcommons.unl.edu/biosciherpetology/15

This Article is brought to you for free and open access by the Papers in the Biological Sciences at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Papers in Herpetology by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

41	
uthors Iichael R. Rocl Iazzotti	nford, Jeffrey M. Lemm, Kenneth Krysko, Louis A. Somma, Robert W. Hansen, and Frank



## INTRODUCED SPECIES

## Spreading Holiday Spirit and Northwestern Salamanders, *Ambystoma gracile* (Baird 1859) (Caudata: Ambystomatidae), Across the USA

Michael R. Rochford<sup>1</sup>, Jeffrey M. Lemm<sup>2</sup>, Kenneth L. Krysko<sup>3</sup>, Louis A. Somma<sup>3</sup>, Robert W. Hansen<sup>4</sup>, and Frank J. Mazzotti<sup>1</sup>

<sup>1</sup>Fort Lauderdale Research and Education Center, University of Florida, Fort Lauderdale, Florida 33314, USA (MRR: miker@ufl.edu, FJM: fjma@ufl.edu)

<sup>2</sup>San Diego Zoo Institute for Conservation Research, Escondido, California 92027, USA (cjlemm@cox.net)

<sup>3</sup>Division of Herpetology, Florida Museum of Natural History, University of Florida, Gainesville, Florida 32611, USA

(KLK: kenneyk@flmnh.ufl.edu, LAS: somma@ufl.edu)

<sup>4</sup>16333 Deer Path Lane, Clovis, California 93619, USA (hansenranch2@gmail.com)

Introduced species of amphibians and reptiles are of growing conservation concern (Bury and Luckenbach 1976; Krysko et al. 2011; Meshaka 2011). Unintentional introductions result largely from individuals escaping from enclosures or hitch-hiking in cargo or on vehicles, whereas intentional introductions occur primarily because a species is perceived to be of value to humans or from deliberately releasing sick or unwanted animals, including pets (Kraus 2009; Krysko et al. 2011). In Florida, most known introductions are directly related to the pet trade, but about 12% are related to the cargo pathway (Krysko et al. 2011). In this paper, we report the introduction of the Northwestern Salamander, *Ambystoma gracile* (Baird 1859), in Florida from Broward County, and in California from Fresno and San Diego counties, via the cargo pathway.

Ambystoma gracile is indigenous to the North American Cordillera of the Pacific Northwest. It occurs principally west of the hydrologic crest of the Cascades, including the Pacific Ranges and intervening lowlands, from extreme coastal southeastern Alaska, including Chichagof Island, southward, including Vancouver Island, to Sonoma County in northwestern California (Duellman and Sweet 1999; Hoffman et al. 2003; Shaffer 2005). This species may be paedomorphic or metamorphose to terrestrial adults (Snyder 1956). Neotenic larvae may reach 105 mm snout-vent length (SVL) and are commonly found in coniferous forests and meadows (Hoffman et al. 2003).

On 12 December 2014, Eli, Tommy, Anthony, and Erin Khouri collected one adult *Ambystoma gracile* (UF-Herpetology 174498; 7.8 cm SVL, 8.5 cm TL; Fig. 1) found in a Noble Fir (*Abies procera*) Christmas tree at their home in Fort Lauderdale, Broward County, Florida. This tree originated in Oregon and was purchased from Happy Holidays Christmas Trees, Fort Lauderdale, Broward

County, Florida (26.14753°N, 80.120498°W). This represents the first known voucher for this species in Florida.

On 14 December 2014, David Tobler obtained one adult A. gracile (UF-Herpetology 174922; 5.8 cm SVL, 11.1 cm TL) from a Christmas tree lot at The Home Depot, 3272 West Shaw Avenue, Fresno, Fresno County, California (36.80954°N, 119.850770°W). On 6 February 2015, JML received one adult A. gracile (UF-Herpetology 174919; 6.67 cm SVL, 12.83 cm TL; Fig. 2) that was recovered from a box of pine boughs shipped to the Walter Anderson Nursery, 12755 Danielson Court, Poway, San Diego County, California, in mid-December (32.94015°N, 117.046631°W). The box was shipped from Hiawatha Evergreens, 681 East Johns Prairie Road, Shelton, Mason County, Washington (47.24183°N, 123.10382°W). For three consecutive years beginning in about 2000, A. gracile was found in Christmas trees at The Home Depot, 12185 Carmel Mountain Road, San Diego, San Diego County, California (32.98467°N, 117.07501°W), and given to a nearby pet store (J.M. Jones, pers. comm.). Another A. gracile was collected in Santa Barbara, Santa Barbara County, California in about 2009 after falling out of a Christmas tree (S. Sweet, pers. comm.).



**Fig. 1.** Northwestern Salamander, *Ambystoma gracile* (UF-Herpetology 174498), from Fort Lauderdale, Broward County, Florida. Photograph Erin Khouri.



Fig. 2. Northwestern Salamander, *Ambystoma gracile* (UF-Herpetology 174919), from Walter Anderson Nursery, Poway, California. Photograph by J.M. Lemm.

However, no voucher has been deposited. Although *A. gracile* is native to California, these introductions occurred over 300 km from the southernmost part of its native range.

These records represent stage 2 (after Colautti and MacIsaac 2004) introductions. Given the late November and December distribution of conifers originating in the Pacific Northwest, A. gracile could be introduced to many areas across the continental USA. The Northern Pacific Treefrog, Pseudacris regilla (Baird and Girard 1852) was intercepted in Anchorage, Municipality of Anchorage, Alaska, USA, via inspected Christmas trees originating from Orting, Washington, USA, and biologists have expressed concern that these frogs may carry pathogens harmful to native amphibians elsewhere (Halpin 2009). Introduced salamanders could serve as a pathway for Batrachochytrium salamandrivorans, a virus lethal to salamanders (Martel et al. 2013). The Western Tiger Salamander, Ambystoma mavortium Baird 1850 (formerly A. tigrinum mavortium, and elevated to species status by Shaffer and McKnight [1996]), a congener of *A. gracile*, has invaded California where it negatively affects native amphibian species through predation and hybridization (Ryan et al. 2009). Controlling alien species early in the invasion process decreases costs and increases effectiveness and efficiency (Kraus 2009). Increased tree screening prior to shipment might prevent future introductions. Areas where A. gracile might be capable of establishing populations should consider sourcing conifers from regions where this species is absent to prevent similar ecological impacts.

### Acknowledgments

We are grateful to Jake R. Edwards and Edward Mercer for assistance procuring UF-Herpetology 174498; Sean from El Plantio nursery and Celine from Walter Anderson nursery for securing UF-Herpetology 174919; David Tobler for data on UF-Herpetology 174922; Eli, Tommy, Anthony, and Erin Khouri for data on UF-Herpetology 174498. Andrea Adams and Greg Pauly made helpful suggestions that improved this manuscript. We thank Greg Pauly and David Wake for confirming the identifications of UF-Herpetology 174498 and UF-Herpetology 174919.

### Literature Cited

- Bury, R.B. and R.A. Luckenbach. 1976. Introduced amphibians and reptiles in California. *Biological Conservation* 10:1–14.
- Colautti, R.I. and H.J. MacIsaac. 2004. A neutral terminology to define 'invasive' species. Diversity and Distributions 10:135–141.
- Duellman, W.E. and S.S. Sweet. 1999. Distribution patterns of amphibians in the Nearctic region of North America, pp. 31–109. In: W.E. Duellman (ed.), *Patterns of Distribution of Amphibians: A Global Perspective*. Johns Hopkins University Press, Baltimore, Maryland.
- Halpin, J. 2009. Wanted, dead or alive: Frogs riding north on Christmas trees. *Alaska Dispatch News* 2009 (December 17):1–2 (http://www.adn.com/article/20091217/wanted-dead-or-alive-frogs-riding-north-christmas-trees).
- Hoffman, R.L., G.L. Larson, and B.J. Brokes. 2003. Habitat segregation of *Ambystoma gracile* and *Ambystoma macrodactylum* in mountain ponds and lakes, Mount Rainier National Park, Washington, USA. *Journal of Herpetology* 37:24–34.
- Kraus, F. 2009. Alien Reptiles and Amphibians: A Scientific Compendium and Analysis. Springer, Dordrecht, The Netherlands.
- Krysko, K.L., J.P. Burgess, M.R. Rochford, C.R. Gillette, D. Cueva, K.M. Enge, L.A. Somma, J.L. Stabile, D.C. Smith, J.A. Wasilewski, G.N. Kieckhefer III, M.C. Granatosky, and S.V. Nielsen. 2010. Verified non-indigenous amphibians and reptiles in Florida from 1863 through 2010: Outlining the invasion process and identifying invasion pathways and stages. *Zootaxa* 3028:1–64 + MorphoBank Project No. p536 (http://www.morphobank.org/permalink/?P536).
- Martel, A., A. Spitzen-van der Sluijs, M. Blooi, W. Bert, R. Ducatelle, M.C. Fisher, A. Woeltjes, W. Bosman, K. Chiers, F. Bossuyt, and F. Pasmans. 2013. Batrachochytrium salamandrivorans sp. nov. causes lethal chytridiomycosis in amphibians. Proceedings of the National Academy of Sciences of the United States of America 110:15325–15329.
- Meshaka, W.E., Jr. 2011. A runaway train in the making: The exotic amphibians, reptiles, turtles, and crocodilians of Florida. Monograph 1. Herpetological Conservation and Biology 6:1–101.
- Ryan, M.E., J.R. Johnson, and B.M. Fitzpatrick. 2009. Invasive hybrid Tiger Salamander genotypes impact native amphibians. *Proceedings of the National Academy of Sciences of the United States of America* 106:11166–11171.
- Shaffer, H.B. 2005. Ambystoma gracile (Baird, 1859) Northwestern Salamander, pp 609–611. In: M. Lannoo (ed.), Amphibian Declines: The Conservation Status of United States Species. University of California Press, Berkeley.
- Shaffer, H.B. and M.L. McKnight. 1996. The polytypic species revisited: Genetic differentiation and molecular phylogenetics of the Tiger Salamander Ambystoma tigrinum (Amphibia: Caudata) complex. Evolution 50:417–433.