

ASTR 610: Problem Set 3

This problem set consists of 4 problems.

Due date: Fri Oct 23, 2020

Problem 1: Spherical Collapse [6 points]

According to the SC model, the parametric solution to the evolution of a mass shell is

$$r = A(1 - \cos \theta)$$

$$t = B(\theta - \sin \theta)$$

where $A^3 = G M B^2$, which implies that

$$1 + \delta = \frac{9}{2} \frac{(\theta - \sin \theta)^2}{(1 - \cos \theta)^3}$$

Show that at early times (when $\theta \ll 1$) one has that

$$\delta_i = \frac{3}{20} (6\pi)^{2/3} \left(\frac{t_i}{t_{\max}} \right)^{2/3}$$

Hint: use Taylor series expansions.

Problem 2: The Zel'dovich Approximation

In this problem we seek to characterize the displacement $\psi(t)$ defined by

$$\vec{x}(t) = \vec{x}_i + \psi(t)$$

where $\vec{x}(t)$ is the comoving coordinate of a particle. Obviously we have that

$$\psi(t) = \int_{t_i}^t \frac{v(t)}{a(t)} dt$$

where $v(t)$ is the particle's peculiar velocity. Under the Zel'dovich approximation, the gradient of the potential (which defines the direction in which the particle moves), can be written as $\nabla\Phi(t) = f(t)\nabla\Phi_i$, where $f(t)$ is some function (to be determined) of time.

a) [4 points] Use the linearized Euler equation for a pressureless fluid to show that

$$\frac{d}{dt}(a\vec{v}) = -\nabla\Phi$$

b) [6 points] Use the fact that, at early times, the Universe behaves as an EdS cosmology to show that

$$\vec{v} = -\frac{\nabla\Phi_i}{a} \int \frac{D(a)}{a} dt$$

c) [6 points] Use the fact that $D(a)$ is a solution of the linearized fluid equation of a pressureless fluid to show that

$$\frac{D(a)}{a} = \frac{1}{4\pi G \bar{\rho}_i} \frac{d(a^2 \dot{D})}{dt}$$

Hint: you may use that the scale factor is normalized such that $a_i = 1$.

d) [6 points] Use the above results to show that the displacement

$$\psi(t) = -\frac{D(a)}{4\pi G \bar{\rho}_i} \nabla\Phi_i$$

Problem 3: The two-point correlation function and σ_8

Let M be the mass inside a top-hat filter. The expectation value for M , i.e., the average value obtained by putting down the top-hat filter at many different locations, is simply $\langle M \rangle = \bar{\rho} V$ where V is the volume of the top-hat. Similarly, one can show that

$$\langle M^2 \rangle = \langle M \rangle^2 + \frac{\langle M \rangle^2}{V^2} \int_V \xi(|\vec{x}_1 - \vec{x}_2|) d^3\vec{x}_1 d^3\vec{x}_2$$

a) [5 points] Show that the mass variance, $\sigma^2(M)$, can be written as

$$\sigma^2(M) = \frac{3}{R^3} \int_0^R \xi(r) r^2 dr$$

where $R = R(M)$ is the size of the top-hat filter.

b [6 points] The first measurements of the two-point correlation function of galaxies revealed a power-law $\xi(r) = (r/r_0)^\gamma$ with $r_0 = 5h^{-1}\text{Mpc}$ and $\gamma = -1.8$. Under the assumption that galaxies are unbiased tracers of the mass distribution, what does this imply for the value of σ_8 ?

Problem 4: Power spectrum and Mass variance

Let the matter power spectrum be a pure power-law, $P(k) \propto k^n$.

a) [5 points] Using a sharp k -space filter, show that the mass variance $\sigma^2(M) \propto M^\gamma$, and give the relation between γ and n .

b [3 points] Repeat the same exercise as under **(a)**, but this time using a Gaussian filter.

c [3 points] Give the ratio of the mass variances computed using the Gaussian filter and the sharp k -space filter for the case $n = 1$. Do NOT use mathematica (or similar), but first express your answer in terms of a special function, prior to giving the numerical value of the ratio.