


Lecture #14

In Vivo MRS-detectable Metabolites




- Topics
 - ^1H -MRS
 - ^{31}P -MRS
 - ^{13}C -MRS
 - Other nuclei
- Handouts and Reading assignments
 - de Graaf, Chapter 2.
 - de Graaf, Chapter 3, 158-171.

Introduction

MRI: anatomy and structure



MRS: metabolism and function




MR of the Brain

Anatomy




Conventional
MRI

Microvasculature




Perfusion
imaging

Microstructure




DWI/DTI

Cellular function and
metabolism






MRS



Cells


Neuron



General cellular functions

- Turn glucose into energy
- Maintain cellular integrity
 - Ionic balance
 - Osmotic balance
 - Membrane structure
- Perform cell specific functions, e.g. neurons send/receive electrical signals

^1H Brain Spectroscopy







Some general rules of thumb

- In vivo MRS does not see everything! Maudsley paper is rather optimistic.
- Concentration limit (of protons) around 1 mM for NMR-detection
- Macromolecules are non-detectable (T_2 s too short)




^1H Brain Spectroscopy



**GE MRS
phantom**




**In vitro
solutions**






^1H Brain Spectra


We'll start with
"late" echo spectra



N-acetyl Aspartate (NAA)




Biochemical role?




NAA

- Largest peak in a ^1H -MRS brain spectrum
- Neuronal marker
- 2.0 ppm
- Approx 10 mM


Creatine (Cre)



“Cre” peak from both
creatine and
phosphocreatine (often
called referred to as total
creatine peak)




Biochemical role




- Reflects cellular energetics
- 3.0 ppm
- $\text{Cr} + \text{PCr} = \text{tCr}$
- Approx 5-10 mM


Choline (Cho)



“Cho” peak from several choline containing compounds: phosphocholine, glycerphosphocholine, etc.



Biochemical role




Cho


- Largely reflects cell membrane repair and synthesis (acetylcholine component very small)
- 3.2 ppm
- Approx 1-2.5 mM

Example: Brain Tumor



52 y.o male: MRI #1 - rule out stroke, MRI # - tumor?





Lactate (Lac)



Lac




Biochemical role



Lac


- Indicator of anaerobic metabolism
- Doublet at 1.3 ppm
- Approx 0.4 mM (normal brain tissue)

Cerebral infarct
Non-selective SE, TE=288 ms (2/J)



Kamada, et al, *J. N Psych*, 70:675, 2001.

Short TE ^1H -MRS Brain Spectra






1.5T ^1H -MRS In Vivo Brain Spectrum


TE = 35 ms

myo-Inositol (mI)

1.5 T In vivo ^1H Brain Spectrum (TE=35ms)





Biochemical role?




mI

- Suggested as a glial marker.
- Glial cells provide structural support for neurons in addition to involvement in neurotransmitter processes.
- Largest peak at 3.6 ppm
- Approx 4-8 mM

Neurotransmitters




Glutamate is the major excitatory neurotransmitter in the brain




Glutamate (Glu) and Glutamine (Gln)


1.5 T In vivo ^1H Brain
Spectrum (TE=35ms)




Glu




Gln



Glu



Gln




In vivo concentrations

8-10 mM


2-3 mM

γ -aminobutyric acid (GABA)



GABA is the major inhibitory neurotransmitter in the brain
(conc: 1-2 mM)




A real neural net





GABA





Example In vivo GABA (3T)





Muscle and Fat



Prostate




Normal prostate (1.5 T)



Prostate cancer (1.5 T)


Breast

Elevated choline



Breast cancer

2D-J spectrum



Noncancerous lesion

Sensitivity





Table 1.1. NMR properties of commonly encountered nuclei in *in vivo* NMR

Isotope	Spin	Gyromagnetic ratio ($10^7 \text{ rads}^{-1}\text{T}^{-1}$)	NMR frequency at 2.35 T (MHz)	Natural abundance (%)	Relative sensitivity ¹
¹ H	1/2	26.752	100.000	99.985	1.00
² H	1	4.107	15.351	0.015	1.45×10^{-6}
¹³ C	1/2	6.728	25.145	1.108	1.76×10^{-4}
¹⁴ N	1	1.934	7.228	99.630	1.01×10^{-3}
¹⁵ N	1/2	-2.712	10.137	0.370	3.85×10^{-6}
¹⁹ F	1/2	25.181	94.094	100.000	0.833
²³ Na	3/2	7.080	26.466	100.000	9.27×10^{-2}
³¹ P	1/2	10.841	40.481	100.000	6.65×10^{-2}
³⁹ K	3/2	1.250	4.672	93.100	4.75×10^{-4}

¹Relative sensitivity is calculated as the product of NMR sensitivity (proportional to $\gamma^3 I(I+1)$) and the natural abundance.

$$\text{in vivo SNR} \propto \rho \frac{\gamma^2 \hbar^2 B_0}{4kT}$$

^{31}P Spectroscopy



^{13}C Spectroscopy

Table 2.3. Chemical shifts of biological relevant ^{13}C containing metabolites



Compound	Carbon atom					
	C1	C2	C3	C4	C5	C6
Acetate	182.6	24.5				
Alanine	176.6	51.5	17.1			
Aspartate	175.1	53.0	37.4	178.4		
Citrate	179.7	46.8	76.0	46.8	179.7	182.3
Creatine	175.4	37.8	158.0	54.7		
GABA	182.3	35.2	24.6	40.2		
Glycerol	63.6	73.3	63.6			
β -hydroxy butyrate	181.2	47.6	66.8	22.9		
Glucose α	92.7	72.1	73.5	70.4	72.1	61.4
β	96.6	79.9	76.5	70.4	76.5	61.4
Glutamate	175.3	55.5	27.8	34.2	182.0	
Glutamine	174.8	55.0	27.1	31.7	178.4	
Glycine	173.3	42.5				
Glycogen	100.5	—	74.0	78.1	72.1	61.4
Inositol	73.3	73.1	73.3	71.9	75.1	71.9
Lactate	183.3	69.3	21.0			
Malate	182.1	71.7	43.9	180.9		
NAA	179.7	54.0	40.3	179.7	174.3	22.8
Succinate	183.4	35.3	35.3	183.4		
Taurine	48.4	36.2				

Note: very
wide chemical
shift range

All chemical shifts are referenced relative to 3-(trimethylsilyl)-1-propanesulfonic acid at 0.00 ppm.

^{13}C natural abundance is low, but ^{13}C -labeled glucose or acetate infusions can yield important information regarding metabolic fluxes


Example: ^{13}C -glucose Infusion




Gruetter, et al., *J. Neurochem.*, 63:1377-1385, 1994.

Other Nuclei

- Sodium (^{23}Na , spin=3/2)
 - Intracellular 10 mM
 - Extracellular 150 mM



Involved in ionic balances,
generation of action potentials,
regulations of cell volume.



- Fluorine (^{19}F)
 - No endogenous ^{19}F compounds in biological tissue
 - Lots of fluorinated drugs!

Next Lecture: ^1H MRS Methods