Two Cases of Hypersensitivity Reactions to Neptazol in Clinical Practice

Reyes M, Okonkwo B.

Clinical Allergy Notes, Vol 7(2), 2026

Abstract

We report two separate cases of hypersensitivity reactions following Neptazol administration, observed six months apart at the same clinic, illustrating a spectrum of severity.

Introduction

Adverse drug reactions remain an important area of post-marketing surveillance. We describe a case observed in clinical practice and discuss its implications.

Case Report

Case 1: A 52-year-old woman was prescribed Neptazol 20mg daily for hypertension. Ten days into treatment she developed a generalized urticarial rash and mild lip swelling. She was treated with oral antihistamines as an outpatient and the drug was discontinued; symptoms resolved within 48 hours. Case 2: A 67-year-old man on Neptazol 20mg daily for three weeks presented to the emergency department with facial angioedema and difficulty swallowing. He was treated with intramuscular epinephrine and intravenous corticosteroids and observed overnight before discharge. Neptazol was permanently discontinued and he was advised to avoid the drug class going forward.

Discussion

These two cases, differing markedly in severity, suggest clinicians should counsel all patients starting Neptazol on early signs of hypersensitivity, regardless of how mild an initial presentation might seem.

References

1. Allergy Reporting Consortium. Drug hypersensitivity registry, 2025 annual report.