

Two-way fixed effects estimators with heterogeneous treatment effects*

Clément de Chaisemartin[†] Xavier D’Haultfoeulle[‡]

March 5, 2020

Abstract

Linear regressions with period and group fixed effects are widely used to estimate treatment effects. We show that they estimate weighted sums of the average treatment effects (ATE) in each group and period, with weights that may be negative. Due to the negative weights, the linear regression coefficient may for instance be negative while all the ATEs are positive. We propose another estimator that solves this issue. In the two applications we revisit, it is significantly different from the linear regression estimator.

Keywords: linear regressions, fixed effects, heterogeneous treatment effects, difference-in-differences.

JEL Codes: C21, C23

*We are very grateful to Olivier Deschênes, Guido Imbens, Peter Kuhn, Kyle Meng, Jesse Shapiro, Dick Startz, Doug Steigerwald, Clémence Tricaud, Gonzalo Vazquez-Bare, members of the UCSB econometrics research group, and seminar participants at Bergen, CIREQ Econometrics conference, CREST, Goteborg, Gothenburg, Groningen, ITAM, Pompeu Fabra, Stanford, SMU, Tinbergen Institute, UCL, UCLA, UC Davis, UCSB, USC, and Warwick for their helpful comments. Xavier D’Haultfoeulle gratefully acknowledges financial support from the research grants Otelo (ANR-17-CE26-0015-041) and the Labex Ecodec: Investissements d’Avenir (ANR-11-IDEX-0003/Labex Ecodec/ANR-11-LABX-0047).

[†]University of California at Santa Barbara, clementdechaisemartin@ucsb.edu

[‡]CREST-ENSAE, xavier.dhaultfoeulle@ensae.fr

Here is some intuition as to why one weight is negative in this example. It follows from Equation (7) in the proof of Theorem 1 (see also Theorem 1 in Goodman-Bacon, 2018) that in this simple example, $\beta_{fe} = (\text{DID}_1 + \text{DID}_2)/2$, with

$$\begin{aligned}\text{DID}_1 &= E(Y_{2,2}) - E(Y_{2,1}) - (E(Y_{1,2}) - E(Y_{1,1})), \\ \text{DID}_2 &= E(Y_{1,3}) - E(Y_{1,2}) - (E(Y_{2,3}) - E(Y_{2,2})).\end{aligned}$$

The first DID compares the evolution of the mean outcome from period 1 to 2 in group 2 and in group 1. The second one compares the evolution of the mean outcome from period 2 to 3 in group 1 and in group 2. The control group in the second DID, group 2, is treated both in the pre and in the post period. Therefore, under the common trends assumption, it follows from Lemma 1 in Appendix A (a similar result appears in Lemma 1 of de Chaisemartin (2011) and in Equation (13) of Goodman-Bacon (2018)) that $\text{DID}_1 = E[\Delta_{2,2}]$, but

$$\text{DID}_2 = E[\Delta_{1,3}] - (E[\Delta_{2,3}] - E[\Delta_{2,2}]).$$

DID_2 is equal to the ATE in group 1 period 3, minus the change in group 2's ATE between periods 2 and 3. Intuitively, the mean outcome of groups 1 and 2 may follow different trends from period 2 to 3 either because group 1 becomes treated, or because group 2's ATE changes. The intuition that negative weights arise because $\hat{\beta}_{fe}$ uses treated observations as controls also appears in Borusyak and Jaravel (2017).

We now generalize the previous illustration by characterizing the (g, t) cells whose ATEs are weighted negatively by β_{fe} .

Proposition 1 *Suppose that Assumption 1 holds and for all $t \geq 2$ $N_{g,t}/N_{g,t-1}$ does not vary across g . Then, for all (g, t, t') such that $D_{g,t} = D_{g,t'} = 1$, $D_{.,t} > D_{.,t'}$ implies $w_{g,t} < w_{g,t'}$. Similarly, for all (g, g', t) such that $D_{g,t} = D_{g',t} = 1$, $D_{g,.} > D_{g',.}$ implies $w_{g,t} < w_{g',t}$.*

Proposition 1 shows that β_{fe} is more likely to assign a negative weight to periods where a large fraction of groups are treated, and to groups treated for many periods. Then, negative weights are a concern when treatment effects differ between periods with many versus few treated groups, or between groups treated for many versus few periods.

Proposition 1 has interesting implications in staggered adoption designs, a special case of sharp designs defined as follows.

Assumption 6 (*Staggered adoption designs*) *For all g , $D_{g,t} \geq D_{g,t-1}$ for all $t \geq 2$.*

Assumption 6 is satisfied in applications where groups adopt a treatment at heterogeneous dates (see e.g. Athey and Stern, 2002). In that design, Borusyak and Jaravel (2017) show that β_{fe} is more likely to assign a negative weight to treatment effects at the last periods of the panel.