

# Difference-in-Differences with Multiple Time Periods\*

Brantly Callaway<sup>†</sup>

Pedro H. C. Sant’Anna<sup>‡</sup>

December 1, 2020

## Abstract

In this article, we consider identification, estimation, and inference procedures for treatment effect parameters using Difference-in-Differences (DiD) with (i) multiple time periods, (ii) variation in treatment timing, and (iii) when the “parallel trends assumption” holds potentially only after conditioning on observed covariates. We show that a family of causal effect parameters are identified in staggered DiD setups, even if differences in observed characteristics create non-parallel outcome dynamics between groups. Our identification results allow one to use outcome regression, inverse probability weighting, or doubly-robust estimands. We also propose different aggregation schemes that can be used to highlight treatment effect heterogeneity across different dimensions as well as to summarize the overall effect of participating in the treatment. We establish the asymptotic properties of the proposed estimators and prove the validity of a computationally convenient bootstrap procedure to conduct asymptotically valid simultaneous (instead of pointwise) inference. Finally, we illustrate the relevance of our proposed tools by analyzing the effect of the minimum wage on teen employment from 2001–2007. Open-source software is available for implementing the proposed methods.

**JEL:** C14, C21, C23, J23, J38.

**Keywords:** Difference-in-Differences, Dynamic Treatment Effects, Doubly Robust, Event Study, Variation in Treatment Timing, Treatment Effect Heterogeneity, Semi-Parametric.

---

\*First complete version: March 23, 2018. A previous version of this paper has been circulated with the title “Difference-in-Differences with Multiple Time Periods and an Application on the Minimum Wage and Employment”. We thank the Editor, the Associate Editor, two anonymous referees, Stéphane Bonhomme, Carol Caetano, Sebastian Calonico, Xiaohong Chen, Clément de Chaisemartin, Xavier D’Haultfoeuille, Bruno Ferman, John Gardner, Andrew Goodman-Bacon, Federico Gutierrez, Sukjin Han, Hugo Jales, Andrew Johnston, Vishal Kamat, Qi Li, Tong Li, Jason Lindo, Catherine Maclean, Matt Masten, Magne Mogstad, Tom Mroz, Aureo de Paula, Jonathan Roth, Donald Rubin, Bernhard Schmidpeter, Yuya Sasaki, Na’Ama Shenhav, Tymon Słoczyński, Sebastian Tello-Trillo, Alex Torgovitsky, Jeffrey Wooldridge, Haiqing Xu and several seminar and conference audiences for comments and suggestions. Code to implement the methods proposed in the paper is available in the R package `did` which is available on CRAN.

<sup>†</sup>Department of Economics, University of Georgia. Email: brantly.callaway@uga.edu

<sup>‡</sup>Department of Economics, Vanderbilt University. Email: pedro.h.santanna@vanderbilt.edu

**Theorem 1.** *Let Assumptions 1, 2, 3 and 6 hold.*

(i) *If Assumption 4 holds, then, for all  $g$  and  $t$  such that  $g \in \mathcal{G}_\delta$ ,  $t \in \{2, \dots, \mathcal{T} - \delta\}$  and  $t \geq g - \delta$ ,*

$$ATT(g, t) = ATT_{ipw}^{nev}(g, t; \delta) = ATT_{or}^{nev}(g, t; \delta) = ATT_{dr}^{nev}(g, t; \delta).$$

(ii) *If Assumption 5 holds, then, for all  $g$  and  $t$  such that  $g \in \mathcal{G}_\delta$ ,  $t \in \{2, \dots, \mathcal{T} - \delta\}$  and  $g - \delta \leq t < \bar{g} - \delta$ ,*

$$ATT(g, t) = ATT_{ipw}^{ny}(g, t; \delta) = ATT_{or}^{ny}(g, t; \delta) = ATT_{dr}^{ny}(g, t; \delta).$$

Theorem 1 is the first main result of this paper. It provides powerful identification results that extend the DiD identification results based on the outcome regression approach of Heckman et al. (1997, 1998), the IPW approach of Abadie (2005), and the DR approach of Sant’Anna and Zhao (2020) to the multiple-periods, multiple groups setup. In other words, Theorem 1 says that, from an identification point of view, one can recover the  $ATT(g, t)$ ’s by exploiting different parts of the data generating process: the OR approach only relies on modeling the conditional expectation of the outcome evolution for the comparison groups, the IPW approach relies on modeling the conditional probability of being in group  $g$ , whereas the DR approach exploits both OR and IPW components.

In order to extend the results of Heckman et al. (1997, 1998), Abadie (2005), and Sant’Anna and Zhao (2020) to the multiple groups, multiple periods framework, we have to address two different challenges: one associated with an appropriate reference time period and one associated with an appropriate comparison group. Theorem 1 highlights how a solution to these challenges is directly connected to the limited anticipation and the conditional parallel trends assumptions. More specifically, Theorem 1 says that we can use the time period  $t = g - \delta - 1$  as an appropriate reference time period under Assumption 3 and either Assumption 4 or 5. This is the most recent time period when untreated potential outcomes are observed for units in group  $g$ . Interestingly, the more treatment anticipation is allowed (i.e., the higher  $\delta$  is), the further back in time one needs to go.<sup>9</sup> Theorem 1 also suggests that the choice of comparison group is directly tied to the conditional parallel trends assumption one makes: under Assumption 4, one can use “never treated” units as a fixed comparison group for all “eventually treated” units; whereas, under Assumption 5, one can use the “not-yet-treated by time  $t + \delta$ ” units as a valid comparison group for those who are first treated at time  $g$ . In this latter case, Theorem 1 also highlights that when all units eventually gets treated ( $\bar{g} < \infty$ ), one is only able to identify the  $ATT(g, t)$ ’s for time periods before the last treated group “effectively” starts their treatment, i.e.,  $t < \bar{g} - \delta$ . In this case, one can not identify the  $ATT(g, t)$  for the last treated cohort, too.

Finally, we note that when pre-treatment covariates play no role in identification (i.e., Assumptions 3, 4, and 5 hold unconditionally on  $X$ ), (2.2)-(2.4) collapse to

$$ATT_{unc}^{nev}(g, t; \delta) = E[Y_t - Y_{g-\delta-1} | G_g = 1] - E[Y_t - Y_{g-\delta-1} | C = 1], \quad (2.8)$$

and (2.5)-(2.7) collapse to

$$ATT_{unc}^{ny}(g, t; \delta) = E[Y_t - Y_{g-\delta-1} | G_g = 1] - E[Y_t - Y_{g-\delta-1} | D_{t+\delta} = 0]. \quad (2.9)$$

---

<sup>9</sup>As mentioned in Remark 1, as one allows  $\delta$  to increase, Assumptions 4 and 5 becomes more restrictive.