

Pizza Sales

Basic:

1. Retrieve the total number of orders placed.

select

count(order_id) as total_orders

from orders

Results		Messages	
	total_orders		
1	4		

2. Calculate the total revenue generated from pizza sales.

select

round(sum(o.quantity*p.price),2) as total_sales

from order_details o

join pizzas p

on p.pizza_id = o.pizza_id

Results		Messages	
	total_sales		
1	817860.05		

3. Identify the highest-priced pizza.

select top 1

pt.name_,

p.price

from pizza_types pt

```

join pizzas p
on pt.pizza_type_id = p.pizza_type_id
order by p.price desc

```

Results		Messages
	name_	price
1	The Greek Pizza	35.95

4. Identify the most common pizza size ordered.

```

select top 1
count(o.order_details_id) as order_count
from pizzas p
join order_details o
on p.pizza_id = o.order_details_id
group by p.size
order by order_count desc

```

	size	count(order_details.order_details_id)
▶	M	15385
	L	18526
	S	14137
	XL	544
	XXL	28

5. List the top 5 most ordered pizza types along with their quantities.

```

select top 5
pt.name_,
sum(o.quantity) as quantity
from pizza_types pt
join pizzas p

```

```

on pt.pizza_type_id = p.pizza_type_id
join order_details o
on o.pizza_id = p.pizza_id
group by pt.name_ order by quantity

```

Results		Messages
	name_	quantity
1	The Brie Carre Pizza	490
2	The Mediterranean Pizza	934
3	The Calabrese Pizza	937
4	The Spinach Supreme Pizza	950
5	The Soppressata Pizza	961

Intermediate

- Join the necessary tables to find the total quantity of each pizza category ordered

```

SELECT pt.category,
       SUM(o.quantity) AS quantity
FROM pizza_types pt
JOIN pizzas p
  ON pt.pizza_type_id = p.pizza_type_id
JOIN order_details o
  ON o.pizza_id = p.pizza_id
GROUP BY pt.category
ORDER BY quantity DESC;

```

Results		Messages
	category	quantity
1	Classic	13529
2	Supreme	11987
3	Veggie	11649
4	Chicken	11050
5	Mushroom	1359

7. Determine the distribution of orders by hour of the day

```
SELECT DATEPART(HOUR, order_time) AS hour,  
       COUNT(order_id) AS order_count  
FROM orders  
GROUP BY DATEPART(HOUR, order_time);
```

Results		Messages
	hour	order_count
1	23	28
2	15	1468
3	9	1
4	12	2520
5	21	1198
6	18	2399
7	10	8
8	19	2009
9	13	2455
10	22	663
11	16	1920
12	17	2336
13	11	1231
14	20	1642
15	14	1472

8. Join relevant tables to find the category-wise distribution of pizzas.

```
SELECT category,  
       COUNT(name_) AS total_types  
FROM pizza_types  
GROUP BY category;
```

Results		Messages
	category	total_types
1	Mushroom	1
2	Chicken	6
3	Classic	7
4	Supreme	9
5	Veggie	9

9. Group the orders by date and calculate the average number of pizzas ordered per day.

```
SELECT avg(quantity) as avg_quantity
FROM (
    SELECT orders.order_data,
    SUM(order_details.quantity) AS quantity
    FROM orders
    JOIN order_details ON orders.order_id =
order_details.order_id
    GROUP BY orders.order_data
) AS order_quantity;
```

Results		Messages
	avg_quantity	
1	138	

10. Determine the top 3 most ordered pizza types based on revenue.

```
SELECT top 3
pizza_types.name_,
SUM(order_details.quantity * pizzas.price) AS
revenue
FROM pizza_types
JOIN pizzas ON pizzas.pizza_type_id =
pizza_types.pizza_type_id
```

JOIN order_details ON order_details.pizza_id =
pizzas.pizza_id
GROUP BY pizza_types.name_
ORDER BY revenue DESC

 Results  Messages

	name_	revenue
1	The Thai Chicken Pizza	43434.25
2	The Barbecue Chicken Pizza	42768
3	The California Chicken Pizza	41409.5