Board Level Exercise

Type (I): Very Short Answer Type Questions:

01 Mark Each

1. Write the principal value of $sec^2(-2)$.

2. If
$$tan^{-1}\sqrt{3} + cot^{-1}(x) = \frac{\pi}{2}$$
, find x .

Type (II): Short Answer Type Questions:

02 Mark Each

1. If
$$sin^{-1}(x) + cos^{-1}(\frac{1}{2}) = \frac{\pi}{2}$$
. then find x .

2. Solve for
$$x$$
: $cos(2sin^{1}x) = \frac{1}{9}, x > 0$.

Type (III): Long Answer Type Questions:

04 Mark Each

1. Solve the following for
$$x: tan^1\left[\frac{1+x}{1-x}\right] = \frac{\pi}{4} + tan^1x, 0 < x < 1.$$

2. Solve the
$$x : cos^{-1}x + sin^{1}(\frac{x}{2}) = \frac{\pi}{6}$$

3. Prove the following :
$$2tan^{-1}\frac{1}{3} + tan^{-1}\frac{1}{7} = \frac{\pi}{4}$$
.

4. Prove the following :
$$cos\left(sin^{-1}\frac{3}{5} + cot^{-1}\frac{3}{2}\right) = \frac{6}{5\sqrt{13}}$$

Type (IV): Very Long Answer Type Questions :

06 Mark Each

1. if
$$tan^{-1}x + tan^{-1}y + tan^{-1}z = \pi$$
, prove that $x + y + z = xyz$.

2. Prove that :
$$sin^{-1}\left(\frac{4}{5}\right) + sin^{-1}\left(\frac{5}{13}\right) + sin^{-1}\left(\frac{16}{65}\right) = \frac{\pi}{2}$$
.

3. Solve the following for
$$x$$
: $tan^{-1}x + 2cot^{-1}x = \frac{2\pi}{3}$

4. Solve for
$$x : tan^{-1}\frac{x}{2} + tan^{-1}\frac{x}{3} = \frac{\pi}{4}; \sqrt{6} > x > 0.$$

5. Prove that :
$$tan^{-1}\frac{1}{4} + tan^{-1}\frac{2}{9} = \frac{1}{2}tan^{-1}\frac{4}{3}$$
.

6. Prove that :
$$2tan^{-1}\frac{3}{4} - tan^{-1}\frac{17}{31} = \frac{\pi}{4}$$
.

7. Solve for
$$x$$
: $tan^{-1}\left(\frac{2x}{1-x^2}\right) + cot^{-1}\left(\frac{1-x^2}{2x}\right) = \frac{\pi}{3}, -1 < x < 1.$

- 8. Solve for $x : tan^{-1}(x+2) + tan^{-1}(x-2) = tan^{-1}\left(\frac{8}{79}\right); x > 0.$
- 9. Prove that : $tan^{-1}(1) + tan^{-1}(2) + tan^{-1}(3) = \pi$.
- 10. Prove the following: $tan^{-1}\left(\frac{3}{4}\right) + tan^{-1}\left(\frac{3}{5}\right) tan^{-1}\left(\frac{8}{19}\right) = \frac{\pi}{4}$.

Exercise # 1

PART - I : SUBJECTIVE QUESTIONS

A Definition, graphs and fundamentals

A-1 Find the simplified value of each of the following inverse trigonometric terms :

(i)
$$sin^{-1}\left(\frac{1}{2}\right)$$

(iii)
$$cosec^{-1}\left(-\frac{1}{2}\right)$$

(ii)
$$cos^{-1}\left(\frac{\sqrt{3}}{2}\right)$$

A-2 Find the simplified value of the following expressions:

(i)
$$sin\left[\frac{\pi}{3} - sin^{-1}\left(-\frac{1}{2}\right)\right]$$

(iii)
$$sin^{-1} \left[\left\{ sin^{-1} \left(\frac{\sqrt{3}}{2} \right) \right\} cos \right]$$

(ii)
$$tan \left[cos^{-1} \frac{1}{2} + tan^{-1} \left(-\frac{1}{\sqrt{3}} \right) \right]$$

A-3 Draw the graph of the following functions:

(i)
$$y = \sin^{-1}(x+1)$$

(iii)
$$y = tan^{-1}(2x - 1)$$

(ii)
$$y = cos^{-1}(3x)$$

A-4 Solve the following inequalities:

(i)
$$sin^{-1}x > -1$$

(iii)
$$\cot^{-1}x < -\sqrt{3}$$

(ii)
$$\cos^{-1}x < 2$$

A-5

- (i) If $\sum_{i=1}^{n} \cos^{-1} \alpha_i = 0$, then find the value of $\sum_{i=1}^{n} i\alpha_i$
- (ii) If $\sum_{i=1}^{2n} \sin^{-1} x_i = n\pi$, then show that $\sum_{i=1}^{2n} x_i = 2n$

B Trig. (trig -1 x), trig -1 (trig x) trig (-x)

B-1 Evaluate the following expressions:

(i)
$$sin\left(cos^{-1}\frac{3}{5}\right)$$

(iv)
$$tan \left(cosec^{-1} \frac{65}{63} \right)$$

(ii)
$$tan\left(cos^{-1}\frac{1}{3}\right)$$

(v)
$$sec\left(tan\left\{tan^{-1}\left(-\frac{\pi}{3}\right)\right\}\right)costan^{-1}sincot^{-1}\frac{1}{2}$$

(iii)
$$cosec \left(sec^{-1} \frac{\sqrt{41}}{5} \right)$$

B-2 Evaluate the following inverse trigonometric expressions:

(i)
$$sin^{-1} \left(sin \frac{7\pi}{6} \right)$$

(iii)
$$cos^{-1}\left(cos\frac{5\pi}{4}\right)$$

(ii)
$$tan^{-1}\left(tan\frac{2\pi}{3}\right)$$

(iv)
$$sec^{-1}\left(sec\frac{7\pi}{4}\right)$$

B-3 Find the value of the following inverse trigonometric expressions:

(i)
$$sin^{-1}(sin4)$$

(iv)
$$\cot^{-1}(\cot(-10))$$

(ii)
$$cos^{-1}(cos10)$$

(v)
$$\cos^{-1}\left(\frac{1}{\sqrt{2}}\left(\cos\frac{9\pi}{10} - \sin9\pi 10\right)\right)$$

(iii)
$$tan^{-1}(tan(-6))$$

B-4

Express $sin^1(sin\theta, cos1(cos\theta, tan1(tan\theta \text{ and } cot1(cot\theta \text{ in terms of linear expression of } \theta \text{ for } \theta \epsilon \left[\frac{3\pi}{2}, 3\pi\right]$

C Property " $\frac{\pi}{2}$ ", Addition and subtraction rule, miscellaneous formula, summation of series

C-1 Find the value of following expressions:

(i)
$$\cot(\tan^{-1}a + \cot^{-1}a)$$

(iii)
$$tan \left[cos^{-1} \left(\frac{3}{4} \right) + sin^{-1} \left(\frac{3}{4} \right) - sec^{-1} 3 \right]$$

(ii)
$$sin(sin^{-1}x + cos^{-1}x), |x| \le 1$$

C-2 Prove that

(i)
$$sin^{-1}\left(\frac{3}{5}\right) + sin^{-1}\left(\frac{8}{17}\right) = sin^{-1}\frac{77}{85}$$

(ii)
$$\cos^{-1}\frac{4}{5} + \cos^{-1}1213 = \cos^{-1}\frac{33}{65}$$

(iii)
$$sin^{-1} \left(\frac{1}{\sqrt{5}} \right) + cot^{-1} 3 = \frac{\pi}{4}$$

(iv)
$$tan^{-1}\left(\frac{1}{3}\right) + tan^{-1}\left(\frac{1}{5}\right) + tan^{-1}\left(\frac{1}{7}\right) + tan^{-1}\left(\frac{1}{8}\right) = \frac{\pi}{4}$$

C-3 Simplify $tan^{-1} \left\{ \frac{1}{2} sin^{-1} \left(\frac{2x}{1+x^2} \right) + \frac{1}{2} cos^{-1} \left(\frac{1-y^2}{1+y^2} \right) \right\}$, if x > y > 1.

C-4 Find the value of $sin^{-1}(cos(sin^{-1}x)) + cos^{-1}(sin(cos^{-1}x))$

D Inverse trigonometric function Equations:

D-1 Solve for x

(i)
$$cos(2sin^{-1}x = \frac{1}{3}$$

(ii)
$$\cot^{-1}x + \tan^{-1}3 = \frac{\pi}{2}$$

D-2 Solve the following equations:

(i)
$$tan^{-1}\left(\frac{x-1}{x-2}\right) + \frac{1}{2}tan^{-1}\left(\frac{x+1}{x+2}\right) = \frac{\pi}{4}$$

(ii)
$$sin^{-1}x + sin^{-1}2x = \frac{2\pi}{3}$$

D-3 Solve the following equations:

(i)
$$tan^{-1}\left(\frac{1-x}{1+x}\right) = \frac{1}{2}tan^{-1}x, (x>0)$$

(ii)
$$3tan^{-1}\left(\frac{1}{2+\sqrt{3}}\right) - tan^{-1}\left(\frac{1}{x}\right) = tan^{-1}\left(\frac{1}{3}\right)$$