

INSTITUTO ALEXANDER FLEMING (IAF)

Flow Cytometry Laboratory

Biochemist in charge: Dra. Andrea Novoa

Crámer 1180 | Phone: (+54-11) 3221-8900, ext. 8029

citometria@alexanderfleming.org

[Translation from Spanish into English — for informational reference]

Name: ADER, MANUEL
Accession Number: A298740
Accession date: April 16, 2026
Ordering physician: [Blank]

IMMUNOPHENOTYPING

SAMPLE: Bone marrow.

METHODOLOGY: A panel of monoclonal antibodies was directly conjugated with fluorescein isothiocyanate (FITC), phycoerythrin (PE), peridinin chlorophyll protein complex (PerCP), PerCP-Cy5.5, allophycocyanin (APC), APC-H7 tandem, PE-Cy7 tandem, Horizon V450 and V500. Data acquisition and analysis were performed using a FACSCanto II flow cytometer with FACSDiva and Infinicyt software.

RESULTS: Based on membrane immunophenotype, relative cell size, and internal complexity, the following populations were identified: Total lymphocytes: 2.7%. Plasma cells: <0.1%. Monocytes: 3%. Neutrophils: 21.3%. Eosinophils: <0.1%. Basophils: <0.1%. Mast cells: <0.01%. Nucleated erythroid cells: 46.9%.

PATHOLOGICAL CELLS: 26.1% myeloid blasts of medium size. CD45+, CD34+, HLA-DR++, CD117+, CD38++, CD13++, CD16(-), CD11b(-), CD10(-), CD71-/+ , CD64(-), CD14(-), CD36(-), CD105(-), CD33+, CD15-/+ , CD56(-), CD7-/(42%).

OBSERVATIONS: Neutrophils show marked dyshematopoietic features in maturation patterns CD13 / CD16 / CD11b / CD10. Monocytes show a predominance of immature forms (85% of the total monocytic lineage) with partial CD13 expression. The erythroid lineage is increased, with heterogeneous expression of CD36 and CD71.

CONCLUSIONS: Presence of a myeloid blast population (26.1%) similar to those found in previous studies. A final diagnostic conclusion requires correlation with the clinical presentation and additional complementary studies.

[Illegible signature]

Dra. Paula Tempra

Biochemist

National License No. 9824

[Illegible signature]

Dra. Andrea Novoa

Biochemist

National License No. 8693