The Journal of Open Source Software

WetlandMapper: A Python package for automatic wetland mapping, dynamics classification, and cover-type characterisation from multispectral time-series data

Manudeo Singh [1¶]

1 Department of Geography and Earth Sciences, Aberystwyth University, Aberystwyth, Wales, UK ¶ Corresponding author

DOI: 10.xxxxxx/draft

Software

- Review
- Repository
- Archive

Editor: Open Journals

Reviewers:

- @openjournals

Submitted: 01 January 1970
Published: unpublished

License
Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License (CC BY 4.0).

DRAFT

Summary

`WetlandMapper` is an open-source Python package that operationalises two peer-reviewed remote-sensing frameworks for automated wetland analysis from multispectral satellite data, now expanded with comprehensive spectral index computation, terrain analysis, and enhanced visualisation capabilities. The package is available on PyPI (`pip install wetlandmapper`) and archived at Zenodo (Singh, 2026).

The first framework is a **wetland dynamics classification** method (Singh & Sinha, 2022a) that maps floodplain wetlands at basin scale from a multi-temporal MNDWI time series and classifies each wetland pixel into one of six temporal dynamics classes: *Persistent*, *New*, *Intensifying*, *Diminishing*, *Lost*, and *Intermittent*.

The second framework is a **Wetland Cover Type (WCT) classification** method (Singh et al., 2022) that combines multiple spectral indices to characterise the biophysical surface composition of wetland pixels into stable, ecologically interpretable cover types including open clear water, turbid water, aquatic vegetation, and moist soil.

The package now includes comprehensive **spectral index computation** supporting seven indices (MNDWI, NDWI, NDVI, NDTI, AWEIsh, AWEInsh) for flexible water detection and vegetation analysis, a **terrain analysis module** for topographic corrections and artifact masking, and **visualisation utilities** for interactive plotting of results.

These modules address complementary monitoring questions: the dynamics module characterises *when and how* wetland inundation is changing over time, the WCT module characterises *what* is present at the wetland surface at any given time, the spectral indices provide flexible water and vegetation detection, and the terrain module enables topographic corrections for improved accuracy in hilly terrain. All operate on `xarray` `DataArray` objects (Hoyer & Hamman, 2017), enabling dask-backed parallel processing of large raster archives. Users may supply their own pre-processed imagery or retrieve analysis-ready surface-reflectance data directly from Google Earth Engine (GEE; Gorelick et al. (2017)) via an integrated optional acquisition submodule that supports all Landsat missions (4, 5, 7, 8, and 9), Sentinel-2, MODIS, and shapefile-path or GeoJSON area-of-interest inputs.

Statement of Need

Wetlands cover roughly 5–8percent of Earth's land surface and deliver ecosystem services valued at tens of trillions of dollars annually, yet global wetland area has declined by approximately

35percent since 1970 (Davidson, 2014; Ramsar Convention Secretariat, 2018). Operational monitoring at basin to continental scales — tracking both inundation dynamics and biophysical surface conditions — is essential for conservation management, restoration prioritisation, and international reporting obligations under the Ramsar Convention.

Existing tools for remote sensing-based wetland analysis suffer from one or more of the following limitations: (1) they characterise only binary inundation extent and do not resolve temporal dynamics or surface cover composition (Pekel et al., 2016); (2) they are embedded in proprietary platforms (ArcGIS Model Builder, GEE JavaScript API) that resist integration into scripted, reproducible workflows; (3) they address either dynamics or cover characterisation but not both within a single interoperable framework; (4) their methods are distributed as single-use scripts rather than tested, documented software libraries with version histories; (5) they lack comprehensive spectral index libraries for flexible water and vegetation detection; and (6) they do not account for topographic effects that can confound wetland classification in hilly terrain.

`WetlandMapper` addresses all six gaps. It provides a fully Pythonic, open-source library that unifies the dynamics-classification method of Singh & Sinha (2021) and Singh & Sinha (2022a) — previously dependent on ArcGIS — and the WCT method of Singh et al. (2022) — previously distributed only as GEE JavaScript code and an ArcGIS toolbox — while adding comprehensive spectral index computation, terrain analysis for topographic corrections, visualisation utilities, and a flexible data-ingestion pathway that supports both user-supplied imagery and automated GEE retrieval. Both core methods require no labelled training data and operate on any multispectral archive from which the required indices can be computed.

State of the Field

The JRC Global Surface Water dataset (Pekel et al., 2016) characterises long-term open-water occurrence globally at 30 m resolution but does not distinguish wetland surface types. Machine-learning approaches achieve high classification accuracy but require labelled training data that are rarely available at regional scales (Mahdavi et al., 2018; Slagter et al., 2020). SAR-based methods handle cloud cover but require specialist pre-processing workflows (**?**) and site-specific thresholding. Most tools lack comprehensive spectral index libraries or terrain correction capabilities needed for robust wetland analysis across diverse landscapes.

The methods unified in `WetlandMapper` occupy a practical middle ground — no training labels, applicable to any cloud-free multispectral archive, and producing ecologically interpretable outputs — while providing comprehensive spectral index computation, terrain analysis for topographic corrections, and visualisation utilities. `WetlandMapper` makes these capabilities accessible in a reusable, tested Python library for the first time.

Software Design and Methods

Wetland Dynamics Classification

This module implements the basin-scale inventory and hydrodynamics framework of Singh & Sinha (2022a). Given a multi-temporal MNDWI raster stack, each time step is first thresholded to a binary water-presence layer:

$$W_t = \begin{cases} 1 & \text{if } \mathrm{MNDWI}_t > \tau \\ 0 & \text{otherwise} \end{cases}$$

where $\tau = 0$ by default (positive MNDWI indicates a water-dominated pixel; Xu (2006)). Three summary statistics are then derived across the full time series of length T, split by a configurable temporal window n:

$$W_{\text{percent}} = \frac{\sum_{t=1}^{T} W_t}{T} \times 100, \quad W_{\text{historic}} = \sum_{t=1}^{n} W_t, \quad W_{\text{recent}} = \sum_{t=T-n+1}^{T} W_t$$

The temporal change signal $\Delta W = W_{\text{recent}} - W_{\text{historic}}$ is used together with W_{percent} and two user-adjustable wet-frequency thresholds (θ_{wet}, θ_{persis}) to assign each pixel to one of six dynamics classes:

Class	Primary condition
Persistent	$W_{\text{percent}} \geq \theta_{\text{persis}}$
New	$\Delta W = +n$
Intensifying	$W_{\text{percent}} \geq \theta_{\text{wet}};\ 0 < \Delta W < n$
Diminishing	$W_{\text{percent}} \geq \theta_{\text{wet}};\ -n < \Delta W < 0$
Lost	$\Delta W = -n$
Intermittent	$W_{\text{percent}} \geq \theta_{\text{wet}}$; no directional trend
Non-wetland	$W_{\text{percent}} < \theta_{\text{wet}}$

The entire classification is implemented using vectorised `xr.where` operations, enabling chunked parallel execution via Dask with no Python-level pixel iteration.

Wetland Cover Type Classification

This module implements the multi-index WCT framework of Singh et al. (2022). Three spectral indices are computed from the same multispectral image:

$$\text{MNDWI} = \frac{G - \text{SWIR}}{G + \text{SWIR}}, \quad \text{NDVI} = \frac{\text{NIR} - R}{\text{NIR} + R}, \quad \text{NDTI} = \frac{R - G}{R + G}$$

MNDWI delineates the extent of surface water; NDVI quantifies vegetation presence and density; NDTI quantifies water turbidity. Their combined spectral signatures partition wetland pixels into five biophysically distinct cover types. Two classification implementations are provided: the original quartile-based combination code method (`classify_wct_ema`), and an improved continuous-threshold variant (`classify_wct`) that allows sub-quartile calibration for different sensors or seasons.

Spectral Index Computation

The package provides comprehensive spectral index computation supporting seven indices for flexible water detection and vegetation analysis:

- **MNDWI**: Modified Normalised Difference Water Index (Xu, 2006)
- **NDWI**: Normalised Difference Water Index (McFeeters, 1996)
- **NDVI**: Normalised Difference Vegetation Index
- **NDTI**: Normalised Difference Turbidity Index
- **AWEIsh**: Automated Water Extraction Index with shadow suppression (Feyisa et al., 2014)
- **AWEInsh**: Automated Water Extraction Index without shadow suppression (Feyisa et al., 2014)

Users can compute individual indices or use `compute_indices()` for the core WCT indices (MNDWI, NDVI, NDTI) or `compute_water_indices()` for comprehensive water detection across all available water indices.

DRAFT

Terrain Analysis

The terrain analysis module provides topographic corrections essential for accurate wetland classification in hilly or mountainous terrain where slope and topographic position can confound spectral signatures. Functions include:

- **Slope computation**: Local slope calculation from digital elevation models
- **Topographic Position Index (TPI)**: Relative topographic position within a local neighborhood
- **Local range**: Local elevation variability for terrain complexity assessment
- **Terrain artifact masking**: Automated masking of steep slopes that may cause classification errors in wetland detection

Data Acquisition and Temporal Aggregation

Data Acquisition and Temporal Aggregation

The optional `wetlandmapper.gee` submodule supports all five Landsat missions (4, 5, 7, 8, 9), Sentinel-2, and MODIS. A `"LandsatAll"` option automatically merges available missions for any requested date range with harmonised band names, enabling long-record analyses from 1982 to the present day. `"MODISAll"` provides similar functionality for MODIS Terra and Aqua missions. An optional `use_slc_off` parameter controls whether Landsat 7 images acquired after the 2003 Scan Line Corrector failure (which cause ~22percent data gaps per scene) are included. Areas of interest may be provided as a GeoJSON dict, a shapefile path, or a GeoJSON file path; multi-feature shapefiles are dissolved to a single boundary automatically.

Server-side temporal compositing reduces data transfer volume for long time series: users may request one median composite per year, month, or meteorological season (DJF / MAM / JJA / SON) rather than every individual scene. The same temporal aggregation functionality is also available client-side via `aggregate_time()`, which operates on any `xarray` `DataArray` or `Dataset` regardless of data source.

Dependencies and Installation

`WetlandMapper` requires Python $\geq$ 3.9. Core dependencies are `numpy` (Harris et al., 2020), `xarray` (Hoyer & Hamman, 2017), and `rioxarray` (Snow & others, 2022). The GEE submodule additionally requires `earthengine-api`, `rasterio`, `xee`, `dask`, and `geopandas`. The plotting submodule requires `matplotlib` (Hunter, 2007). Installation:

```
pip install wetlandmapper                # core
pip install "wetlandmapper[gee]"         # with GEE + shapefile support
pip install "wetlandmapper[plot]"        # with visualisation utilities
pip install "wetlandmapper[all]"         # complete installation
```

Detailed platform-specific installation instructions, including GEE authentication setup, are provided in `INSTALL.md` in the repository.

Usage

A minimal end-to-end example for each workflow:

```python
import xarray as xr
from wetlandmapper import compute_mndwi, classify_dynamics
from wetlandmapper import compute_indices, classify_wct_ema
from wetlandmapper import compute_water_indices, compute_slope
from wetlandmapper.plotting import plot_dynamics, plot_wct
from wetlandmapper.gee import fetch
```

DRAFT

```python
# --- Dynamics: fetch annual composites from all Landsat missions ---
mndwi = fetch("study_area.shp", "1984-01-01", "2023-12-31",
              sensor="LandsatAll", temporal_aggregation="annual")
dynamics = classify_dynamics(mndwi, nYear=3,
                             thresholdWet=25, thresholdPersis=75)
dynamics.rio.to_raster("wetland_dynamics.tif")

# --- WCT: single composite → 5 biophysical cover types
# --- ds_composite is an xarray multispectral dataset (a Landsat .tiff)
ds_composite = xr.load_dataset('Landsat.tiff')
indices = compute_indices(ds_composite, green_band="B3", red_band="B4",
                          nir_band="B5", swir_band="B6")
wct = classify_wct_ema(indices)
wct.rio.to_raster("wetland_cover_types.tif")

# --- Comprehensive water detection with all available indices ---
water_indices = compute_water_indices(ds_composite,
                                      blue_band="B2", green_band="B3",
                                      red_band="B4", nir_band="B5",
                                      swir_band="B6", swir2_band="B7")
water_indices  # Contains MNDWI, NDWI, AWEIsh, AWEInsh

# --- Terrain analysis for topographic corrections ---
dem = xr.open_dataset("elevation.nc")["elevation"]
slope = compute_slope(dem)
# Apply terrain masking to reduce false positives in steep areas

# --- Visualisation ---
plot_dynamics(dynamics, title="Wetland Dynamics Classification")
plot_wct(wct, title="Wetland Cover Types")
```

A Jupyter notebook demonstrating both workflows on synthetic data, with full GEE acquisition, terrain analysis, and interactive visualisation sections, is included at

That study demonstrated that frequency-based temporal aggregation can effectively distinguish permanently inundated, seasonally active, and recently changed wetlands at basin scale, providing a validated inventory framework applicable to wetland restoration prioritisation (Singh & Sinha, 2022b).

The WCT module is derived from Singh et al. (2022), where the MNDWI–NDVI–NDTI combination was validated across three Ramsar-listed wetlands in contrasting geomorphic and climatic settings — Kaabar Tal (Ganga floodplain), Chilika Lagoon (coastal), and Nal Sarovar (semi-arid) — demonstrating that the WCTs are stable in space and time, meaning comparable biophysical conditions correspond to the same WCT irrespective of geographic location.

By unifying both workflows in a single, tested Python package with flexible data ingestion, `WetlandMapper` enables application of these frameworks globally in fully scripted, reproducible workflows, without dependence on proprietary GEE or ArcGIS environments.

Acknowledgements

The author acknowledges the ISRO–IIT Kanpur Space Technology Cell and WWF-India for funding the original research underlying both methods, and the Google Earth Engine team for providing cloud-computing access. The author is a Newton International Fellow funded by The Royal Society, London, and gratefully acknowledges this support. The facilities and support provided by Aberystwyth University are also duly acknowledged.

AI Assistance

Development of this software package was assisted by Claude (Anthropic), an AI language model, for tasks including code scaffolding, packaging configuration, continuous integration setup, docstring writing, and debugging. The scientific methodology, classification algorithms, spectral index thresholds, and validation are based entirely on the author's prior peer-reviewed work (Singh et al., 2022; Singh & Sinha, 2021, 2022a), and all scientific content, design decisions, and results presented here are the author's own.

References

Davidson, N. C. (2014). How much wetland has the world lost? Long-term and recent trends in global wetland area. *Marine and Freshwater Research*, *65*(10), 934–941. https://doi.org/10.1071/MF14173

Feyisa, G. L., Meilby, H., Fensholt, R., & Proud, S. R. (2014). Automated water extraction index: AWE for surface water mapping AW3D. *Remote Sensing of Environment*, *140*, 23–35. https://doi.org/10.1016/j.rse.2013.08.029

Gorelick, N., Hancher, M., Dixon, M., Ilyushchenko, S., Thau, D., & Moore, R. (2017). Google Earth Engine: Planetary-scale geospatial analysis for everyone. *Remote Sensing of Environment*, *202*, 18–27. https://doi.org/10.1016/j.rse.2017.06.031

Harris, C. R., Millman, K. J., Walt, S. J. van der, & others. (2020). Array programming with NumPy. *Nature*, *585*(7825), 357–362. https://doi.org/10.1038/s41586-020-2649-2

Hoyer, S., & Hamman, J. J. (2017). Xarray: N-D labeled arrays and datasets in Python. *Journal of Open Research Software*, *5*(1), 10. https://doi.org/10.5334/jors.148

Hunter, J. D. (2007). Matplotlib: A 2D graphics environment. *Computing in Science & Engineering*, *9*(3), 90–95. https://doi.org/10.1109/MCSE.2007.55

Mahdavi, S., Salehi, B., Granger, J., Amani, M., Brisco, B., & Huang, W. (2018). Remote sensing for wetland classification: A comprehensive review. *GIScience & Remote Sensing*, *55*(5), 623–658. https://doi.org/10.1080/15481603.2017.1419602

McFeeters, S. K. (1996). The use of the Normalized Difference Water Index (NDWI) in the delineation of open water features. *International Journal of Remote Sensing*, *17*(7), 1425–1432. https://doi.org/10.1080/01431169608948714

Pekel, J.-F., Cottam, A., Gorelick, N., & Belward, A. S. (2016). High-resolution mapping of global surface water and its long-term changes. *Nature*, *540*(7633), 418–422. https://doi.org/10.1038/nature20584

Ramsar Convention Secretariat. (2018). *Global wetland outlook: State of the world's wetlands and their services to people*. Ramsar Convention Secretariat. https://www.global-wetland-outlook.ramsar.org

Singh, M. (2026). *WetlandMapper: Automatic wetland mapping, dynamics classification, and cover-type characterisation from multispectral time-series data* (Version v1.0.0). Zenodo. https://doi.org/10.5281/zenodo.18967177

Singh, M., Allaka, S., Gupta, P. K., Patel, J. G., & Sinha, R. (2022). Deriving wetland-cover types (WCTs) from integration of multispectral indices based on Earth observation data. *Environmental Monitoring and Assessment*, *194*(12), 878. https://doi.org/10.1007/s10661-022-10541-7

Singh, M., & Sinha, R. (2021). Hydrogeomorphic indicators of wetland health inferred from multi-temporal remote sensing data for a new ramsar site (kaabar tal), india. *Ecological Indicators*, *127*(1), 107739. https://doi.org/10.1016/j.ecolind.2021.107739

DRAFT

Singh, M., & Sinha, R. (2022a). A basin-scale inventory and hydrodynamics of floodplain wetlands based on time-series of remote sensing data. *Remote Sensing Letters*, *13*(1), 1–13. https://doi.org/10.1080/2150704X.2021.1980919

Singh, M., & Sinha, R. (2022b). Integrating hydrological connectivity in a process–response framework for restoration and monitoring prioritisation of floodplain wetlands in the ramganga basin, india. *Water*, *14*(21), 3520. https://doi.org/10.3390/w14213520

Slagter, B., Tsendbazar, N.-E., Vollrath, A., & Reiche, J. (2020). Mapping wetland characteristics using temporally dense Sentinel-1 and Sentinel-2 data: A case study in the St. Lucia wetlands, South Africa. *International Journal of Applied Earth Observation and Geoinformation*, *86*, 102009. https://doi.org/10.1016/j.jag.2019.102009

Snow, A. D., & others. (2022). *Rioxarray: Geospatial xarray extension powered by rasterio*. Zenodo. https://doi.org/10.5281/zenodo.4570456

Xu, H. (2006). Modification of normalised difference water index (NDWI) to enhance open water features in remotely sensed imagery. *International Journal of Remote Sensing*, *27*(14), 3025–3033. https://doi.org/10.1080/01431160600589179

DRAFT