

AMCS 394E: Contemp. Topics in Computational Science.

Computing with the finite element method

David I. Ketcheson and Manuel Quezada de Luna



Strong and weak forms of a PDE

Given a PDE, for example

$$-\frac{d^2 u(x)}{dx^2} = f(x), \quad \forall x \in \Omega = (0, 1),$$

how can we find an approximation of its solution?

Strong and weak forms of a PDE

From the previous lectures:

Strong and weak forms of a PDE

From the previous lectures:

- * The solution $u(x)$ belongs to some infinite dimensional space \tilde{V} .

Strong and weak forms of a PDE

From the previous lectures:

- * The solution $u(x)$ belongs to some infinite dimensional space \tilde{V} .
- * We want to consider

$$V_h = \{v \in C^0(\Omega) \mid v|_K \in \mathbb{P}^p(K)\}$$

Strong and weak forms of a PDE

From the previous lectures:

- * The solution $u(x)$ belongs to some infinite dimensional space \tilde{V} .
- * We want to consider

$$V_h = \{v \in C^0(\Omega) \mid v|_K \in \mathbb{P}^p(K)\}$$

- * We look for a solution $u_h(x) \in V_h$; i.e., we approximate $u(x)$ via continuous piecewise polynomials.


Strong and weak forms of a PDE

From the previous lectures:

- * The solution $u(x)$ belongs to some infinite dimensional space \tilde{V} .
- * We want to consider

$$V_h = \{v \in C^0(\Omega) \mid v|_K \in \mathbb{P}^p(K)\}$$

- * We look for a solution $u_h(x) \in V_h$; i.e., we approximate $u(x)$ via continuous piecewise polynomials.



Strong and weak forms of a PDE

From the previous lectures:

Strong and weak forms of a PDE

From the previous lectures:

- * We have a basis $B = \{\varphi_i\}_{i=0}^{\dim(V_h)}$ for V_h .

Strong and weak forms of a PDE

From the previous lectures:

- * We have a basis $B = \{\varphi_i\}_{i=0}^{\dim(V_h)}$ for V_h .
- * We assume for now the basis is interpolatory: $\varphi_i(x_j) = \delta_{ij}$.

Strong and weak forms of a PDE

From the previous lectures:

- * We have a basis $B = \{\varphi_i\}_{i=0}^{\dim(V_h)}$ for V_h .
- * We assume for now the basis is interpolatory: $\varphi_i(x_j) = \delta_{ij}$.
- * We expand the solution $u_h(x)$ as follows:

$$u_h(x) = \sum_j U_j \varphi_j(x),$$

where (for this interpolatory basis) $U_j = u_h(x_j)$.

Strong and weak forms of a PDE

From the previous lectures:

- * We have a basis $B = \{\varphi_i\}_{i=0}^{\dim(V_h)}$ for V_h .
- * We assume for now the basis is interpolatory: $\varphi_i(x_j) = \delta_{ij}$.
- * We expand the solution $u_h(x)$ as follows:

$$u_h(x) = \sum_j U_j \varphi_j(x),$$

where (for this interpolatory basis) $U_j = u_h(x_j)$.

NOTE:

In order to represent the solution $u_h(x)$, we only need the (finitely many) coordinates U_j . These coordinates are indeed the solution of our linear algebraic systems.

Strong and weak forms of a PDE

The idea of a **weak formulation** is to consider a “less restrictive” version of the PDE.

Strong and weak forms of a PDE

The idea of a **weak formulation** is to consider a “less restrictive” version of the PDE.

Say I want to compare two functions $f(x)$ and $g(x)$; i.e., are they ‘equal’ in some domain Ω ?

Strong and weak forms of a PDE

The idea of a **weak formulation** is to consider a “less restrictive” version of the PDE.

Say I want to compare two functions $f(x)$ and $g(x)$; i.e., are they ‘equal’ in some domain Ω ?

$$\text{If } f(x) = g(x), \quad \forall x \in \Omega$$

then clearly $f(x) = g(x)$.

Strong and weak forms of a PDE

The idea of a **weak formulation** is to consider a “less restrictive” version of the PDE.

Say I want to compare two functions $f(x)$ and $g(x)$; i.e., are they ‘equal’ in some domain Ω ?

$$\text{If } f(x) = g(x), \quad \forall x \in \Omega$$

then clearly $f(x) = g(x)$.

Problem:

This is too restrictive.

Strong and weak forms of a PDE

The idea of a **weak formulation** is to consider a “less restrictive” version of the PDE.

Say I want to compare two functions $f(x)$ and $g(x)$; i.e., are they ‘equal’ in some domain Ω ?

$$\text{If } f(x) = g(x), \quad \forall x \in \Omega$$

then clearly $f(x) = g(x)$.

Problem:

This is too restrictive.

Question:

How can we say $f(x) = g(x)$ in a weaker sense?

Strong and weak forms of a PDE

How can we say $f(x) = g(x)$ in a weaker sense (than pointwise)?

Strong and weak forms of a PDE

How can we say $f(x) = g(x)$ in a weaker sense (than pointwise)?

We can weaken the question by asking if applying a functional to both equations yields the same number. For instance, does

$$\int_{\Omega} f(x) dx = \int_{\Omega} g(x) dx, \quad \text{hold?}$$

Strong and weak forms of a PDE

How can we say $f(x) = g(x)$ in a weaker sense (than pointwise)?

We can weaken the question by asking if applying a functional to both equations yields the same number. For instance, does

$$\int_{\Omega} f(x) dx = \int_{\Omega} g(x) dx, \quad \text{hold?}$$

Problem:

This is too weak. For instance,

$$1, \quad 2x, \quad 3x^2, \quad \dots, \quad (n+1)x^n$$

have the same integral over $\Omega := (0, 1)$.

Strong and weak forms of a PDE

How can we say $f(x) = g(x)$ in a weaker sense (than pointwise)?

We can define that $f(x) = g(x)$ if

$$\int_{\Omega} f(x) dx = \int_{\Omega} g(x) dx \quad \text{and} \quad \int_{\Omega} x f(x) dx = \int_{\Omega} x g(x) dx,$$

and/or consider more functionals.

Strong and weak forms of a PDE

How can we say $f(x) = g(x)$ in a weaker sense (than pointwise)?

We can define that $f(x) = g(x)$ if

$$\int_{\Omega} f(x) dx = \int_{\Omega} g(x) dx \quad \text{and} \quad \int_{\Omega} x f(x) dx = \int_{\Omega} x g(x) dx,$$

and/or consider more functionals.

Question:

For a given (vector) space V , what is a good set of functionals to weakly define that $f(x) = g(x)$?

Strong and weak forms of a PDE

Given a vector space V with some inner product $\langle \cdot, \cdot \rangle$,
 $f, g \in V$ are weakly equal if

$$\langle f - g, v \rangle = 0, \quad \forall v \in V$$

Strong and weak forms of a PDE

Given a vector space V with some inner product $\langle \cdot, \cdot \rangle$,
 $f, g \in V$ are weakly equal if

$$\langle f - g, v \rangle = 0, \quad \forall v \in V$$

Example

If $V = L^2(\Omega)$, f and g are weakly equal if

$$\int_{\Omega} (f - g)v \, dx = 0, \quad \forall v \in V$$

Strong and weak forms of a PDE

Given a vector space V with some inner product $\langle \cdot, \cdot \rangle$,
 $f, g \in V$ are weakly equal if

$$\langle f - g, v \rangle = 0, \quad \forall v \in V$$

Example

If $V = L^2(\Omega)$, f and g are weakly equal if

$$\int_{\Omega} (f - g)v \, dx = 0, \quad \forall v \in V$$

Note

If $B = \{\varphi\}_i$ is a basis for V , then $f = g$ if

$$\int_{\Omega} (f - g)\varphi_i \, dx = 0, \quad \forall \varphi_i \in B$$

Strong and weak forms of a PDE

The **strong form** of a PDE is the differential form; e.g.,

$$-\frac{d^2 u(x)}{dx^2} = f(x), \quad \forall x \in \Omega = (0, 1)$$

Strong and weak forms of a PDE

The **strong form** of a PDE is the differential form; e.g.,

$$-\frac{d^2 u(x)}{dx^2} = f(x), \quad \forall x \in \Omega = (0, 1)$$

The **weak form** of a PDE holds in a weak sense; e.g.,

$$-\int_0^1 \frac{d^2 u(x)}{dx^2} v dx = \int_0^1 f(x) v dx, \quad \forall v \in V$$

Strong and weak forms of a PDE

The **strong form** of a PDE is the differential form; e.g.,

$$-\frac{d^2 u(x)}{dx^2} = f(x), \quad \forall x \in \Omega = (0, 1)$$

The **weak form** of a PDE holds in a weak sense; e.g.,

$$-\int_0^1 \frac{d^2 u(x)}{dx^2} v dx = \int_0^1 f(x) v dx, \quad \forall v \in V$$

or, integrating by parts:

$$\int_0^1 \frac{du}{dx} \frac{dv}{dx} dx + BCs = \int_0^1 f(x) v dx, \quad \forall v \in V$$

Strong and weak forms of a PDE

Why and when should we integrate by parts?

Strong and weak forms of a PDE

Why and when should we integrate by parts?

- * If we expect the solution to be not “sufficiently” smooth.

Strong and weak forms of a PDE

Why and when should we integrate by parts?

- * If we expect the solution to be not “sufficiently” smooth.
- * If we need to impose boundary conditions via a numerical flux.

Strong and weak forms of a PDE

Why and when should we integrate by parts?

- * If we expect the solution to be not “sufficiently” smooth.
- * If we need to impose boundary conditions via a numerical flux.
- * We need to consider the finite dimensional space we aim to use.

Strong and weak forms of a PDE

Examples of strong to weak forms

Finite dimensional approximation of the weak form

Strong formulation:

Consider the following PDE in strong form

$$-\frac{d^2 u}{dx^2} = f(x), \quad \forall x \in \Omega = (0, 1),$$

with

$$du/dx|_{x=1} = g_r, \quad \text{and} \quad du/dx|_{x=0} = g_l,$$

as boundary conditions.

Finite dimensional approximation of the weak form

Strong formulation:

Consider the following PDE in strong form

$$-\frac{d^2 u}{dx^2} = f(x), \quad \forall x \in \Omega = (0, 1),$$

with

$$du/dx|_{x=1} = g_r, \quad \text{and} \quad du/dx|_{x=0} = g_l,$$

as boundary conditions.

Weak formulation:

Finite dimensional approximation of the weak form

Strong formulation:

Consider the following PDE in strong form

$$-\frac{d^2 u}{dx^2} = f(x), \quad \forall x \in \Omega = (0, 1),$$

with

$$du/dx|_{x=1} = g_r, \quad \text{and} \quad du/dx|_{x=0} = g_l,$$

as boundary conditions.

Weak formulation:

- * Get the weak form of the PDE and apply the boundary conditions.

Finite dimensional approximation of the weak form

Strong formulation:

Consider the following PDE in strong form

$$-\frac{d^2 u}{dx^2} = f(x), \quad \forall x \in \Omega = (0, 1),$$

with

$$du/dx|_{x=1} = g_r, \quad \text{and} \quad du/dx|_{x=0} = g_l,$$

as boundary conditions.

Weak formulation:

- * Get the weak form of the PDE and apply the boundary conditions.
- * What is an appropriate space for the solution?

Finite dimensional approximation of the weak form

Weak formulation:

$$\int_0^1 \frac{du}{dx} \frac{d\varphi}{dx} dx - [g_r \varphi(1) - g_l \varphi(0)] = \int_0^1 f(x) \varphi dx, \quad \forall \varphi \in V$$

where $V = H^1(\Omega)$.

Finite dimensional approximation of the weak form

Weak formulation:

$$\int_0^1 \frac{du}{dx} \frac{d\varphi}{dx} dx - [g_r \varphi(1) - g_l \varphi(0)] = \int_0^1 f(x) \varphi dx, \quad \forall \varphi \in V$$

where $V = H^1(\Omega)$.

- * The weak solution $u \in V$ is an infinite dimensional solution.

Finite dimensional approximation of the weak form

Weak formulation:

$$\int_0^1 \frac{du}{dx} \frac{d\varphi}{dx} dx - [g_r \varphi(1) - g_l \varphi(0)] = \int_0^1 f(x) \varphi dx, \quad \forall \varphi \in V$$

where $V = H^1(\Omega)$.

- * The weak solution $u \in V$ is an infinite dimensional solution.
- * We need to approximate u via a finite dimensional space.

Finite dimensional approximation of the weak form

Weak formulation:

$$\int_0^1 \frac{du}{dx} \frac{d\varphi}{dx} dx - [g_r \varphi(1) - g_l \varphi(0)] = \int_0^1 f(x) \varphi dx, \quad \forall \varphi \in V$$

where $V = H^1(\Omega)$.

- * The weak solution $u \in V$ is an infinite dimensional solution.
- * We need to approximate u via a finite dimensional space.
- * We can use the space space of continuous piecewise polynomials

$$V_h = \{\varphi \in C^0(\Omega) \mid \varphi_K \in \mathbb{P}^p(K)\}$$

Finite dimensional approximation of the weak form

Finite dimensional weak formulation:

Find $u_h \in V_h$ such that

$$\int_0^1 \frac{d\mathbf{u}_h}{dx} \frac{d\varphi}{dx} dx - [g_r \varphi(1) - g_l \varphi(0)] = \int_0^1 f(x) \varphi dx, \quad \forall \varphi \in V_h$$

Finite dimensional approximation of the weak form

Finite dimensional weak formulation:

Find $u_h \in V_h$ such that

$$\underbrace{\int_0^1 \frac{du_h}{dx} \frac{d\varphi}{dx} dx - [g_r \varphi(1) - g_l \varphi(0)]}_{=: a(u_h, \varphi)} = \underbrace{\int_0^1 f(x) \varphi dx}_{=: F(\varphi)}, \quad \forall \varphi \in V_h$$

Finite dimensional approximation of the weak form

Finite dimensional weak formulation:

Find $u_h \in V_h$ such that

$$\underbrace{\int_0^1 \frac{du_h}{dx} \frac{d\varphi}{dx} dx - [g_r \varphi(1) - g_l \varphi(0)]}_{=: a(u_h, \varphi)} = \underbrace{\int_0^1 f(x) \varphi dx}_{=: F(\varphi)}, \quad \forall \varphi \in V_h$$

The finite element problem is then to find $u_h \in V_h$ such that

$$a(u_h, \varphi) = F(\varphi), \quad \forall \varphi \in V_h$$

Galerkin orthogonality

Let $u \in V$ and $u_h, \varphi \in V_h$ and consider

$$a(u, \varphi) - a(u_h, \varphi) = F(\phi) - F(\phi) = 0$$

Galerkin orthogonality

Let $u \in V$ and $u_h, \varphi \in V_h$ and consider

$$a(u, \varphi) - a(u_h, \varphi) = F(\phi) - F(\phi) = 0$$

Then, we get

$$a(\underbrace{u - u_h}_{=e_h}, \varphi) = 0, \quad \forall \varphi \in V_h,$$

which means that the error e_h is orthogonal to V_h .

Finite element projection

Given a general function $f(x)$. We want to represent $f(x)$ in V_h .
That is, find

$$f_h(x) \in V_h \quad \text{s.t.} \quad f_h(x) = f(x)$$

in a weak sense in V_h .

From finite dimensional weak form to a linear algebra problem

Given a finite dimensional weak formulation, we need to “translate” the problem to a (non)linear algebra problem.

From finite dimensional weak form to a linear algebra problem

Given a finite dimensional weak formulation, we need to “translate” the problem to a (non)linear algebra problem.

Steps:

From finite dimensional weak form to a linear algebra problem

Given a finite dimensional weak formulation, we need to “translate” the problem to a (non)linear algebra problem.

Steps:

- i. Consider the weak form $\forall v \in V_h$.

From finite dimensional weak form to a linear algebra problem

Given a finite dimensional weak formulation, we need to “translate” the problem to a (non)linear algebra problem.

Steps:

- i. Consider the weak form $\forall v \in V_h$.
- ii. Consider $v = \phi_i \in B = \{\phi_j\}_{j=1}^{\dim(V_h)}$.

From finite dimensional weak form to a linear algebra problem

Given a finite dimensional weak formulation, we need to “translate” the problem to a (non)linear algebra problem.

Steps:

- i. Consider the weak form $\forall v \in V_h$.
- ii. Consider $v = \phi_i \in B = \{\phi_j\}_{j=1}^{\dim(V_h)}$.
- iii. Plug $u_h = \sum_j U_j \phi_j$ into the finite dim. weak form.

From finite dimensional weak form to a linear algebra problem

Given a finite dimensional weak formulation, we need to “translate” the problem to a (non)linear algebra problem.

Steps:

- i. Consider the weak form $\forall v \in V_h$.
- ii. Consider $v = \phi_i \in B = \{\phi_j\}_{j=1}^{\dim(V_h)}$.
- iii. Plug $u_h = \sum_j U_j \phi_j$ into the finite dim. weak form.
- iv. Write down the linear algebra problem.

From finite dimensional weak form to a linear algebra problem

Given a finite dimensional weak formulation, we need to “translate” the problem to a (non)linear algebra problem.

Steps:

- i. Consider the weak form $\forall v \in V_h$.
- ii. Consider $v = \phi_i \in B = \{\phi_j\}_{j=1}^{\dim(V_h)}$.
- iii. Plug $u_h = \sum_j U_j \phi_j$ into the finite dim. weak form.
- iv. Write down the linear algebra problem.

Note:

Always test again ϕ_i and always expand u_h using ϕ_j , then i and j will represent rows and columns, respectively.

From finite dimensional weak form to a linear algebra problem

Given a finite dimensional weak formulation, we need to “translate” the problem to a (non)linear algebra problem.

Consider, for example, the finite element problem

$$\int_0^1 \frac{du_h}{dx} \frac{d\varphi}{dx} dx = \int_0^1 f(x)\varphi dx + [g_r\varphi(1) - g_l\varphi(0)] =: F(\phi)$$

From finite dimensional weak form to a linear algebra problem

Given a finite dimensional weak formulation, we need to “translate” the problem to a (non)linear algebra problem.

Consider, for example, the finite element problem

$$\int_0^1 \frac{du_h}{dx} \frac{d\varphi}{dx} dx = \int_0^1 f(x)\varphi dx + [g_r\varphi(1) - g_l\varphi(0)] =: F(\phi)$$

Given a basis $B = \{\varphi_i\}_{i=0}^{\dim(V_h)}$ for V_h . Choose φ to be φ_i .

From finite dimensional weak form to a linear algebra problem

Given a finite dimensional weak formulation, we need to “translate” the problem to a (non)linear algebra problem.

Consider, for example, the finite element problem

$$\int_0^1 \frac{du_h}{dx} \frac{d\varphi}{dx} dx = \int_0^1 f(x)\varphi dx + [g_r\varphi(1) - g_l\varphi(0)] =: F(\phi)$$

Given a basis $B = \{\varphi_i\}_{i=0}^{\dim(V_h)}$ for V_h . Choose φ to be φ_i .
Since $u_h \in V_h$, we have

$$\int_0^1 \frac{d}{dx} \left(\sum_j U_j \varphi_j \right) \frac{d\varphi_i}{dx} dx = \sum_j U_j \int_0^1 \frac{d\varphi_j}{dx} \frac{d\varphi_i}{dx} dx = F(\varphi_i)$$

From finite dimensional weak form to a linear algebra problem

Given a finite dimensional weak formulation, we need to “translate” the problem to a (non)linear algebra problem.

Consider, for example, the finite element problem

$$\int_0^1 \frac{du_h}{dx} \frac{d\varphi}{dx} dx = \int_0^1 f(x)\varphi dx + [g_r\varphi(1) - g_l\varphi(0)] =: F(\phi)$$

Given a basis $B = \{\varphi_i\}_{i=0}^{\dim(V_h)}$ for V_h . Choose φ to be φ_i .
Since $u_h \in V_h$, we have

$$\int_0^1 \frac{d}{dx} \left(\sum_j U_j \varphi_j \right) \frac{d\varphi_i}{dx} dx = \sum_j U_j \underbrace{\int_0^1 \frac{d\varphi_j}{dx} \frac{d\varphi_i}{dx} dx}_{=: S_{ij}} = \underbrace{F(\varphi_i)}_{=: F_i}$$

From finite dimensional weak form to a linear algebra problem

More examples:

Strong, weak and finite dimensional problems

Consider the following PDE in strong form:

$$u_t + \nabla \cdot (\mathbf{a}u) - \nabla \cdot \mu \nabla u = 0,$$

$$u = u_B,$$

$$\partial_{\mathbf{n}} u = f(x),$$

$$u(\mathbf{a} \cdot \mathbf{n}) - \mu \partial_{\mathbf{n}} u = g(x),$$

$$\forall x \in \Gamma_D,$$

$$\forall x \in \Gamma_N,$$

$$\forall x \in \Gamma_R$$

Strong, weak and finite dimensional problems

Consider the following PDE in strong form:

$$\begin{aligned}u_t + \nabla \cdot (\mathbf{a}u) - \nabla \cdot \mu \nabla u &= 0, \\u &= u_B, & \forall x \in \Gamma_D, \\ \partial_{\mathbf{n}} u &= f(x), & \forall x \in \Gamma_N, \\ u(\mathbf{a} \cdot \mathbf{n}) - \mu \partial_{\mathbf{n}} u &= g(x), & \forall x \in \Gamma_R\end{aligned}$$

The corresponding weak form is given by

$$\partial_t \int_{\Omega} u \varphi dx + \int_{\Omega} \nabla \cdot (\mathbf{a}u) \varphi dx - \int_{\Omega} \nabla \cdot (\mu \nabla u) \varphi dx = 0, \quad \forall \varphi \in V$$

Strong, weak and finite dimensional problems

Consider the following PDE in strong form:

$$\begin{aligned}u_t + \nabla \cdot (\mathbf{a}u) - \nabla \cdot \mu \nabla u &= 0, \\u &= u_B, & \forall \mathbf{x} \in \Gamma_D, \\ \partial_{\mathbf{n}} u &= f(x), & \forall \mathbf{x} \in \Gamma_N, \\ u(\mathbf{a} \cdot \mathbf{n}) - \mu \partial_{\mathbf{n}} u &= g(x), & \forall \mathbf{x} \in \Gamma_R\end{aligned}$$

The corresponding weak form is given by

$$\partial_t \int_{\Omega} u \varphi dx + \int_{\Omega} \nabla \cdot (\mathbf{a}u) \varphi dx - \int_{\Omega} \nabla \cdot (\mu \nabla u) \varphi dx = 0, \quad \forall \varphi \in V$$

$$\int_{\Omega} \nabla \cdot (\mathbf{a}u) \varphi dx = - \int_{\Omega} u(\mathbf{a} \cdot \nabla \varphi) dx + \int_{\partial\Omega} u(\mathbf{a} \cdot \mathbf{n}) \varphi ds$$

Strong, weak and finite dimensional problems

Consider the following PDE in strong form:

$$\begin{aligned}u_t + \nabla \cdot (\mathbf{a}u) - \nabla \cdot \mu \nabla u &= 0, \\u &= u_B, & \forall \mathbf{x} \in \Gamma_D, \\ \partial_{\mathbf{n}} u &= f(\mathbf{x}), & \forall \mathbf{x} \in \Gamma_N, \\ u(\mathbf{a} \cdot \mathbf{n}) - \mu \partial_{\mathbf{n}} u &= g(\mathbf{x}), & \forall \mathbf{x} \in \Gamma_R\end{aligned}$$

The corresponding weak form is given by

$$\begin{aligned}\partial_t \int_{\Omega} u \varphi d\mathbf{x} + \int_{\Omega} \nabla \cdot (\mathbf{a}u) \varphi d\mathbf{x} - \int_{\Omega} \nabla \cdot (\mu \nabla u) \varphi d\mathbf{x} &= 0, \quad \forall \varphi \in V \\ - \int_{\Omega} \nabla \cdot (\mu \nabla u) \varphi d\mathbf{x} &= \int_{\Omega} \mu \nabla u \cdot \nabla \varphi d\mathbf{x} - \int_{\partial\Omega} \mu (\nabla u \cdot \mathbf{n}) \varphi ds\end{aligned}$$

Strong, weak and finite dimensional problems

The weak form is given by

$$\partial_t \int_{\Omega} u \varphi dx - \int_{\Omega} u (\mathbf{a} \cdot \nabla \varphi) dx + \int_{\Omega} \mu \nabla u \cdot \nabla \varphi dx + B(u, \varphi) = 0, \quad \forall \varphi \in V$$

where $B(u, \varphi)$ is an operator that accounts for the boundary conditions.

Strong, weak and finite dimensional problems

The weak form is given by

$$\partial_t \int_{\Omega} u \varphi dx - \int_{\Omega} u (\mathbf{a} \cdot \nabla \varphi) dx + \int_{\Omega} \mu \nabla u \cdot \nabla \varphi dx + B(u, \varphi) = 0, \quad \forall \varphi \in V$$

where $B(u, \varphi)$ is an operator that accounts for the boundary conditions.

What is the space V ?

Strong, weak and finite dimensional problems

The weak form is given by

$$\partial_t \int_{\Omega} u \varphi dx - \int_{\Omega} u(\mathbf{a} \cdot \nabla \varphi) dx + \int_{\Omega} \mu \nabla u \cdot \nabla \varphi dx + B(u, \varphi) = 0, \quad \forall \varphi \in V$$

where $B(u, \varphi)$ is an operator that accounts for the boundary conditions.

What is the space V ?

$$V = \{\varphi \in H^1(\Omega) \mid \varphi = 0 \text{ in } \Gamma_D\}.$$

Strong, weak and finite dimensional problems

The weak form is given by

$$\partial_t \int_{\Omega} u \varphi dx - \int_{\Omega} u(\mathbf{a} \cdot \nabla \varphi) dx + \int_{\Omega} \mu \nabla u \cdot \nabla \varphi dx + B(u, \varphi) = 0, \quad \forall \varphi \in V$$

where $B(u, \varphi)$ is an operator that accounts for the boundary conditions.

The boundary conditions are imposed via

$$B(u, \varphi) = \int_{\partial\Omega} [u(\mathbf{a} \cdot \mathbf{n}) - \mu(\nabla u \cdot \mathbf{n})] \varphi ds$$

Strong, weak and finite dimensional problems

The weak form is given by

$$\partial_t \int_{\Omega} u \varphi dx - \int_{\Omega} u(\mathbf{a} \cdot \nabla \varphi) dx + \int_{\Omega} \mu \nabla u \cdot \nabla \varphi dx + B(u, \varphi) = 0, \quad \forall \varphi \in V$$

where $B(u, \varphi)$ is an operator that accounts for the boundary conditions.

The boundary conditions are imposed via

$$\begin{aligned} B(u, \varphi) &= \int_{\partial\Omega} [u(\mathbf{a} \cdot \mathbf{n}) - \mu(\nabla u \cdot \mathbf{n})] \varphi ds \\ &= \int_{\Gamma_N} [u(\mathbf{a} \cdot \mathbf{n}) - \mu f] \varphi ds + \int_{\Gamma_R} g \varphi ds \end{aligned}$$

Note that the integral in Γ_D vanishes since $\varphi = 0$ in Γ_D .

Strong, weak and finite dimensional problems

The **discrete** weak form is given by

$$\partial_t \int_{\Omega} \mathbf{u}_h \varphi dx - \int_{\Omega} \mathbf{u}_h (\mathbf{a} \cdot \nabla \varphi) dx + \int_{\Omega} \mu \nabla \mathbf{u}_h \cdot \nabla \varphi dx + B(\mathbf{u}_h, \varphi) = 0, \quad \forall \varphi \in V_h$$

Strong, weak and finite dimensional problems

To get the **linear algebra** problem, choose $\phi = \phi_i \in B$. Then we get

$$\partial_t \int_{\Omega} u_h \varphi_i dx - \int_{\Omega} u_h (\mathbf{a} \cdot \nabla \varphi_i) dx + \int_{\Omega} \mu \nabla u_h \cdot \nabla \varphi_i dx + B(u_h, \varphi_i) = 0$$

Strong, weak and finite dimensional problems

To get the **linear algebra** problem, choose $\phi = \phi_i \in B$. Then we get

$$\partial_t \int_{\Omega} u_h \phi_i dx - \int_{\Omega} u_h (\mathbf{a} \cdot \nabla \phi_i) dx + \int_{\Omega} \mu \nabla u_h \cdot \nabla \phi_i dx + B(u_h, \phi_i) = 0$$

Or in matrix form

$$\sum_j \dot{U}_j M_{ij} - \sum_j U_j K_{ij} + \sum_j U_j S_{ij} + BCs = 0,$$

where M_{ij} , K_{ij} and S_{ij} are the entries of the mass, transport and stiffness matrix.

Strong, weak and finite dimensional problems

During the implementation we need to consider the time integration method.

Integration by parts (IBP) in multiple dimensions

Divergence theorem:

$$\int_{\Omega} \nabla \cdot \mathbf{a} dx = \int_{\partial\Omega} \mathbf{a} \cdot \mathbf{n} dx$$

Integration by parts (IBP) in multiple dimensions

Divergence theorem:

$$\int_{\Omega} \nabla \cdot \mathbf{a} dx = \int_{\partial\Omega} \mathbf{a} \cdot \mathbf{n} dx$$

IBP based on the divergence:

$$\int_{\Omega} (\nabla \cdot \mathbf{a}) \varphi dx = - \int_{\Omega} \mathbf{a} \cdot \nabla \varphi dx + \int_{\partial\Omega} (\mathbf{a} \cdot \mathbf{n}) \varphi dx$$