

AMCS 394E: Contemp. Topics in Computational Science. Computing with the finite element method

David I. Ketcheson and Manuel Quezada de Luna



What have we seen so far?

Approximation theory:

What have we seen so far?

Approximation theory:

- * How do the solutions of PDEs look?

What have we seen so far?

Approximation theory:

- * How do the solutions of PDEs look?
- * How can we approximate these solutions?

What have we seen so far?

Approximation theory:

- * How do the solutions of PDEs look?
- * How can we approximate these solutions?
- * What is the error we make in this approximation?

What have we seen so far?

Approximation theory:

- * How do the solutions of PDEs look?
- * How can we approximate these solutions?
- * What is the error we make in this approximation?

Approximation of PDEs:

What have we seen so far?

Approximation theory:

- * How do the solutions of PDEs look?
- * How can we approximate these solutions?
- * What is the error we make in this approximation?

Approximation of PDEs:

- * Why do we go from a strong to a weak formulation?

What have we seen so far?

Approximation theory:

- * How do the solutions of PDEs look?
- * How can we approximate these solutions?
- * What is the error we make in this approximation?

Approximation of PDEs:

- * Why do we go from a strong to a weak formulation?
- * How do we get a finite dimensional approx. of the weak form?

What have we seen so far?

Approximation theory:

- * How do the solutions of PDEs look?
- * How can we approximate these solutions?
- * What is the error we make in this approximation?

Approximation of PDEs:

- * Why do we go from a strong to a weak formulation?
- * How do we get a finite dimensional approx. of the weak form?
- * How do we go from a finite dimensional approx. of the weak form to a linear algebra problem?

What have we seen so far?

Example:

Consider the following equation:

$$\begin{aligned}u - \Delta u &= 0, & \forall x \in \Omega, \\u &= u_D, & \forall x \in \Gamma_D, \\\partial_{\mathbf{n}} u &= b(x), & \forall x \in \Gamma_N.\end{aligned}$$

What have we seen so far?

Example:

Consider the following equation:

$$\begin{aligned}u - \Delta u &= 0, & \forall x \in \Omega, \\u &= u_D, & \forall x \in \Gamma_D, \\\partial_{\mathbf{n}} u &= b(x), & \forall x \in \Gamma_N.\end{aligned}$$

What is the weak formulation?

$$\int_{\Omega} u \varphi dx + \int_{\Omega} \nabla u \cdot \nabla \varphi dx - \int_{\partial\Omega} (\nabla u \cdot \mathbf{n}) \varphi ds = 0, \quad \forall \varphi \in V$$

What have we seen so far?

Example:

Consider the following equation:

$$\begin{aligned}u - \Delta u &= 0, & \forall x \in \Omega, \\u &= u_D, & \forall x \in \Gamma_D, \\\partial_{\mathbf{n}} u &= b(x), & \forall x \in \Gamma_N.\end{aligned}$$

What is the weak formulation?

$$\int_{\Omega} u \varphi dx + \int_{\Omega} \nabla u \cdot \nabla \varphi dx - \int_{\partial\Omega} b \varphi ds = 0, \quad \forall \varphi \in V$$

What have we seen so far?

Example:

Consider the following equation:

$$\begin{aligned}u - \Delta u &= 0, & \forall x \in \Omega, \\u &= u_D, & \forall x \in \Gamma_D, \\\partial_{\mathbf{n}} u &= b(x), & \forall x \in \Gamma_N.\end{aligned}$$

What is the weak formulation?

$$\int_{\Omega} u \varphi dx + \int_{\Omega} \nabla u \cdot \nabla \varphi dx - \int_{\partial\Omega} b \varphi ds = 0, \quad \forall \varphi \in V$$

where

$$V = \{v \in H^1(\Omega) \mid v = 0 \text{ if } x \in \Gamma_D\}.$$

What have we seen so far?

What is the discrete weak formulation?

$$\int_{\Omega} u_h \varphi dx + \int_{\Omega} \nabla u_h \cdot \nabla \varphi dx = \int_{\partial\Omega} b \varphi ds, \quad \forall \varphi \in V_h$$

What have we seen so far?

What is the discrete weak formulation?

$$\int_{\Omega} u_h \varphi dx + \int_{\Omega} \nabla u_h \cdot \nabla \varphi dx = \int_{\partial\Omega} b \varphi ds, \quad \forall \varphi \in V_h$$

where

$$V_h = \{v \in C^0(\Omega) \mid v_K \in \mathbb{P}^p\}.$$

What have we seen so far?

What is the discrete weak formulation?

$$\int_{\Omega} u_h \varphi dx + \int_{\Omega} \nabla u_h \cdot \nabla \varphi dx = \int_{\partial\Omega} b \varphi ds, \quad \forall \varphi \in V_h$$

where

$$V_h = \{v \in C^0(\Omega) \mid v_K \in \mathbb{P}^p\}.$$

How do we get a linear algebra problem?

Consider and plug $\phi = \phi_i$ and $u_h = \sum_j U_j \varphi_j$ into the equation.

$$\sum_j U_j \int_{\Omega} \varphi_i \varphi_j dx + \sum_j U_j \int_{\Omega} \nabla \varphi_i \cdot \nabla \varphi_j dx = \int_{\partial\Omega} b \varphi_i ds.$$

What have we seen so far?

What is the discrete weak formulation?

$$\int_{\Omega} u_h \varphi dx + \int_{\Omega} \nabla u_h \cdot \nabla \varphi dx = \int_{\partial\Omega} b \varphi ds, \quad \forall \varphi \in V_h$$

where

$$V_h = \{v \in C^0(\Omega) \mid v_K \in \mathbb{P}^p\}.$$

How do we get a linear algebra problem?

Consider and plug $\phi = \phi_i$ and $u_h = \sum_j U_j \varphi_j$ into the equation.

$$\sum_j U_j \underbrace{\int_{\Omega} \varphi_i \varphi_j dx}_{=M_{ij}} + \sum_j U_j \underbrace{\int_{\Omega} \nabla \varphi_i \cdot \nabla \varphi_j dx}_{=S_{ij}} = \underbrace{\int_{\partial\Omega} b \varphi_i ds}_{=F(\phi_i)}.$$

What have we seen so far?

What is the discrete weak formulation?

$$\int_{\Omega} u_h \varphi dx + \int_{\Omega} \nabla u_h \cdot \nabla \varphi dx = \int_{\partial\Omega} b \varphi ds, \quad \forall \varphi \in V_h$$

where

$$V_h = \{v \in C^0(\Omega) \mid v_K \in \mathbb{P}^p\}.$$

How do we get a linear algebra problem?

Consider and plug $\phi = \phi_i$ and $u_h = \sum_j U_j \varphi_j$ into the equation.

$$\sum_j U_j \left[\underbrace{\int_{\Omega} \varphi_i \varphi_j dx}_{=M_{ij}} + \underbrace{\int_{\Omega} \nabla \varphi_i \cdot \nabla \varphi_j dx}_{=S_{ij}} \right] = \underbrace{\int_{\partial\Omega} b \varphi_i ds}_{=F(\phi_i)}.$$

Linear algebra problem from a FE discretization

We want to solve

$$\sum_j U_j \underbrace{\int_{\Omega} \varphi_i \varphi_j + \nabla \varphi_i \cdot \nabla \varphi_j dx}_{=A_{ij}} = \underbrace{\int_{\partial\Omega} b \varphi_i ds}_{=F(\phi_i)}, \quad i = 1, \dots, \dim(V_h).$$

Linear algebra problem from a FE discretization

We want to solve

$$\sum_j U_j \underbrace{\int_{\Omega} \varphi_i \varphi_j + \nabla \varphi_i \cdot \nabla \varphi_j dx}_{=A_{ij}} = \underbrace{\int_{\partial\Omega} b \varphi_i ds}_{=F(\phi_i)}, \quad i = 1, \dots, \dim(V_h).$$

Recall the space V_h and that $\phi_i \in V_h$

$$V_h = \{\varphi \in C^0(\Omega) \mid \varphi|_K \in \mathbb{P}^p\}$$

Linear algebra problem from a FE discretization

We want to solve

$$\sum_j U_j \underbrace{\int_{\Omega} \varphi_i \varphi_j + \nabla \varphi_i \cdot \nabla \varphi_j dx}_{=A_{ij}} = \underbrace{\int_{\partial\Omega} b \varphi_i ds}_{=F(\phi_i)}, \quad i = 1, \dots, \dim(V_h).$$

Recall the space V_h and that $\phi_i \in V_h$

$$V_h = \{\varphi \in C^0(\Omega) \mid \varphi|_K \in \mathbb{P}^p\}$$

Assume $\Omega = (0, 1)$ and consider $p = 1$

Linear algebra problem from a FE discretization

We want to solve

$$\sum_j U_j \underbrace{\int_{\Omega} \varphi_i \varphi_j + \nabla \varphi_i \cdot \nabla \varphi_j dx}_{=A_{ij}} = \underbrace{\int_{\partial\Omega} b \varphi_i ds}_{=F(\phi_i)}, \quad i = 1, \dots, \dim(V_h).$$

Recall the space V_h and that $\phi_i \in V_h$

$$V_h = \{\varphi \in C^0(\Omega) \mid \varphi|_K \in \mathbb{P}^p\}$$

Assume $\Omega = (0, 1)$ and consider $p = 1$

- * How do the shape (or basis) functions look like?

Linear algebra problem from a FE discretization

We want to solve

$$\sum_j U_j \underbrace{\int_{\Omega} \varphi_i \varphi_j + \nabla \varphi_i \cdot \nabla \varphi_j dx}_{=A_{ij}} = \underbrace{\int_{\partial\Omega} b \varphi_i ds}_{=F(\phi_i)}, \quad i = 1, \dots, \dim(V_h).$$

Recall the space V_h and that $\phi_i \in V_h$

$$V_h = \{\varphi \in C^0(\Omega) \mid \varphi|_K \in \mathbb{P}^p\}$$

Assume $\Omega = (0, 1)$ and consider $p = 1$

- * How do the shape (or basis) functions look like?
- * Let's consider a discretization of Ω .

Linear algebra problem from a FE discretization

We want to solve

$$\sum_j U_j \underbrace{\int_{\Omega} \varphi_i \varphi_j + \nabla \varphi_i \cdot \nabla \varphi_j dx}_{=A_{ij}} = \underbrace{\int_{\partial\Omega} b \varphi_i ds}_{=F(\phi_i)}, \quad i = 1, \dots, \dim(V_h).$$

Recall the space V_h and that $\phi_i \in V_h$

$$V_h = \{\varphi \in C^0(\Omega) \mid \varphi|_K \in \mathbb{P}^p\}$$

Assume $\Omega = (0, 1)$ and consider $p = 1$

- * How do the shape (or basis) functions look like?
- * Let's consider a discretization of Ω .
- * Let's compute the matrices and RHS.

Linear algebra problem from a FE discretization

Second example:

Let's consider the 1D equation (from the first week):

$$\partial_t u + a \partial_x u - \mu \partial_{xx} u = r(x), \quad \forall x \in \Omega = (0, 1),$$

with $\mu > 0$.

Linear algebra problem from a FE discretization

Second example:

Let's consider the 1D equation (from the first week):

$$\partial_t u + a \partial_x u - \mu \partial_{xx} u = r(x), \quad \forall x \in \Omega = (0, 1),$$

with $\mu > 0$.

Get the finite element spatial semi-discretization:

Linear algebra problem from a FE discretization

Second example:

Let's consider the 1D equation (from the first week):

$$\partial_t u + a \partial_x u - \mu \partial_{xx} u = r(x), \quad \forall x \in \Omega = (0, 1),$$

with $\mu > 0$.

Get the finite element spatial semi-discretization:

- * Consider piecewise linear continuous finite elements.

Linear algebra problem from a FE discretization

Second example:

Let's consider the 1D equation (from the first week):

$$\partial_t u + a \partial_x u - \mu \partial_{xx} u = r(x), \quad \forall x \in \Omega = (0, 1),$$

with $\mu > 0$.

Get the finite element spatial semi-discretization:

- * Consider piecewise linear continuous finite elements.
- * What is the discrete weak formulation?

Linear algebra problem from a FE discretization

Second example:

Let's consider the 1D equation (from the first week):

$$\partial_t u + a \partial_x u - \mu \partial_{xx} u = r(x), \quad \forall x \in \Omega = (0, 1),$$

with $\mu > 0$.

Get the finite element spatial semi-discretization:

- * Consider piecewise linear continuous finite elements.
- * What is the discrete weak formulation?
- * Compute the mass, transport and stiffness matrices.

Linear algebra problem from a FE discretization

Second example:

Let's consider the 1D equation (from the first week):

$$\partial_t u + a \partial_x u - \mu \partial_{xx} u = r(x), \quad \forall x \in \Omega = (0, 1),$$

with $\mu > 0$.

Get the finite element spatial semi-discretization:

- * Consider piecewise linear continuous finite elements.
- * What is the discrete weak formulation?
- * Compute the mass, transport and stiffness matrices.
- * Express the equation for the i -th DoF.

Linear algebra problem from a FE discretization

Second example:

Let's consider the 1D equation (from the first week):

$$\partial_t u + a \partial_x u - \mu \partial_{xx} u = r(x), \quad \forall x \in \Omega = (0, 1),$$

with $\mu > 0$.

Get the finite element spatial semi-discretization:

- * Consider piecewise linear continuous finite elements.
- * What is the discrete weak formulation?
- * Compute the mass, transport and stiffness matrices.
- * Express the equation for the i -th DoF.
- * Compare this versus the finite difference formulation.

Summary of what we did

Given the discrete weak formulation we proceeded as follows:

Summary of what we did

Given the discrete weak formulation we proceeded as follows:

- * Identify each matrix and vectors we needed to compute.

Summary of what we did

Given the discrete weak formulation we proceeded as follows:

- * Identify each matrix and vectors we needed to compute.
- * To compute the matrices and vectors we needed to perform integrals on the domain.

Summary of what we did

Given the discrete weak formulation we proceeded as follows:

- * Identify each matrix and vectors we needed to compute.
- * To compute the matrices and vectors we needed to perform integrals on the domain.
- * We broke the **integration over** the domain in **elements**.

Summary of what we did

Given the discrete weak formulation we proceeded as follows:

- * Identify each matrix and vectors we needed to compute.
- * To compute the matrices and vectors we needed to perform integrals on the domain.
- * We broke the integration over the domain in elements.
- * We considered any node i and its shape function φ_i .

Summary of what we did

Given the discrete weak formulation we proceeded as follows:

- * Identify each matrix and vectors we needed to compute.
- * To compute the matrices and vectors we needed to perform integrals on the domain.
- * We broke the **integration over** the domain in **elements**.
- * We **considered any node i** and its shape function φ_i .

Note: if the mesh is not uniform, we need to consider all nodes independently.

Summary of what we did

Given the discrete weak formulation we proceeded as follows:

- * Identify each matrix and vectors we needed to compute.
- * To compute the matrices and vectors we needed to perform integrals on the domain.
- * We broke the **integration over** the domain in **elements**.
- * We **considered any node i** and its shape function φ_i .
Note: if the mesh is not uniform, we need to consider all nodes independently.
- * To compute the matrices, **for a given node i , we considered all j -shape functions**.

Summary of what we did

Given the discrete weak formulation we proceeded as follows:

- * Identify each matrix and vectors we needed to compute.
- * To compute the matrices and vectors we needed to perform integrals on the domain.
- * We broke the **integration over** the domain in **elements**.
- * We **considered any node i** and its shape function φ_i .
Note: if the mesh is not uniform, we need to consider all nodes independently.
- * To compute the matrices, **for a given node i , we considered all j -shape functions**.
- * We performed the **integration based on a reference element**.

Quadratures

We approximate the integrals via quadratures. That is (for 1D),

$$\int_{-1}^1 f(x) dx \approx \sum_{q=1}^{N_q} w_q f(x_q),$$

where N_q is the number of quadrature points.

Quadratures

We approximate the integrals via quadratures. That is (for 1D),

$$\int_{-1}^1 f(x) dx \approx \sum_{q=1}^{N_q} w_q f(x_q),$$

where N_q is the number of quadrature points.

Gaussian quadratures

Quadratures

We approximate the integrals via quadratures. That is (for 1D),

$$\int_{-1}^1 f(x) dx \approx \sum_{q=1}^{N_q} w_q f(x_q),$$

where N_q is the number of quadrature points.

Gaussian quadratures

- * In finite elements, it is common to use Gaussian quadratures.

Quadratures

We approximate the integrals via quadratures. That is (for 1D),

$$\int_{-1}^1 f(x) dx \approx \sum_{q=1}^{N_q} w_q f(x_q),$$

where N_q is the number of quadrature points.

Gaussian quadratures

- * In finite elements, it is common to use Gaussian quadratures.
- * The quad points $\{x_q\}$ are the roots of a polynomial.

Quadratures

We approximate the integrals via quadratures. That is (for 1D),

$$\int_{-1}^1 f(x) dx \approx \sum_{q=1}^{N_q} w_q f(x_q),$$

where N_q is the number of quadrature points.

Gaussian quadratures

- * In finite elements, it is common to use Gaussian quadratures.
- * The quad points $\{x_q\}$ are the roots of a polynomial.
- * Gauss-Legendre: $\{x_q\}$ are the roots of Legendre polynomials.

Quadratures

We approximate the integrals via quadratures. That is (for 1D),

$$\int_{-1}^1 f(x) dx \approx \sum_{q=1}^{N_q} w_q f(x_q),$$

where N_q is the number of quadrature points.

Gaussian quadratures

- * In finite elements, it is common to use Gaussian quadratures.
- * The quad points $\{x_q\}$ are the roots of a polynomial.
- * Gauss-Legendre: $\{x_q\}$ are the roots of Legendre polynomials.
- * Gauss-Legendre quadrature is exact for polynomials of degree up to $2N_q - 1$.

Quadratures: Gauss-Legendre


Quadrature points

The quadrature points $\{x_1, \dots, x_{N_q}\}$ are the roots of the Legendre polynomials of degree N_q .

Quadratures: Gauss-Legendre

Quadrature points

The quadrature points $\{x_1, \dots, x_{N_q}\}$ are the roots of the Legendre polynomials of degree N_q .



Quadratures: Gauss-Legendre

Quadrature weights

The weights are defined so that polynomials of degree up to $2N_q - 1$ are integrated exactly.

Quadratures: Gauss-Legendre

Quadrature weights

The weights are defined so that polynomials of degree up to $2N_q - 1$ are integrated exactly.

Example:

Consider piecewise linear poly. to solve the advection-diffusion eqn.

Quadratures: Gauss-Legendre

Quadrature weights

The weights are defined so that polynomials of degree up to $2N_q - 1$ are integrated exactly.

Example:

Consider piecewise linear poly. to solve the advection-diffusion eqn.

- * What is the highest degree of polynomials that I need to integrate?

Quadratures: Gauss-Legendre

Quadrature weights

The weights are defined so that polynomials of degree up to $2N_q - 1$ are integrated exactly.

Example:

Consider piecewise linear poly. to solve the advection-diffusion eqn.

- * What is the highest degree of polynomials that I need to integrate?
- * The highest degree is given by the mass matrix.

Quadratures: Gauss-Legendre

Quadrature weights

The weights are defined so that polynomials of degree up to $2N_q - 1$ are integrated exactly.

Example:

Consider piecewise linear poly. to solve the advection-diffusion eqn.

- * What is the highest degree of polynomials that I need to integrate?
- * The highest degree is given by the mass matrix.
- * The integrand of the mass matrix is quadratic.

Quadratures: Gauss-Legendre

Quadrature weights

The weights are defined so that polynomials of degree up to $2N_q - 1$ are integrated exactly.

Example:

Consider piecewise linear poly. to solve the advection-diffusion eqn.

- * What is the highest degree of polynomials that I need to integrate?
- * The highest degree is given by the mass matrix.
- * The integrand of the mass matrix is quadratic.
- * Then I need $2N_q - 1 \geq 2$. Therefore, $N_q \geq 3/2 \implies N_q = 2$.

Quadratures: Gauss-Legendre

Quadrature weights

The weights are defined so that polynomials of degree up to $2N_q - 1$ are integrated exactly.

Example:

Consider piecewise linear poly. to solve the advection-diffusion eqn.

- * What is the highest degree of polynomials that I need to integrate?
- * The highest degree is given by the mass matrix.
- * The integrand of the mass matrix is quadratic.
- * Then I need $2N_q - 1 \geq 2$. Therefore, $N_q \geq 3/2 \implies N_q = 2$.
- * With $N_q = 2$, I can integrate exactly polynomials up to degree 3.

Quadratures: Gauss-Legendre

Quadrature weights

The weights are defined so that polynomials of degree up to $2N_q - 1$ are integrated exactly.

Example:

Consider piecewise linear poly. to solve the advection-diffusion eqn.

- * What is the highest degree of polynomials that I need to integrate?
- * The highest degree is given by the mass matrix.
- * The integrand of the mass matrix is quadratic.
- * Then I need $2N_q - 1 \geq 2$. Therefore, $N_q \geq 3/2 \implies N_q = 2$.
- * With $N_q = 2$, I can integrate exactly polynomials up to degree 3.

Verify that I can integrate third order polynomials with $N_q = 2$.

The finite element (FE) loop

The kernel of most FE codes is the FE loop

The finite element (FE) loop

The kernel of most FE codes is the FE loop

- * Loop on elements. We do this since we broke the integral in element integrals.

The finite element (FE) loop

The kernel of most FE codes is the FE loop

- * Loop on elements. We do this since we broke the integral in element integrals.
- * Compute element based integrals (vectors and matrices).

The finite element (FE) loop

The kernel of most FE codes is the FE loop

- * Loop on elements. We do this since we broke the integral in element integrals.
- * Compute element based integrals (vectors and matrices).
- * Combine (or assemble) the element integrals to global integrals.

The finite element (FE) loop

The kernel of most FE codes is the FE loop

- * Loop on elements. We do this since we broke the integral in element integrals.
- * Compute element based integrals (vectors and matrices).
- * Combine (or assemble) the element integrals to global integrals.

How to compute the element based integrals?

For each element, proceed as follows:

The finite element (FE) loop

The kernel of most FE codes is the FE loop

- * Loop on elements. We do this since we broke the integral in element integrals.
- * Compute element based integrals (vectors and matrices).
- * Combine (or assemble) the element integrals to global integrals.

How to compute the element based integrals?

For each element, proceed as follows:

- * Compute all quantities with respect to the reference element.

The finite element (FE) loop

The kernel of most FE codes is the FE loop

- * Loop on elements. We do this since we broke the integral in element integrals.
- * Compute element based integrals (vectors and matrices).
- * Combine (or assemble) the element integrals to global integrals.

How to compute the element based integrals?

For each element, proceed as follows:

- * Compute all quantities with respect to the reference element.
- * Loop on quadrature points. We do this since we approximate the integrals via quadratures.

The finite element (FE) loop

The kernel of most FE codes is the FE loop

- * Loop on elements. We do this since we broke the integral in element integrals.
- * Compute element based integrals (vectors and matrices).
- * Combine (or assemble) the element integrals to global integrals.

How to compute the element based integrals?

For each element, proceed as follows:

- * Compute all quantities with respect to the reference element.
- * Loop on quadrature points. We do this since we approximate the integrals via quadratures.
- * Loop on i -DoFs.

The finite element (FE) loop

The kernel of most FE codes is the FE loop

- * Loop on elements. We do this since we broke the integral in element integrals.
- * Compute element based integrals (vectors and matrices).
- * Combine (or assemble) the element integrals to global integrals.

How to compute the element based integrals?

For each element, proceed as follows:

- * Compute all quantities with respect to the reference element.
- * Loop on quadrature points. We do this since we approximate the integrals via quadratures.
- * Loop on i -DoFs.
- * Loop on j -DoFs and more indices if needed.

The finite element (FE) loop

The FE loop looks as follows:

The finite element (FE) loop

The FE loop looks as follows:

for $K = 1$ to N_{el} **do**

end for

The finite element (FE) loop

The FE loop looks as follows:

for $K = 1$ to N_{el} **do**

 Compute quantities wrt reference element.

end for

The finite element (FE) loop

The FE loop looks as follows:

for $K = 1$ to N_{el} do

 Compute quantities wrt reference element.

for $q = 1$ to N_q do

end for

end for

The finite element (FE) loop

The FE loop looks as follows:

for $K = 1$ to N_{el} **do**

 Compute quantities wrt reference element.

for $q = 1$ to N_q **do**

for $i = 1$ to DoFs per element **do**

end for

end for

end for

The finite element (FE) loop

The FE loop looks as follows:

for $K = 1$ to N_{el} do

 Compute quantities wrt reference element.

for $q = 1$ to N_q do

for $i = 1$ to DoFs per element do

 Compute element based vectors.

end for

end for

end for

The finite element (FE) loop

The FE loop looks as follows:

for $K = 1$ to N_{el} do

 Compute quantities wrt reference element.

for $q = 1$ to N_q do

for $i = 1$ to DoFs per element do

 Compute element based vectors.

for $j = 1$ to DoFs per element do

end for

end for

end for

end for

The finite element (FE) loop

The FE loop looks as follows:

for $K = 1$ to N_{el} do

 Compute quantities wrt reference element.

for $q = 1$ to N_q do

for $i = 1$ to DoFs per element do

 Compute element based vectors.

for $j = 1$ to DoFs per element do

 Compute element based matrices.

end for

end for

end for

end for

The finite element (FE) loop

The FE loop looks as follows:

for $K = 1$ to N_{el} **do**

 Compute quantities wrt reference element.

for $q = 1$ to N_q **do**

for $i = 1$ to DoFs per element **do**

 Compute element based vectors.

for $j = 1$ to DoFs per element **do**

 Compute element based matrices.

end for

end for

 Assemble from local to global operators.

end for

end for