

## Homework 3

### Exercise 1: projection of a smooth function to $C^0$ finite element spaces

Consider the function

$$f(x) = \sin^4(2\pi x) \sin^4(2\pi y), \quad \forall (x, y) \in [0, 1]^2,$$

and project it to a finite element space using triangular elements and piecewise linear polynomials. Consider  $N_{\text{el}} = 100, 400, 1600, 6400$  elements. For each refinement level compute the errors

$$E_1 := \int_0^1 |f(x) - f_h(x)| dx, \quad E_2 := \sqrt{\int_0^1 (f(x) - f_h(x))^2 dx},$$

where  $f_h(x)$  is the projection. Report the four plots and the convergence table.

### Exercise 2: projection of a non-smooth function to $C^0$ finite element spaces

Repeat exercise 1 but considering

$$f(x) = \begin{cases} 1, & \text{if } x \in [0.35, 0.65] \text{ and } y \in [0.35, 0.65], \\ 0, & \text{otherwise} \end{cases}$$

instead.

### Exercise 3: solve the Poisson equation

Consider the two-dimensional Poisson equation

$$\begin{aligned} -\Delta u &= f, & \Omega &= [0, 1]^2, \\ u &= 0, & \forall x &\in \partial\Omega. \end{aligned}$$

where

$$f = \begin{cases} 1, & \text{if } y > 0.5 + 0.25 \sin(4\pi x), \\ -1, & \text{if } y \leq 0.5 + 0.25 \sin(4\pi x). \end{cases}$$

Consider a finite element space using triangular elements and piecewise linear polynomials. Solve the equation using  $N_{\text{el}} = 100, 400, 1600, 6400$  elements. Report the four plots.

**Exercise 4: solve the Eikonal equation in a unit square**

In this exercise, you can use either quadrilateral or triangular elements with piecewise bilinear or linear, respectively, polynomials. The domain is  $\Omega = [0, 1]^2$ . Consider

$$\tilde{u}(x, y) = \begin{cases} h, & \text{if } \sqrt{(x - 0.5)^2 + (y - 0.5)^2} \leq 0.25, \\ -h, & \text{if } \sqrt{(x - 0.5)^2 + (y - 0.5)^2} > 0.25, \end{cases}$$

where  $h = \Delta x = \Delta y$  is the mesh size. Use the lumped mass matrix to project  $\tilde{u}$  into the finite element space to obtain

$$\tilde{u}_h(x, y) = \sum_j \tilde{U}_j \phi_j(x, y).$$

Consider the following discrete weak formulation:

$$\alpha m_i^{\Gamma(\tilde{u}_h)} (U_i - \tilde{U}_i) + \int_{\Omega} S_{\epsilon}(\tilde{u}_h) [|\nabla u_h| - 1] \phi d\mathbf{x} + ch \int_{\Omega} \nabla u_h \cdot \nabla \phi d\mathbf{x} = 0, \quad \forall \phi \in V_h \quad (1)$$

where  $\alpha = 10^{10}$ ,  $c = 0.5$ ,  $m_i^{\Gamma(\tilde{u}_h)} = \int_{\Omega} \delta_{\epsilon}(\tilde{u}_h) \phi_i d\mathbf{x}$ , and

$$H_{\epsilon}(u) = \begin{cases} 0, & \text{if } u \leq -\epsilon, \\ \frac{1}{2} \left[ 1 + \frac{u}{\epsilon} + \frac{1}{\pi} \sin \left( \frac{\pi u}{\epsilon} \right) \right], & \text{if } -\epsilon \leq u \leq \epsilon, \\ 1, & \text{if } u \geq \epsilon, \end{cases}$$

$$S_{\epsilon}(u) = 2H_{\epsilon}(u) - 1,$$

$$\delta_{\epsilon}(u) = H'_{\epsilon}(u),$$

$$\epsilon = 0.1h.$$

Solve (1) using  $N_{\text{el}} = 400, 1600, 6400$  elements. Report the three plots.