

AMCS 394E: Contemp. Topics in Computational Science.

Computing with the finite element method

David I. Ketcheson and Manuel Quezada de Luna



Solution of PDEs

The solution of a PDE is defined at every point in a given domain Ω .

Solution of PDEs

The solution of a PDE is defined at every point in a given domain Ω .

For example:

The solution $u(x)$ of

$$-\Delta u = f(x), \quad \forall x \in (0, 1)$$

describes some distribution (e.g., of heat) at every point between $(0, 1)$.

Solution of PDEs

The solution of a PDE is defined at every point in a given domain Ω .

Problems:

Solution of PDEs

The solution of a PDE is defined at every point in a given domain Ω .

Problems:

- * Need infinite processing power to get the solution at every point.

Solution of PDEs

The solution of a PDE is defined at every point in a given domain Ω .

Problems:

- * Need infinite processing power to get the solution at every point.
- * Even if I know the exact solution, I can't store or represent the solution in any computer.

Solution of PDEs

The solution of a PDE is defined at every point in a given domain Ω .

Problems:

- * Need infinite processing power to get the solution at every point.
- * Even if I know the exact solution, I can't store or represent the solution in any computer.

Alternatives:

Solution of PDEs

The solution of a PDE is defined at every point in a given domain Ω .

Problems:

- * Need infinite processing power to get the solution at every point.
- * Even if I know the exact solution, I can't store or represent the solution in any computer.

Alternatives:

- * Represent the solution in finitely many points.

Solution of PDEs

The solution of a PDE is defined at every point in a given domain Ω .

Problems:

- * Need infinite processing power to get the solution at every point.
- * Even if I know the exact solution, I can't store or represent the solution in any computer.

Alternatives:

- * Represent the solution in finitely many points.
- * Use finite information to approximate the solution.

Solution of PDEs

The solution of a PDE is defined at every point in a given domain Ω .

Questions:

Solution of PDEs

The solution of a PDE is defined at every point in a given domain Ω .

Questions:

- * how do I approximate the solution?

Solution of PDEs

The solution of a PDE is defined at every point in a given domain Ω .

Questions:

- * how do I approximate the solution?
- * how do I obtain the solution?

How can we approximate the solution?

In general we can consider local or global approximations.

How can we approximate the solution?

In general we can consider local or global approximations.

Global approximations:

For example via Taylor series:

$$\begin{aligned} f(x) &= \sum_{n=0}^{\infty} \frac{f^{(n)}(x_0)}{n!} (x - x_0)^n \\ &\approx \sum_{n=0}^N \alpha_n (x - x_0)^n \end{aligned}$$


Exercise:

Consider $f(x) = \sin(x)$ and approximate it via Taylor series.

How can we approximate the solution?

Exercise:

Consider $f(x) = \sin(x)$ and approximate it via Taylor series.




How can we approximate the solution?

In general we can consider local or global approximations.

Local approximations:

For example, via local (linear) polynomial approximations.




How can we approximate the solution?

In general we can consider local or global approximations.

Local approximations:

For example, via local (linear) polynomial approximations.




How can we approximate the solution?

In general we can consider local or global approximations.

Local approximations:

For example, via local (linear) polynomial approximations.




How can we approximate the solution?

In general we can consider local or global approximations.

Local approximations:

For example, via local (linear) polynomial approximations.



How can we approximate the solution?

In general we can consider local or global approximations.

Other approximations:

We can consider other approximations like Fourier series.

How can we approximate the solution?

In general we can consider local or global approximations.

Other approximations:

We can consider other approximations like Fourier series.

Questions:

How can we approximate the solution?

In general we can consider local or global approximations.

Other approximations:

We can consider other approximations like Fourier series.

Questions:

- * Why can we consider polynomials?

How can we approximate the solution?

In general we can consider local or global approximations.

Other approximations:

We can consider other approximations like Fourier series.

Questions:

- * Why can we consider polynomials?
- * Should we use global or local approximations?

Some definitions

Vector space

Is a set of objects that can be added together and multiplied by scalars (of a given field) and yield a new object also in the set.

Some definitions

Vector space

Is a set of objects that can be added together and multiplied by scalars (of a given field) and yield a new object also in the set.

Examples:

Some definitions

Vector space

Is a set of objects that can be added together and multiplied by scalars (of a given field) and yield a new object also in the set.

Examples:

- * Real numbers.

Some definitions

Vector space

Is a set of objects that can be added together and multiplied by scalars (of a given field) and yield a new object also in the set.

Examples:

- * Real numbers.
- * Euclidean spaces like \mathbb{R}^2 or \mathbb{R}^3 .

Some definitions

Vector space

Is a set of objects that can be added together and multiplied by scalars (of a given field) and yield a new object also in the set.

Examples:

- * Real numbers.
- * Euclidean spaces like \mathbb{R}^2 or \mathbb{R}^3 .
- * Set of continuous functions $C^0(\Omega)$.

Some definitions

Vector space

Is a set of objects that can be added together and multiplied by scalars (of a given field) and yield a new object also in the set.

Examples:

- * Real numbers.
- * Euclidean spaces like \mathbb{R}^2 or \mathbb{R}^3 .
- * Set of continuous functions $C^0(\Omega)$.

Important concepts of a vector space:

Basis and dimension.

Basis of a vector space

A set of elements in a (vector) space V is a basis of the space if every element in V can be written as a linear combination of the basis.

Basis of a vector space

A set of elements in a (vector) space V is a basis of the space if every element in V can be written as a linear combination of the basis.

Meaning:

Consider the set $B = \{\varphi_i\}_{i \in I}$ and any $v \in V$.

Basis of a vector space

A set of elements in a (vector) space V is a basis of the space if every element in V can be written as a linear combination of the basis.

Meaning:

Consider the set $B = \{\varphi_i\}_{i \in I}$ and any $v \in V$.

Then, B is a basis of V if we can write v as follows:

$$v = \sum_i a_i \varphi_i,$$

where a_i are the coordinates of v .

Basis of a vector space

A set of elements in a (vector) space V is a basis of the space if every element in V can be written as a linear combination of the basis.

Meaning:

Consider the set $B = \{\varphi_i\}_{i \in I}$ and any $v \in V$.

Then, B is a basis of V if we can write v as follows:

$$v = \sum_i a_i \varphi_i,$$

where a_i are the coordinates of v .

Requirement:

The elements of the basis B must be linearly independent.

Dimension of a vector space

The dimension of the vector space is the number of elements in the basis. Note that any basis of V has the same number of elements.

Examples of vector spaces and basis

Example 1:

Three dimensional euclidean space.

- * Elements in the space are all vectors in \mathbb{R}^3 .
- * Example of a basis is the set of orthonormal vectors $(\hat{i}, \hat{j}, \hat{k})$.
- * Dimension of the space is 3.

Examples of vector spaces and basis

Example 2:

Polynomial space of degree p .

- * Elements in the space are all polynomials with degree at most p .
- * Example of a basis is the set of monomials $\{1, x, x^2, \dots, x^p\}$.
- * Dimension of the space is $p + 1$.

Examples of vector spaces and basis

Example 3:

Space of infinitely smooth functions in 1D: $C^\infty(\mathbb{R})$.

- * Elements in the space are all 1D infinitely smooth functions.
- * Examples of basis functions are:
 - set of monomials $\{1, x, x^2, \dots\}$.
 - trigonometric functions $\{e^{2\pi i n x}\}_{n=-\infty}^{\infty}$.
- * Dimension of the space is infinity.

Examples of vector spaces and basis

Example 4:

square integrable functions in 1D: $L^2(\mathbb{R})$.

- * Elements in the space are all 1D functions s.t.: $\int_{\mathbb{R}} f^2(x) dx < \infty$
- * Examples of basis functions are:
 - set of monomials $\{1, x, x^2, \dots\}$.
 - trigonometric functions $\{e^{2\pi i n x}\}_{n=-\infty}^{\infty}$.
- * Dimension of the space is infinity.

Some definitions

Dense set

A set S is dense in V if every element in V is either in S or is a limit point of S .

Limit point of a set S

The limit point of S is any point that can be approx. by points in S .

Example:

The rationals are dense in the real numbers.

Why do we even consider polynomials in the first place?

Weierstrass approximation theorem:

Let $f(x)$ be a real valued and continuous function in an interval $[a, b]$.
Then for any $\epsilon > 0$, there is a polynomial $p(x)$ in $[a, b]$ s.t.

$$|f(x) - p(x)| < \epsilon, \quad \forall x \in [a, b]$$

Why do we even consider polynomials in the first place?

Weierstrass approximation theorem:

Let $f(x)$ be a real valued and continuous function in an interval $[a, b]$.
Then for any $\epsilon > 0$, there is a polynomial $p(x)$ in $[a, b]$ s.t.

$$|f(x) - p(x)| < \epsilon, \quad \forall x \in [a, b]$$

What does this mean?

Why do we even consider polynomials in the first place?

Weierstrass approximation theorem:

Let $f(x)$ be a real valued and continuous function in an interval $[a, b]$. Then for any $\epsilon > 0$, there is a polynomial $p(x)$ in $[a, b]$ s.t.

$$|f(x) - p(x)| < \epsilon, \quad \forall x \in [a, b]$$

What does this mean?

- * Polynomials are dense in the space of real valued cont. functions.

Why do we even consider polynomials in the first place?

Weierstrass approximation theorem:

Let $f(x)$ be a real valued and continuous function in an interval $[a, b]$. Then for any $\epsilon > 0$, there is a polynomial $p(x)$ in $[a, b]$ s.t.

$$|f(x) - p(x)| < \epsilon, \quad \forall x \in [a, b]$$

What does this mean?

- * Polynomials are dense in the space of real valued cont. functions.
- * $f(x)$ can be uniformly approx. to any accuracy by a polynomial.
(uniformly means at every point)

Why do we even consider polynomials in the first place?

Weierstrass approximation theorem:

Let $f(x)$ be a real valued and continuous function in an interval $[a, b]$. Then for any $\epsilon > 0$, there is a polynomial $p(x)$ in $[a, b]$ s.t.

$$|f(x) - p(x)| < \epsilon, \quad \forall x \in [a, b]$$

What does this mean?

- * Polynomials are dense in the space of real valued cont. functions.
- * $f(x)$ can be uniformly approx. to any accuracy by a polynomial.

Should we consider global or local polynomials?

We need to consider:

Why do we even consider polynomials in the first place?

Weierstrass approximation theorem:

Let $f(x)$ be a real valued and continuous function in an interval $[a, b]$. Then for any $\epsilon > 0$, there is a polynomial $p(x)$ in $[a, b]$ s.t.

$$|f(x) - p(x)| < \epsilon, \quad \forall x \in [a, b]$$

What does this mean?

- * Polynomials are dense in the space of real valued cont. functions.
- * $f(x)$ can be uniformly approx. to any accuracy by a polynomial.

Should we consider global or local polynomials?

We need to consider:

- * Smoothness properties of the function

Why do we even consider polynomials in the first place?

Weierstrass approximation theorem:

Let $f(x)$ be a real valued and continuous function in an interval $[a, b]$. Then for any $\epsilon > 0$, there is a polynomial $p(x)$ in $[a, b]$ s.t.

$$|f(x) - p(x)| < \epsilon, \quad \forall x \in [a, b]$$

What does this mean?

- * Polynomials are dense in the space of real valued cont. functions.
- * $f(x)$ can be uniformly approx. to any accuracy by a polynomial.

Should we consider global or local polynomials?

We need to consider:

- * Smoothness properties of the function
- * Computational cost.

$L^2(\Omega)$ space

Given a domain Ω , the $L^2(\Omega)$ space is “more general” than $C^0(\Omega)$. This space is quite commonly considered in finite elements.

$L^2(\Omega)$ space

Given a domain Ω , the $L^2(\Omega)$ space is “more general” than $C^0(\Omega)$. This space is quite commonly considered in finite elements.

Definition:

Is a normed vector space composed of the set of all squared integrable functions.

$L^2(\Omega)$ space

Given a domain Ω , the $L^2(\Omega)$ space is “more general” than $C^0(\Omega)$. This space is quite commonly considered in finite elements.

Definition:

Is a normed vector space composed of the set of all squared integrable functions.

Relevance within finite elements:

Since $L^2(\Omega)$ is more general than $C^0(\Omega)$, solutions with discontinuities can be represented by the L^2 space.

$L^2(\Omega)$ space

Properties:

$L^2(\Omega)$ space

Properties:

- * $C^0(\Omega)$ is dense (w.r.t. the L^2 norm) in $L^2(\Omega)$.

$L^2(\Omega)$ space

Properties:

- * $C^0(\Omega)$ is dense (w.r.t. the L^2 norm) in $L^2(\Omega)$.

In simple words, ...

we can approximate functions in $L^2(\Omega)$ via
(sequences of) continuous functions.

Piecewise polynomial approximation of $L^2(\Omega)$

Summary so far:

Piecewise polynomial approximation of $L^2(\Omega)$

Summary so far:

- * We consider L^2 spaces to represent solutions of PDEs.

Piecewise polynomial approximation of $L^2(\Omega)$


Summary so far:

- * We consider L^2 spaces to represent solutions of PDEs.
- * We consider sequences of continuous functions to approximate the solution of the PDEs.

Piecewise polynomial approximation of $L^2(\Omega)$

Summary so far:


- * We consider L^2 spaces to represent solutions of PDEs.
- * We consider sequences of continuous functions to approximate the solution of the PDEs.



Piecewise polynomial approximation of $L^2(\Omega)$

Summary so far:


- * We consider L^2 spaces to represent solutions of PDEs.
- * We consider sequences of continuous functions to approximate the solution of the PDEs.



Piecewise polynomial approximation of $L^2(\Omega)$

Summary so far:


- * We consider L^2 spaces to represent solutions of PDEs.
- * We consider sequences of continuous functions to approximate the solution of the PDEs.



Piecewise polynomial approximation of $L^2(\Omega)$

Summary so far:

- * We consider L^2 spaces to represent solutions of PDEs.
- * We consider sequences of continuous functions to approximate the solution of the PDEs.



Piecewise polynomial approximation of $L^2(\Omega)$

The (red) sequence of piecewise linear functions approximates $f(x)$.

Piecewise polynomial approximation of $L^2(\Omega)$

The (red) sequence of piecewise linear functions approximates $f(x)$.

What is the error of this approximation?

Piecewise polynomial approximation of $L^2(\Omega)$

The (red) sequence of piecewise linear functions approximates $f(x)$.

What is the error of this approximation?

- * Consider the interval $[a, b]$.

Piecewise polynomial approximation of $L^2(\Omega)$

The (red) sequence of piecewise linear functions approximates $f(x)$.

What is the error of this approximation?

- * Consider the interval $[a, b]$.
- * Let $f(x)$ be the function to be approximated.

Piecewise polynomial approximation of $L^2(\Omega)$

The (red) sequence of piecewise linear functions approximates $f(x)$.

What is the error of this approximation?

- * Consider the interval $[a, b]$.
- * Let $f(x)$ be the function to be approximated.
- * Let $f_h(x)$ be a piecewise polynomial approximation over $[a, b]$.

Piecewise polynomial approximation of $L^2(\Omega)$

The (red) sequence of piecewise linear functions approximates $f(x)$.

What is the error of this approximation?

- * Consider the interval $[a, b]$.
- * Let $f(x)$ be the function to be approximated.
- * Let $f_h(x)$ be a piecewise polynomial approximation over $[a, b]$.

Then,

$$\|f(x) - f_h(x)\|_{L^2([a,b])} \leq C(b-a)^{p+1},$$

where p is the degree of the polynomial and C is a constant independent of u and $|b-a|$.

Common piecewise polynomial spaces used in finite elements

The most common (finite dimensional) space in finite elements is

$$\underbrace{V_h := \{v \in C^0(\Omega) \mid v|_K \in \mathbb{P}^p(K) + \text{BCs}\}}_{\text{(finite dim.) space of the finite element soln.}} \subset \underbrace{V := \{v \in L^2(\Omega) + \text{BCs}\}}_{\text{(infinite dim.) space of the exact soln.}}$$

where K is a cell in the mesh.

Common piecewise polynomial spaces used in finite elements

The most common (finite dimensional) space in finite elements is

$$\underbrace{V_h := \{v \in C^0(\Omega) \mid v|_K \in \mathbb{P}^p(K) + \text{BCs}\}}_{\text{(finite dim.) space of the finite element soln.}} \subset \underbrace{V := \{v \in L^2(\Omega) + \text{BCs}\}}_{\text{(infinite dim.) space of the exact soln.}}$$

where K is a cell in the mesh.

Basis functions for V_h .

Common piecewise polynomial spaces used in finite elements

The most common (finite dimensional) space in finite elements is

$$\underbrace{V_h := \{v \in C^0(\Omega) \mid v|_K \in \mathbb{P}^p(K) + \text{BCs}\}}_{\text{(finite dim.) space of the finite element soln.}} \subset \underbrace{V := \{v \in L^2(\Omega) + \text{BCs}\}}_{\text{(infinite dim.) space of the exact soln.}}$$

where K is a cell in the mesh.

Basis functions for V_h .

- * We need $\dim(V_h)$ linearly independent functions $\varphi \in V_h$.

Common piecewise polynomial spaces used in finite elements

The most common (finite dimensional) space in finite elements is

$$\underbrace{V_h := \{v \in C^0(\Omega) \mid v|_K \in \mathbb{P}^p(K) + \text{BCs}\}}_{\text{(finite dim.) space of the finite element soln.}} \subset \underbrace{V := \{v \in L^2(\Omega) + \text{BCs}\}}_{\text{(infinite dim.) space of the exact soln.}}$$

where K is a cell in the mesh.

Basis functions for V_h .

- * We need $\dim(V_h)$ linearly independent functions $\varphi \in V_h$.
- * Consider the 'hat' functions.

Common piecewise polynomial spaces used in finite elements

The most common (finite dimensional) space in finite elements is

$$\underbrace{V_h := \{v \in C^0(\Omega) \mid v|_K \in \mathbb{P}^p(K) + \text{BCs}\}}_{\text{(finite dim.) space of the finite element soln.}} \subset \underbrace{V := \{v \in L^2(\Omega) + \text{BCs}\}}_{\text{(infinite dim.) space of the exact soln.}}$$

where K is a cell in the mesh.

Basis functions for V_h .

- * We need $\dim(V_h)$ linearly independent functions $\varphi \in V_h$.
- * Consider the ‘hat’ functions.
- * Interpolatory versus non-interpolatory basis functions.

Common piecewise polynomial spaces used in finite elements

Finite element solution

Given the basis $\{\varphi_j, j = 1, \dots, \dim(V_h)\}$, the finite element solution is

$$u_h(x) = \sum_{n=1}^{\dim(V_h)} a_n \varphi_n(x),$$

where a_i are called degrees of freedom (DoFs) or control points.

Common piecewise polynomial spaces used in finite elements

Finite element solution

Given the basis $\{\varphi_i, j = 1, \dots, \dim(V_h)\}$, the finite element solution is

$$u_h(x) = \sum_{n=1}^{\dim(V_h)} a_i \varphi_i(x),$$

where a_i are called degrees of freedom (DoFs) or control points.

Note: if the basis is interpolatory; i.e., if $\varphi_i(x_j) = \delta_{ij}$, then

$$a_i = u_h(x_i)$$