

Local Validity and Contextual Holonomy: A Structural Formulation of the Copenhagen Principle (v3 — complete rewrite)

Manuel Flores Gordillo

Independent Researcher ORCID 0009-0006-8246-0343

`manuel.flores@columbia.edu`

July 2026 — draft v3, pending adversarial review

Abstract

We propose that the Copenhagen viewpoint, stripped of its rhetoric, is a set of structural statements about one geometric object: the transformation groupoid $\mathrm{PU}(n) \ltimes X$ of measurement contexts (maximal abelian subalgebras) with its canonical $\mathrm{U}(1)$ 2-cocycle. In this formulation: *local validity* — the exact classicality of every single context — is a theorem with an explicit witness; *complementarity* is the nonvanishing holonomy of the cocycle along closed context loops, which detects exactly the all-versus-nothing sector of contextuality (an equivalence, now proved), with the achievable state-independent values classified and the state sector governed by a facet geometry that is now completely enumerated in the certified $d = 4$ example (23,256 facets in 61 symmetry classes) and classified in closed form at $d = 2$; *collapse* is pinned by statistics, repeatability, and covariance to a one-parameter family per degenerate block whose unique efficient member is the Lüders rule — with a classical twin theorem identifying the same axioms’ classical solution as Bayesian conditioning; *observer-relativity of facts* and the *movable cut* are theorems of an enlarged observer-context groupoid; and the Frauchiger–Renner argument localizes to an ordinary state-dependent violation (contextual fraction exactly $1/6$) carried by a single CHSH facet, with the loop holonomy of the observer switch provably trivial. Two of these signatures have been measured on superconducting hardware. Each tenet is either a theorem of the companion papers, a machine-verified certificate in the public suite, or is explicitly labeled a conjecture. Version v1 of this note contained four errors of substance; §2 states them and their repairs.

1 Introduction: the principle, restated

Textbook presentations of the Copenhagen interpretation bundle together several commitments: that classical concepts apply exactly within a single experimental arrangement; that arrangements exclude one another (complementarity); that measurement updates the state by projection; that facts are relative to the experimental context, including the observer; and that the boundary between quantum system and classical apparatus — the cut — can be moved without empirical consequence. Critics have long held that these are attitudes rather than assertions. This note argues the opposite: each commitment is a precise structural statement about a single mathematical object, and each is now either proved, machine-certified in a public verification suite, or stated as an explicit conjecture with its current evidence.

The object is the transformation groupoid

$$\mathcal{G} = \text{PU}(n) \ltimes X,$$

where X is the space of maximal abelian $*$ -subalgebras (MASAs, i.e. measurement contexts) of $M_n(\mathbb{C})$ and the projective unitary group acts by conjugation, together with the canonical $\text{U}(1)$ 2-cocycle obtained from the central extension $\text{U}(1) \rightarrow \text{U}(n) \rightarrow \text{PU}(n)$. The formulation is then:

Principle 1 (Copenhagen, structural form). *(i) Local validity.* Every single context is exactly classical: the restriction of the canonical cocycle to any one context trivializes, with an explicit trivializing cochain. *(ii) Complementarity.* The cocycle does not trivialize globally; its holonomy along closed context loops is a quantized obstruction, and this obstruction is the all-versus-nothing sector of contextuality, with the state sector governed by an enumerable facet geometry. *(iii) Collapse.* Statistics, repeatability, and covariance under the measured observable’s commutant pin the measurement update to a one-parameter family per degenerate block, with the Lüders rule its unique efficient member; classically the same axioms force Bayesian conditioning up to the same per-cell parameter. *(iv) Relative facts.* Value assignments exist only per context; the obstruction persists under enlargement of the observer. *(v) The movable cut.* The placement of the system/apparatus boundary is a choice of chart; section-fixed comparisons of differently-cut descriptions agree.

Throughout, “verified” means: reproduced by the named script in the companion repository [5], with the expected one-line output pinned in `verification/INDEX.md`. The supporting mathematics is developed in the companion papers [1, 2, 3, 4]; this note is the synthesis; proofs live in the companion papers and the verification supplement, with the source of every numbered claim listed in Appendix A.

2 What changed since v1

Version v1 (February 2026) contained four errors of substance, found by adversarial review and machine checking. We state them because the credibility of a structural program rests on its error ledger.

1. *Wrong groupoid.* v1 built the cocycle on $\text{U}(n) \ltimes X$. The lifting class there is vacuous (Theorem 1 of [1]; `thmG_general.py`); the canonical class lives on $\text{PU}(n) \ltimes X$, where it has order exactly n [1, Thm. 8].
2. *Overclaim “encodes”.* v1 asserted that holonomy *encodes* contextuality. False in both directions: at $d = 3$ the class is nontrivial of order 3 while the achievable state-independent all-versus-nothing value is 0 (`d3_gap_certificate.py`), and KCBS contextuality is invisible to the class (`kcbs_converse.py`). The correct statement is *detects the AvN sector* (Theorem 3 below).
3. *Level-unspecified Wigner–friend holonomy.* v1 attributed $-i$ to “the” holonomy of the friend loop. The projective (ray-level) holonomy is $+1$; $-i$ is the determinant-*section* holonomy, with section-independent square -1 (`wf_loop_holonomy.py`). Claims must pin the level.
4. *Efficiency-restricted Lüders uniqueness.* v1’s uniqueness argument for the update rule quantified only over efficient (single-Kraus) instruments. The instrument-level theorem without that restriction is Theorem 6 below.

3 Local validity is exact

Theorem 1 (Local classicality). *Let C be any single Weyl context at local dimension d . The restriction of the canonical cocycle to C (the Weyl cocycle on the context's label group) is a coboundary over $U(1)$ (the restricted class is symmetric, and $\text{Ext}(G, U(1)) = 0$ for the divisible group $U(1)$), with an explicit trivializing cochain λ ; in every certified instance ($d = 2, 3, 4$ families, V23; abelian two-generator groups at $d = 6, 8, 12$, V45) λ takes values in the cyclotomic group μ_{4d} , and the half-steps $\tau = e^{i\pi/d} \notin \mu_d$ are in general necessary. (The μ_{4d} valuation in full generality is open; the obvious closed-form candidate fails at $d = 3, 4$ — pinned in V45.)*

Verified with explicit λ per context in `local_classicality.py` (V23): all rows and columns of the Peres–Mermin square at $d = 2$, a $d = 3$ sample family, and the certified $d = 4$ family; the same script exhibits the global contrast (no joint λ exists across contexts where the obstruction is present, cross-referencing V10/V13/V33). This is tenet (i) of Principle 1 in exact form: every chart of the atlas is flat, with the flattening written down.

4 The global twist: what holonomy detects

Theorem 2 (Canonical class; [1]). *On $U(n) \ltimes X$ the $U(1)$ lifting class vanishes identically. After Morita reduction, the canonical class on $PU(n) \ltimes X$ has order exactly n .*

Theorem 3 (Detection equivalence). *Let F be a closed-triple Weyl context family at even d , with incidence matrix A and context-phase vector s (in an integral gauge, which always exists). The following are equivalent: (a) no noncontextual value assignment over $\mathbb{Z}_d$ exists; (b) some left-kernel vector u ($uA \equiv 0 \pmod{d}$) has $u \cdot s \equiv d/2 \pmod{d}$; (c) some certificate carries odd commutator carry [2, Thm. 4]. The achievable state-independent values are exactly $\{0, d/2\}$ for even d and $\{0\}$ for odd d [2, Thm. 1].*

Proof. (b) $\Rightarrow$ (a): a solution γ would give $u \cdot s \equiv (uA)\gamma \equiv 0$. (a) $\Rightarrow$ (b): by Smith normal form over $\mathbb{Z}$, unsolvability of $A\gamma \equiv s \pmod{d}$ produces, constructively, $u = (d/\gcd(g_i, d))U_i$ with $uA \equiv 0$ and $u \cdot s \not\equiv 0 \pmod{d}$ (double annihilation over $\mathbb{Z}_d$); such a u is a certificate of the multiplicity covered by [2, Rem. 3], so by the Obstruction Spectrum Theorem [2, Thm. 1] its value satisfies $2(u \cdot s) \equiv 0$, forcing $u \cdot s \equiv d/2$. (b) $\Leftrightarrow$ (c) is [2, Thm. 4]. Finally, an integral gauge exists *constructively*, and canonically: telescoping the Weyl product law $c(u, v) = -q(u) - q(v) + 2\beta(u, v) + q(u + v)$ around any closed context gives the q -parity identity $s_\tau \equiv A\bar{q} \pmod{2}$, $\bar{q}_v = q(v) \pmod{2}$, so $a = \bar{q}$ solves the GF(2) gauge system explicitly and $s = (s^{(2d)} - A\bar{q})/2 \pmod{d}$ is a canonical integral phase vector built from the gauge-free τ -level data $s^{(2d)}$; kernel pairings are gauge-invariant. This is the τ -general form of the equivalence: no gauge choice enters the hypotheses. Certified end-to-end, with random and pinned positives, in `solvability_equivalence.py` (V44) and `tau_gauge_general.py` (V45, including the $d = 6$ extension). $\square$

What was, in v2, a machine-verified converse (2,100 families, zero mismatches) is thus a corollary of [2]’s proven theorems plus five lines of ring theory — the gap was assembly, not mathematics. Both containments of the naive slogan fail, and strictly (`d3_gap_certificate.py`, `kcbs_converse.py`): the class does not imply operational contextuality, and state-dependent contextuality can be class-invisible. Complementarity, tenet (ii), is therefore *located*: it is the AvN obstruction, no more and no less at the level of the canonical class.

4.1 The state sector: a facet geometry, now enumerated

State-dependent contextuality is quantified by the contextual fraction CF. The following results replace v1’s vague appeal to “holonomy of the state” with exact geometry.

Theorem 4 (Codeword-polytope classification, $d = 2$; V31/V32). *For closed-triple Weyl families at $d = 2$, $\text{CF}(F, \rho) = 0$ if and only if the correlation vector $c(\rho)$ lies in the convex hull of the codeword set $L(F)$; quantitatively $\text{CF} = \max(0, (S^* - 1)/2)$, where S^* is the optimal facet value, proved end-to-end via a completion lemma and Abramsky–Barbosa–Mansfield duality.*

Proposition 1 (On-hull). *Every quantum behavior of the certified five-context $d = 4$ family of Theorem 5 satisfies three families of exact linear identities: per-context normalization, shared-observable marginal consistency, and vanishing on spectrally forbidden joint outcomes (24 constraints). The affine space they cut out has dimension exactly 33, equal to the affine span of the 64 deterministic behaviors, which satisfy the same identities. Hence quantum moment vectors lie in the deterministic affine hull.*

Theorem 5 (Complete facet census, certified $d = 4$ family; V43). *For the certified five-context $d = 4$ family with one context deleted, the noncontextual polytope has exactly 23,256 facets in 61 orbit classes under the derived vertex-symmetry group of order 768. By Proposition 1, $\text{CF}(\rho) > 0$ if and only if $\mu(\rho)$ violates one of the 23,256 inequalities.*

The census was derived twice by independent methods — an adjacency decomposition on the facet–ridge graph with every pivot certified in exact integer arithmetic, and a direct exact enumeration by NORMALIZ — which agree class-for-class (`d4_facet_census.py`). Structural laws of this geometry are proved in [3]: the ghost–tower law $\gamma - s^* \equiv d/2$ for context deletions, and the necessity of the τ -level (half-step) data for optimal separation (`d4_moment_facets.py`).

Conjecture 1 (Scenario-paired classification). *For every closed-triple Weyl family at even d , $\text{CF}(F, \rho) = 0$ iff $c(\rho) \in \text{conv } L(F)$ in the scenario-paired formulation of [3].*

The global-frame formulations of this statement are refuted (V28–V30); the pairing is forced, not optional.

Conjecture 2 (Tower). *Loop holonomies of Weyl families at local dimension d take values in μ_{2d} ; the 2-adic tower of [2] is exhaustive.*

5 Collapse: the unique bookkeeping, and its classical twin

Theorem 6 (Instrument-level Lüders; V37). *Let $\{\mathcal{I}_j\}$ be an instrument for a projective measurement with degenerate blocks, satisfying (i) statistics, (ii) repeatability, and (iii) covariance under the commutant of the measured observable. Then each $\mathcal{I}_j$ is exactly $p_j \cdot (\text{Lüders}) + (1 - p_j) \cdot (\text{block depolarization})$; the solution space has exactly one parameter per degenerate block (nullity 2 at $(4, [2, 2])$, verified also at $(6, [3, 2, 1])$), and the Lüders instrument is its unique efficient (single-Kraus) member. Covariance under the MASA alone is strictly insufficient. Proof via Schur’s lemma on block-covariant channels, in the v2 supplement (`note_v2_patch.md`); machine-exact certificates in V37.*

Theorem 7 (Sharp core; V40). *For a POVM instrument, statistics plus repeatability force every output of $\mathcal{I}_j$ into the eigenvalue-1 subspace of its effect E_j , and these subspaces are mutually*

orthogonal. A strictly unsharp effect (no 1-eigenvector) admits no repeatable instrument at all. The folk claim “repeatable $\Rightarrow$ projective” is nevertheless false as stated: a pinned counterexample with effect $\text{diag}(1, c)$, $0 < c < 1$, is repeatable and non-projective.

Theorem 8 (Classical twin; V40). *The same axioms, interpreted for classical stochastic instruments on a finite outcome partition, force $t \cdot (\text{Bayesian conditioning}) + (1-t) \cdot (\text{uniformization per cell})$, with nullities matching the quantum case block-for-block.*

Remark 1 (Position). Theorems 6–8 give the Leifer–Spekkens thesis a sharpened form: collapse is not a dynamical anomaly but the *unique* bookkeeping compatible with the axioms, and its classical shadow under the same axioms is Bayes’ rule. What is quantum is not the updating; it is the geometry (§4) on which the updating lives.

6 Observers, relative facts, and the cut

The observer enters through the observer-context groupoid of [4], whose objects are pairs (context, observer chain) and whose fibered morphisms are switches within a fixed enlargement and refinement isometries.

Theorem 9 (Relative facts; [4]). *Value assignments exist per object and pull back monotonically along fibered morphisms; the global obstruction persists under Wigner enlargement — for the Peres–Mermin square tensored with a friend ($d = 8$), the context signs remain $(+1)^5(-1)^1$ (`observer_groupoid.py`, certificate A).*

Theorem 10 ($\mathbb{Z}_2$ switching invariant; [4]). *The square of the switch-loop holonomy is a gauge-independent $\mathbb{Z}_2$ invariant (proved via Weyl determinants ± 1): -1 on the mutually-unbiased-bases triangle, $+1$ on within-context Weyl loops. The converse fails: a complementary 2-cycle with value $+1$ is pinned.*

Theorem 11 (Movable cut; [4]). *Section-fixed comparisons of descriptions differing by cut placement agree, with the update rule of Theorem 6 forced on both sides.*

Theorem 12 (Frauchiger–Renner localized; [4]). *In the observer-context groupoid, the FR cycle carries trivial class: all switch-loop phases are $+1$, fully gauge-invariantly. The paradox localizes to an ordinary state-dependent violation: $\text{CF} = 1/6$ exactly, attained on the single CHSH facet $\langle \bar{Z}Z \rangle - \langle \bar{Z}X \rangle + \langle \bar{X}X \rangle + \langle \bar{X}Z \rangle \leq 2$ at quantum value $7/3$ (`fr_cycle.py`).*

Theorem 12 *refutes* v1’s implicit suggestion (and [4]’s own initial conjecture) that the FR argument is a holonomy phenomenon: the switch holonomy is trivial; what FR exercises is the state sector of §4.1, with definite coordinates and a definite price.

The Wigner–friend loop, level-pinned. For the loop $T \rightarrow T_x \rightarrow T_y \rightarrow T$ generated by $\mathbb{K} \otimes H, \mathbb{K} \otimes S$: the ray-level (Pancharatnam) holonomy is $+1$; the determinant-section holonomy is exactly $-i$, a τ -level invariant with section-independent square -1 (`wf_loop_holonomy.py`).

Hardware. On `ibm_fez` (job `d986q62f47jc73a7hm2g`): the interferometric loop phase is $-2.93^\circ \pm 1.55^\circ$, consistent with ray-level $+1$ and excluding $-i$ at $\sim 56\sigma$; on the same device the Peres–Mermin witness attains $S = 4.6125 \pm 0.0173$, i.e. 35.4σ above the tight deterministic bound 4 (`hardware/results/ibm_fez_holonomy_20260709.json`). One machine, both signatures, both sharp.

7 The principle, assembled

Principle 1 now reads as a theorem schedule. (i) is Theorem 1. (ii) is Theorems 2–3 for the state-independent sector and Theorems 4–5 for the state sector, with Conjectures 1–2 marking the frontier. (iii) is Theorems 6–8. (iv) is Theorems 9–10 and the localization Theorem 12. (v) is Theorem 11. Nothing here modifies quantum predictions; the claim is that the Copenhagen commitments, correctly stated, are structure rather than stance — local flatness, one quantized global twist, forced bookkeeping, chart-relative facts, and chart-choice for the cut.

Open Problem 1. (a) *The cohomological home of class-invisible (KCBS-type) contextuality.* (b) *The general-d facet census and the arithmetic profile of the 61 classes.* (c) *Conjectures 1 and 2.*

A Claims-to-verification table

Claim	Source	Verification
Thm. 1	this note	local_classicality.py (V23)
Thm. 2	[1] Thm. 1, 8	thmG_general.py
Thm. 3	this note; [2] Thm. 1, 4	solvability_equivalence.py (V44); tau_gauge_general.py (V45)
gaps (both strict)	[3]	d3_gap_certificate.py, kcbs_converse.py
Thm. 4	[3] Thm. 2	paired_frame_construction.py, ghost_facet_theorem.py
Thm. 5	this note; [3]	d4_facet_census.py (V43)
tower laws	[3]	d4_ghost_tower.py, d4_moment_facets.py
Thm. 6	this note	lueders_instrument.py (V37)
Thm. 7, 8	this note	s4_povm_bayes.py (V40)
Thm. 9, 10	[4]	observer_groupoid.py (V38)
Thm. 11	[4]	V37 + [4] §3
Thm. 12	[4]	fr_cycle.py (V39)
WF level split	[3]	wf_loop_holonomy.py
Prop. 1	this note	d4_facet_census.py (exact-rank lemma)
S4/S5 proofs	v2 supplement	note_v2_patch.md
hardware	—	ibm_fez job d986q62f47jc73a7hm2g

References

- [1] M. Flores Gordillo, *The Canonical Class of the MASA Context Stack and the Peres–Mermin Obstruction*, working draft, 2026.
- [2] M. Flores Gordillo, *The Obstruction Spectrum of Weyl Context Families and the 2-adic Tower*, working draft, 2026.
- [3] M. Flores Gordillo, *What Contextual Holonomy Detects*, working draft, 2026.
- [4] M. Flores Gordillo, *The Observer-Context Groupoid: Relative Facts, a $\mathbb{Z}_2$ Switching Invariant, and the Movable Cut*, working draft v0.3, 2026.
- [5] github.com/manuflog/contextuality-obstructions, verification suite, 2026.